

Genome-wide analyses for personality traits identify six genomic loci and show correlations with psychiatric disorders

Min-Tzu Lo¹, David A Hinds², Joyce Y Tung², Carol Franz³, Chun-Chieh Fan^{1,4}, Yunpeng Wang⁵⁻⁷, Olav B Smeland^{6,7}, Andrew Schork^{1,4}, Dominic Holland⁵, Karolina Kauppi^{1,8}, Nilotpal Sanyal¹, Valentina Escott-Price⁹, Daniel J Smith¹⁰, Michael O'Donovan⁹, Hreinn Stefansson¹¹, Gyda Bjornsdottir¹¹, Thorgerir E Thorgerirsson¹¹, Kari Stefansson¹¹, Linda K McEvoy¹, Anders M Dale^{1,3,5}, Ole A Andreassen^{6,7} & Chi-Hua Chen¹

Personality is influenced by genetic and environmental factors¹ and associated with mental health. However, the underlying genetic determinants are largely unknown. We identified six genetic loci, including five novel loci^{2,3}, significantly associated with personality traits in a meta-analysis of genome-wide association studies ($N = 123,132$ – $260,861$). Of these genome-wide significant loci, extraversion was associated with variants in *WSCD2* and near *PCDH15*, and neuroticism with variants on chromosome 8p23.1 and in *L3MBTL2*. We performed a principal component analysis to extract major dimensions underlying genetic variations among five personality traits and six psychiatric disorders ($N = 5,422$ – $18,759$). The first genetic dimension separated personality traits and psychiatric disorders, except that neuroticism and openness to experience were clustered with the disorders. High genetic correlations were found between extraversion and attention-deficit-hyperactivity disorder (ADHD) and between openness and schizophrenia and bipolar disorder. The second genetic dimension was closely aligned with extraversion-introversion and grouped neuroticism with internalizing psychopathology (e.g., depression or anxiety).

The five-factor model (FFM) of personality, also known as the 'Big Five', is commonly used to measure individual differences in personality. It models personality according to five broad domains⁴. Extraversion (versus introversion) reflects talkativeness, assertiveness and a high activity level. Neuroticism (versus emotional stability) reflects negative affect, such as anxiety and depression. Agreeableness (versus antagonism) measures cooperativeness and compassion. Conscientiousness (versus undependability) indicates diligence and self-discipline. Openness to experience (versus being closed to experience)

captures intellectual curiosity and creativity^{4,5}. Personality phenotypes, measured by various questionnaires, are represented by continuous quantitative scores for each of the five traits⁴.

A meta-analysis of twin and family studies found that approximately 40% of the variance in personality could be attributed to genetic factors¹. Genome-wide association studies (GWAS) have discovered several variants associated with FFM traits⁶⁻⁸. Neuroticism was reported to be associated with an intronic variant in *MAG11* ($P = 9.26 \times 10^{-9}$, $N = 63,661$)⁷, conscientiousness with an intronic variant in *KATNAL2* ($P = 4.9 \times 10^{-8}$, $N = 17,375$)⁶, and openness with variants near *RASA1* ($P = 2.8 \times 10^{-8}$, $N = 17,375$)⁶ and *PTPRD* ($P = 1.67 \times 10^{-8}$, $N = 1,089$)⁸. Additionally, recent UK Biobank studies ($N = 106,716$ – $170,908$) yielded several SNPs associated with neuroticism^{2,3}.

Information collected by the consumer genomics company 23andMe contains well-phenotyped data on personality, as all participants were evaluated with the same personality inventory (Online Methods). Thus, the 23andMe data offer an opportunity to identify additional genetic variants. We performed a meta-analysis based on GWAS summary statistics to identify genetic variants associated with FFM traits. We included participants with European ancestry from 23andMe ($N = 59,225$) and two samples (GPC-1 and GPC-2) from the Genetics of Personality Consortium (GPC)^{6,7}. GPC-1 ($N = 17,375$)⁶ contains data on agreeableness, conscientiousness and openness, whereas GPC-2 ($N = 63,661$)⁷ contains information on extraversion and neuroticism.

Summary statistics of GWAS from 23andMe (**Supplementary Data Sets 1–5**) were combined with the two GPC samples separately, yielding totals of 76,600 and 122,886 subjects for the discovery-stage 1 sample. Eight linkage disequilibrium (LD)-independent SNPs ($LD r^2 < 0.05$) exceeded genome-wide significance ($P < 5 \times 10^{-8}$) in the discovery meta-analysis (**Table 1** and **Fig. 1**).

¹Department of Radiology, University of California, San Diego, La Jolla, California, USA. ²23andMe, Inc., Mountain View, California, USA. ³Department of Psychiatry, University of California, San Diego, La Jolla, California, USA. ⁴Department of Cognitive Science, University of California, San Diego, La Jolla, California, USA. ⁵Department of Neurosciences, University of California, San Diego, La Jolla, California, USA. ⁶NORMENT, KG Jebsen Centre for Psychosis Research, Institute of Clinical Medicine, University of Oslo, Oslo, Norway. ⁷Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway. ⁸Department of Radiation Sciences, Umea University, Sweden. ⁹MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University, Cardiff, UK. ¹⁰Institute of Health and Wellbeing, University of Glasgow, Glasgow, UK. ¹¹deCODE Genetics/Amgen, Reykjavik, Iceland. Correspondence should be addressed to C.-H.C. (chc101@ucsd.edu).

Received 22 July; accepted 2 November; published online 5 December 2016; doi:10.1038/ng.3736

Table 1 LD-independent genetic variants significantly associated with personality traits

SNP	Chr.	Closest gene (region)	A1/A2	Frq.	Discovery-stage 1						Replication-stage 2						Combined analysis							
					23andMe (N = 59,200)			GPC (N = 17,375 and 63,661) ^b			Combined analysis			23andMe replication (N = 39,500)			deCODE (N = 7,100)			UK Biobank (N = 91,370)		P	N	R ² (%)
					β	SE	P	β	SE	P	N	β	SE	P	β	SE	P	β	SE	P				
Conscientiousness																								
rs3814424	5q	<i>LINC00461</i> ^a	T/C	0.17	-0.289	0.050	9.75 × 10 ⁻⁹	-0.138	0.131	0.294	2.98 × 10 ⁻⁸	76.551	-0.051	0.051	0.313	-0.005	0.027	0.855	6.19 × 10 ⁻⁷	123,132	0.0202			
Extraversion																								
rs57590327	3p	<i>GRE1</i> (intergenic)	T/G	0.26	0.236	0.054	1.37 × 10 ⁻⁵	0.026	0.006	2.03 × 10 ⁻⁵	1.61 × 10 ⁻⁹	122,886	0.088	0.052	0.091	0.007	0.019	0.713	1.26 × 10 ⁻⁹	169,507	0.0217			
rs2164273	8p	<i>MTMR9</i> (intron)	G/A	0.39	0.179	0.047	1.14 × 10 ⁻⁴	0.024	0.006	4.08 × 10 ⁻⁵	1.79 × 10 ⁻⁸	122,845	0.093	0.045	0.037	0.021	0.018	0.255	1.61 × 10 ⁻⁹	169,466	0.0215			
rs6481128	10q	<i>PCDH15</i> (intergenic)	G/A	0.45	0.205	0.046	7.10 × 10 ⁻⁶	0.018	0.005	0.0010	4.15 × 10 ⁻⁸	122,886	0.154	0.045	5.58 × 10 ⁻⁴	-0.011	0.017	0.528	5.44 × 10 ⁻¹⁰	169,507	0.0227			
rs1426371	12q	<i>WSCD2</i> (intron)	A/G	0.28	-0.308	0.053	4.65 × 10 ⁻⁹	-0.023	0.006	2.56 × 10 ⁻⁴	2.09 × 10 ⁻¹¹	122,886	-0.177	0.051	5.09 × 10 ⁻⁴	-0.037	0.021	0.077	9.54 × 10 ⁻¹⁵	169,507	0.0354			
rs7498702	16p	<i>RBFox1</i> (intron)	C/T	0.29	-0.166	0.050	8.94 × 10 ⁻⁴	-0.026	0.006	1.17 × 10 ⁻⁵	4.73 × 10 ⁻⁸	122,886	-0.006	0.048	0.907	-0.005	0.018	0.777	1.89 × 10 ⁻⁶	169,507	0.0134			
Neuroticism																								
rs981523	8p	<i>XKR6</i> (intergenic)	T/C	0.50	0.250	0.042	2.68 × 10 ⁻⁹	0.022	0.006	1.01 × 10 ⁻⁴	4.25 × 10 ⁻¹²	122,867	0.138	0.042	1.05 × 10 ⁻³	0.032	0.018	0.070	1.04 × 10 ⁻¹⁰	260,861	0.0395			
rs9611519	22q	<i>L3MBTL2</i> (exon) <i>CHADL</i> (intron)	T/C	0.31	0.235	0.046	4.05 × 10 ⁻⁷	0.020	0.007	0.003	1.87 × 10 ⁻⁸	122,867	0.002	0.047	0.966	-0.002	0.023	0.931	0.0015 ^c	9.16 × 10 ⁻⁹	260,861	0.0127		

A1, effect allele; A2, non-effect allele; frq., allele frequency of A1; β , linear regression association coefficient; SE, standard error; N, sample size; β and SE may have varying scales in different cohorts; thus sample-based meta-analyses were used.

^aSNP in non-protein coding region. ^bThe sample sizes of GPC1 and GPC2 are 17,375 and 63,661, respectively. ^cOwing to absence of rs9611519 in the UK Biobank data, a proxy SNP (rs2273085, LD $r^2 = 0.99$) was used.

To evaluate the consistency of association signals between 23andMe and GPC samples, we conducted genome-wide polygenic analyses using LD Score regression to examine genetic correlations (r_g) (ref. 9) of personality traits between the two samples. The estimated r_g values were highly significant ($r_g = 0.86-0.96$), suggesting that genetic effects are consistent and replicable between the samples at the polygenic level (**Supplementary Fig. 1**) and that a considerable number of SNPs below the GWAS significance threshold contain trait-associated genetic effects.

To assess replicability of the eight significant SNPs identified in the discovery-stage 1 sample, we obtained their summary statistics from three independent samples, including an independent 23andMe replication sample, UK Biobank cohort (neuroticism only) and an Icelandic sample from deCODE Genetics (Online Methods and **Table 1**). In the final combined meta-analysis, six SNPs remained GWAS significant. The other two fell just below GWAS significance but had consistent direction of effects in all samples, suggesting that these may be significant in larger samples. Overall, the directions of effects were consistent for all eight SNPs between the discovery and replication tests, except two SNPs in the smaller ($N = 7,137$) deCODE sample.

The strongest associations were detected for neuroticism within a subregion of 8p23.1, which spans ~4 Mb (chr. 8: 8091701-11835712) with highly correlated SNPs in one LD block (**Fig. 2a**). The 8p23.1 region comprises genes related to innate immunity and the nervous system and is considered as a potential hub for cancer and developmental neuropsychiatric disorders¹⁰. Our conditional analysis indicated the presence of multiple associations (conditional $P \sim 10^{-7}$) independent of the top SNP within the 8p23.1 locus, but these were not GWAS significant.

The UK Biobank studies also identified multiple associations with neuroticism in 8p23.1 (refs. 2,3), which were attributed to an inversion polymorphism². Our association signals reside in the same inversion region, with an LD of $r^2 = 0.35$ (LDlink) between the lead SNP found here and that found in the UK Biobank study³. Additionally, we identified an intronic variant of *MTMR9* within 8p23.1 that was associated with extraversion and inversely associated with neuroticism (**Fig. 2b**). Together, these findings provide converging evidence for the association of 8p23.1 with personality.

For extraversion, we found a significant locus on 12q23.3 within *WSCD2*. This locus has been implicated in a GWAS of temperament in bipolar disorder¹¹ and in a linkage analysis¹², suggesting that 12q harbors important alleles for temperament and personality. Another SNP significantly associated with extraversion is near *PCDH15*, which encodes a member of the cadherin superfamily important for calcium-dependent cell-cell adhesion.

All six SNPs discovered here reside in loci for which genome-wide significant associations with other phenotypes have been reported (US National Human Genome Research Institute GWAS catalog). For example, we found a variant associated with neuroticism in *L3MBTL2*, a gene reported to be associated with schizophrenia¹³. Etiologically, neuroticism has been associated with schizophrenia risk¹⁴. Further, *MTMR9*, in which we found a variant associated with extraversion, has been related to response to antipsychotic medications¹⁵. The SNP associated with conscientiousness in the discovery sample, though not significant in the final meta-analysis, was located in a locus linked to educational attainment¹⁶, and high conscientiousness was found to correlate positively with academic performance¹⁷.

These six SNPs were significantly associated with gene expression, and all are listed as expression quantitative trait loci (eQTL) for brain tissues (**Supplementary Table 1**). We performed a Bayesian test¹⁸




Figure 1 Manhattan plots for personality traits in the combined sample of 23andMe and GPC data (discovery-stage1 sample). Sample sizes were as follows: agreeableness, $N = 76,551$; conscientiousness, $N = 76,551$; extraversion, $N = 122,886$; neuroticism, $N = 122,867$; openness, $N = 76,581$. Number of SNPs: agreeableness, $N = 2,165,398$; conscientiousness, $N = 2,166,809$; extraversion, $N = 6,343,667$; neuroticism, $N = 6,337,541$; openness, $N = 2,167,320$.




Figure 2 Regional association plot. (a,b) Distribution of $-\log_{10}(P)$ of SNPs on chr. 8p of the significant SNPs for neuroticism (a) and extraversion (b, top) in the combined discovery analysis. The most significant SNPs (rs6981523 and rs2164273) are shown in purple; otherwise, the colors of the circles denote their correlations (LD r^2) with the top SNP. These SNPs (LD $r^2 = 0.5$ in LDlink) have opposite β signs in GWAS results for neuroticism and extraversion. The opposite signals might be attributable to negative phenotypic association between neuroticism and extraversion. Gene symbols and locations within the region derived from UCSC Genome Browser human hg19 assembly are shown (b, bottom). Regional plots with detailed annotation information for significant SNPs are also shown in **Supplementary Figure 4**.

to examine whether GWAS signals colocalize with eQTL. COLOC-estimated posterior probabilities¹⁸ (Online Methods) indicated that one SNP-associated locus (rs57590327) and its corresponding eQTL (**Supplementary Table 1**) were probably attributable to a common causal variant (posterior probability = 0.76). Another SNP (rs216273) showed evidence of independence with eQTL (posterior probability = 0.75). For the rest of the SNPs, the posterior probability ranged between 0 and 0.45, failing to support any of the specified hypotheses. Our analyses did not show consistent evidence for these SNPs influencing personality traits through gene expression in the brain, but cautious interpretation is warranted owing to the small eQTL sample ($N = 134$).

Beyond identifying single genetic variants that each account for very little phenotypic variance, we estimated SNP-based heritability of the traits. All heritability estimates were significant in the 23andMe discovery sample, with the largest estimate for extraversion ($H^2 = 0.18$) (**Supplementary Table 2**). These findings extend those from a previous heritability analysis of FFM traits ($N = 5,011$), in which SNP-based heritability estimates were significant for neuroticism and openness¹⁹. As expected, SNP-based heritability estimates were lower than those reported in family studies¹.

Relationships among personality traits are also of interest. Although the FFM traits were derived through factor analysis and were thus orthogonal in the original findings, most studies observe some degree of phenotypic correlation between traits¹⁹. Using 23andMe data, we found that neuroticism was inversely correlated with the other personality traits, whereas agreeableness, conscientiousness, extraversion and openness were all positively correlated; all phenotypic correlations were highly significant except that between openness and conscientiousness (**Supplementary Table 3**). Genetic correlation

patterns were congruent with phenotypic correlations, but the associations were more apparent in genetic structure, which reflected shared genetic factors contributing to the correlations (Fig. 3a).

A notable feature of personality is its link with a wide range of social, mental and physical health outcomes⁵. High levels of neuroticism, extraversion and openness have been associated with bipolar disorder²⁰, and high neuroticism has been associated with major depression and anxiety²¹. Low agreeableness has been associated with narcissism, Machiavellianism and psychopathy²². In addition to phenotypic relationships, twin and GWAS studies have demonstrated genetic correlations between personality traits and psychiatric disorders^{3,21,23}, though most focus on neuroticism (Supplementary Note).

We thus sought to quantify the genetic correlations between the five personality traits and six psychiatric disorders from the Psychiatric Genomics Consortium (PGC): schizophrenia ($N = 17,115$), bipolar disorder ($N = 16,731$), major depressive disorder ($N = 18,759$), ADHD ($N = 5,422$) and autism spectrum disorder ($N = 10,263$), and from the Genetic Consortium for Anorexia Nervosa ($N = 17,767$) (Online Methods and Supplementary Table 2). A pairwise genetic correlation matrix (11×11) revealed several significant correlations (Fig. 3a and Supplementary Table 4). For example, neuroticism was highly correlated with depression, and extraversion with ADHD. To complement genetic correlation estimation via LD Score regression⁹, we compared the pattern of GWAS results by assessing whether signs of genetic effects were concordant between the top associations among these traits and disorders. The results of the sign tests of directional effects closely matched the genetic correlations (Supplementary Fig. 2).

Given the moderate and high genetic correlations, we subsequently conducted a principal component analysis (PCA) to extract principal components of genetic variation (Fig. 3b). We projected all phenotypes onto a two-dimensional space spanned by the top two principal components (PC1 and PC2) of genetic variation to summarize the genetic relationships between personality traits and psychiatric disorders. The analysis integrates genomic information with traditionally defined phenotypes to better understand basic dimensions of the full range of human behavior, from typical to pathological, in line with the research strategy of the Research Domain Criteria (RDoC)²⁴.

Our results indicated that openness, bipolar disorder and schizophrenia cluster in the first quadrant (Fig. 3b). Notably, all three share phenotypic commonality in that they have been linked to heightened creativity and dopamine activity^{25,26}. Most personality traits (conscientiousness, agreeableness and extraversion) clustered in the second quadrant. Neuroticism and depression were in the fourth quadrant. Autism and anorexia nervosa were captured by factors in higher dimensions and have relatively low loadings on the first two components (as indicated by short arrows on these two dimensions in Fig. 3b). Notably, ADHD showed a high genetic correlation with extraversion and low correlations with other psychiatric disorders (except bipolar disorder), as also shown in hierarchical clustering analysis, in which ADHD clustered with personality traits rather than psychiatric disorders (Supplementary Fig. 3). This may indicate that ADHD, or some ADHD subtypes, represent a variant of extraversion. Of note, our ADHD data were from individuals ranging in age from 5 to 19 years old. Phenotypically, positive emotionality has been linked with a subgroup of children with ADHD²⁷. Future genetic studies considering ADHD heterogeneity (e.g., subtypes and differences between child and adult forms) may help characterize its diverse etiologies and relationships with personality traits.

Overall, we observed a systematic pattern, with all psychiatric disorders showing positive loadings on PC1, and agreeableness and




Figure 3 Genetic correlations between personality traits (23andMe sample) and psychiatric disorders. **(a)** Heat map illustrating genetic correlations between phenotypes. The values in the color squares correspond to genetic correlations. Asterisks denote genetic correlations significantly different from 0: * $P < 0.05$; ** $P < 0.00091$ (Bonferroni correction threshold). **(b)** Loading plot of personality traits and psychiatric disorders on the first two principal components derived from the genetic correlation matrix in **a**. A small angle between arrows indicates a high correlation between variables, and arrows pointing in opposite directions indicate a negative correlation in the space of the two principal components.

conscientiousness with negative loadings. A combination of low agreeableness and low conscientiousness is thought to reflect Eysenck's psychoticism trait⁴. PC2 was closely aligned with the extraversion-introversion axis. Extraversion has been associated with externalizing traits and behavioral activation, and introversion, with internalizing traits and behavioral inhibition^{28,29}. Internalizing traits (e.g., neuroticism, depression, anxiety and withdrawal)²¹ have negative loadings on PC2. Externalizing traits are predicted by high extraversion, low agreeableness and low conscientiousness²⁹.

These findings provide additional support for shared genetic influences between personality traits and psychiatric disorders^{3,21,23} and for the idea that personality traits and psychiatric disorders exist on a continuum in phenotypic and genomic space^{5,11}. Maladaptive or extreme variants of personality may contribute to the persistence of, or vulnerability to, psychiatric disorders and comorbidity^{5,11,21,23}. Further genomic research in which categorical disease entities are viewed as variants of quantitative dimensions in a polygenic framework may help elucidate this issue³⁰.

Caveats of this study include that the sample size, although large, is underpowered to detect the majority of associated SNPs, given the conservative GWAS significance threshold. Because we used only GWAS summary statistics, we cannot estimate nonadditive genetic variance, such as dominance and epistasis, or genetic contributions from structural (e.g., inversions) or rare variants. Additionally, genetic correlations indicate the degree of shared genetic influences across traits at the genome-wide level, but other studies using different methods are needed to identify specific pleiotropic variants underlying the observed correlations.

In summary, by studying all FFM traits, we found six replicable genetic variants associated with personality, five of which are novel and one of which replicates published findings^{2,3}. We also observed that personality traits are correlated at the genetic level, with neuroticism showing an inverse association with the other traits. Other novel aspects of this study include description of the genetic correlations among five personality traits and six psychiatric disorders and depiction of their relationships through PCA. Personality traits are probably influenced by many genetic variants and gene–environment interactions. Researchers are only beginning to understand the genetics of personality and their relation to psychiatric disorders. The overall effort promises to have great relevance to public health.

URLs. LDlink, <http://analysistools.nci.nih.gov/LDlink/?tab=ldpair>; US National Human Genome Research Institute GWAS catalog, <https://www.ebi.ac.uk/gwas/>; LocusZoom, <http://locuszoom.sph.umich.edu/locuszoom/>; Braineac (UK Brain Expression Consortium), <http://www.braineac.org/>; LD Score regression, <https://github.com/bulik/ldsc>; GCTA-COJO (conditional and joint genome-wide association analysis), <http://cnsgenomics.com/software/gcta/cojo.html>; METAL, <http://csg.sph.umich.edu/abecasis/metal/>; PLINK 1.07, <http://pngu.mgh.harvard.edu/~purcell/plink/>; Ethical and Independent Review Services, <http://www.eandireview.com>.

METHODS

Methods, including statements of data availability and any associated accession codes and references, are available in the [online version of the paper](#).

Note: Any Supplementary Information and Source Data files are available in the online version of the paper.

ACKNOWLEDGMENTS

We thank the customers, research participants and employees of 23andMe for making this work possible. This project was funded by National Institute of Mental Health R01MH100351 (M.-T.L., N.S. and C.-H.C.), NARSAD Young Investigator award (C.-H.C.), South-East Norway Regional Health Authority (2016-064) (O.B.S.), and Research Council of Norway through a FRIPRO Mobility Grant, contract no. 251134 (Y.W.). The FRIPRO Mobility grant scheme (FRICON) is cofunded by the European Union's Seventh Framework Programme for research, technological development and demonstration under Marie Curie grant agreement no. 608695. D.J.S. is supported by a Lister Institute Prize fellowship. The research leading to deCODE results was supported in part by the US National Institutes of Health NIDA (R01-DA017932 and R01-DA034076) and the Innovative Medicines Initiative Joint Undertaking under grant agreement no. 115008, of which resources are composed of a European Federation of Pharmaceutical Industries and Associations (EFPIA) in-kind contribution and financial contribution from the European Union's Seventh Framework Programme (FP7/2007-2013) and EU-funded FP7-People-2011-IAPP grant PsychDPC (GA 28613) (H.S., G.B., T.E.T. and K.S.).

AUTHOR CONTRIBUTIONS

C.-H.C., M.-T.L. and O.A.A. designed the study. M.-T.L. and C.-H.C. analyzed data and wrote the manuscript. D.A.H. and J.Y.T. analyzed the 23andMe data. V.E.-P., D.J.S. and M.O. analyzed the UK Biobank data. H.S., G.B., T.E.T. and K.S.

analyzed the deCODE data. C.F., C.-C.F., Y.W., O.B.S., A.S., D.H., K.K., N.S., L.K.M., A.M.D. and O.A.A. contributed to manuscript preparation. All authors commented on and approved the manuscript.

COMPETING FINANCIAL INTERESTS

The authors declare competing financial interests: details are available in the [online version of the paper](#).

Reprints and permissions information is available online at <http://www.nature.com/reprints/index.html>.

- Vukasović, T. & Bratko, D. Heritability of personality: a meta-analysis of behavior genetic studies. *Psychol. Bull.* **141**, 769–785 (2015).
- Okbay, A. *et al.* Genetic variants associated with subjective well-being, depressive symptoms, and neuroticism identified through genome-wide analyses. *Nat. Genet.* **48**, 624–633 (2016).
- Smith, D.J. *et al.* Genome-wide analysis of over 106,000 individuals identifies 9 neuroticism-associated loci. *Mol. Psychiatry* **21**, 749–757 (2016).
- Goldberg, L.R. The structure of phenotypic personality traits. *Am. Psychol.* **48**, 26–34 (1993).
- Trull, T.J. & Widiger, T.A. Dimensional models of personality: the five-factor model and the DSM-5. *Dialogues Clin. Neurosci.* **15**, 135–146 (2013).
- de Moor, M.H. *et al.* Meta-analysis of genome-wide association studies for personality. *Mol. Psychiatry* **17**, 337–349 (2012).
- de Moor, M.H. *et al.* Meta-analysis of genome-wide association studies for neuroticism, and the polygenic association with major depressive disorder. *JAMA Psychiatry* **72**, 642–650 (2015).
- Kim, H.N. *et al.* Genome-wide association study of the five-factor model of personality in young Korean women. *J. Hum. Genet.* **58**, 667–674 (2013).
- Bulik-Sullivan, B.K. *et al.* LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).
- Tabarés-Seisdedos, R. & Rubenstein, J.L.R. Chromosome 8p as a potential hub for developmental neuropsychiatric disorders: implications for schizophrenia, autism and cancer. *Mol. Psychiatry* **14**, 563–589 (2009).
- Greenwood, T.A., Akiskal, H.S. & Akiskal, K.K. Bipolar Genome Study & Kelseo, J.R. Genome-wide association study of temperament in bipolar disorder reveals significant associations with three novel loci. *Biol. Psychiatry* **72**, 303–310 (2012).
- Green, E. *et al.* Localization of bipolar susceptibility locus by molecular genetic analysis of the chromosome 12q23-q24 region in two pedigrees with bipolar disorder and Darier's disease. *Am. J. Psychiatry* **162**, 35–42 (2005).
- Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511**, 421–427 (2014).
- Van Os, J. & Jones, P.B. Neuroticism as a risk factor for schizophrenia. *Psychol. Med.* **31**, 1129–1134 (2001).
- Åberg, K. *et al.* Genome-wide association study of movement-related adverse antipsychotic effects. *Biol. Psychiatry* **67**, 279–282 (2010).
- Rietveld, C.A. *et al.* Common genetic variants associated with cognitive performance identified using the proxy-phenotype method. *Proc. Natl. Acad. Sci. USA* **111**, 13790–13794 (2014).
- Poropat, A.E. A meta-analysis of adult-rated child personality and academic performance in primary education. *Br. J. Educ. Psychol.* **84**, 239–252 (2014).
- Giambartolomei, C. *et al.* Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet.* **10**, e1004383 (2014).
- Power, R.A. & Pluess, M. Heritability estimates of the Big Five personality traits based on common genetic variants. *Transl. Psychiatry* **5**, e604 (2015).
- Barnett, J.H. *et al.* Personality and bipolar disorder: dissecting state and trait associations between mood and personality. *Psychol. Med.* **41**, 1593–1604 (2011).
- Hettema, J.M., Neale, M.C., Myers, J.M., Prescott, C.A. & Kendler, K.S. A population-based twin study of the relationship between neuroticism and internalizing disorders. *Am. J. Psychiatry* **163**, 857–864 (2006).
- Jakobowitz, S. & Egan, V. The dark triad and normal personality traits. *Pers. Individ. Dif.* **40**, 331–339 (2006).
- Kendler, K.S. & Myers, J. The genetic and environmental relationship between major depression and the five-factor model of personality. *Psychol. Med.* **40**, 801–806 (2010).
- Insel, T. *et al.* Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *Am. J. Psychiatry* **167**, 748–751 (2010).
- DeYoung, C.G. The neuromodulator of exploration: a unifying theory of the role of dopamine in personality. *Front. Hum. Neurosci.* **7**, 762 (2013).
- Power, R.A. *et al.* Polygenic risk scores for schizophrenia and bipolar disorder predict creativity. *Nat. Neurosci.* **18**, 953–955 (2015).
- Karalunas, S.L. *et al.* Subtyping attention-deficit/hyperactivity disorder using temperament dimensions: toward biologically based nosologic criteria. *JAMA Psychiatry* **71**, 1015–1024 (2014).
- Gray, J.A. The psychophysiological basis of introversion-extraversion. *Behav. Res. Ther.* **8**, 249–266 (1970).
- Mezquita, L. *et al.* Five-factor model and internalizing and externalizing syndromes: a 5-year prospective study. *Pers. Individ. Dif.* **79**, 98–103 (2015).
- Plomin, R., Haworth, C.M. & Davis, O.S. Common disorders are quantitative traits. *Nat. Rev. Genet.* **10**, 872–878 (2009).

ONLINE METHODS

23andMe sample. The GWAS summary statistics were obtained from a subset of 23andMe participants. 23andMe uses a survey designed to collect a number of phenotypes, including the personality traits reported here, and the sample has been described previously for other phenotypes^{31,32}. We included only participants ($N = 59,225$) who showed >97% European ancestry as determined by analyzing local ancestry and comparing to three HapMap2 populations³³. Relatedness between participants was examined by a segmental identity-by-descent (IBD) method³⁴ to ensure that only unrelated individuals (sharing less than 700 cM IBD) were included in the sample. All participants included in the analyses provided informed consent and answered surveys online according to a human subject research protocol, which was reviewed and approved by Ethical and Independent Review Services, a private institutional review board accredited by the Association for the Accreditation of Human Research Protection Programs.

Additionally, we obtained independent replication results of GWAS from the 23andMe replication sample. This sample included ~39,500 participants ($N = 39,452$ for conscientiousness, 39,484 for extraversion and 39,488 for neuroticism) who met the inclusion criteria described above.

Genetics of Personality Consortium (GPC) sample. The Genetics of Personality Consortium (GPC) is a large collaboration of GWAS for personality. Summary statistics of the GPC data used in the current study included the first meta-analysis of GWAS (GPC-1)⁶ for three traits (agreeableness, conscientiousness and openness) and the second meta-analysis of GWAS (GPC-2) for neuroticism and extraversion^{7,35,36}. The results of 10 discovery cohorts for GPC-1 and 29 discovery cohorts for GPC-2 are available in the public domain, and consist of 17,375 and 63,661 participants, respectively, with European ancestry across Europe, Australia and United States. These studies were performed with oversight from local ethic committees, and all participants provided informed consent^{6,7,35,36}.

UK Biobank sample. UK Biobank is a large prospective cohort of more than 502,000 participants (aged 40–69 years)³ with genetic data and a wide range of phenotypic data, including social, cognitive, personality (neuroticism trait), life style, and physical health measures collected at baseline recruitment from 2006 to 2010. We used a subsample of this cohort for neuroticism replication. Exclusion criteria included UK Biobank genomic analysis exclusions, relatedness, gender mismatch, non-white UK ancestry and failure of quality control of UK BiLEVE genotyping³, resulting in a sample of 91,370 individuals. Association analysis was conducted using linear regression under a model of additive allelic effects with sex, age, array and the first eight PCs as covariates³. Informed consent was obtained from all participants, and the study was approved by the UK National Health Service National Research Ethics Service³.

deCODE sample. Icelandic participants ($N = 7,137$ for extraversion, 7,136 for neuroticism and 7,129 for conscientiousness) were enrolled in various ongoing deCODE studies administering the Neuroticism–Extraversion–Openness Five-Factor Inventory (NEO-FFI) measure of the Big Five personality traits^{37,38}. All deCODE studies were approved by the appropriate bioethics and data-protection authorities, and all subjects donating blood provided informed consent. The personal identities of participants from whom phenotype information and biological samples were obtained were encrypted by a third-party system overseen by the Icelandic Data Protection Authority³⁹. A generalized form of linear regression that accounts for relatedness between individuals was used to test the correlation between normalized NEO-FFI trait scores and genotypes.

Personality assessment. In the 23andMe sample, individuals completed a web-based implementation of the Big Five Inventory (BFI)^{40,41} that includes 44 questions. Scores for agreeableness, conscientiousness, extraversion, neuroticism and openness were computed using 8 to 10 items per factor⁴⁰.

In GPC-1, scores of personality traits were based on the 60-item NEO-FFI with 12 items per factor^{6,37}. In GPC-2, harmonization of measures for neuroticism and extraversion across 9 inventories and 29 cohorts was performed by applying Item Response Theory (IRT) to avoid personality scores being influenced by the number of items and the specific inventory. Because the

personality measures were not assessed similarly across GPC-2 cohorts, the harmonized or calibrated scores of personality are more comparable, thereby increasing power for meta-analysis of GWAS using fixed-effect models^{7,35,36}. As described in the main text, we found high genetic correlations between 23andMe and GPC samples, suggesting a highly consistent pattern of associations despite the discrepancy in questionnaires (**Supplementary Fig. 1**).

In the UK Biobank sample, neuroticism was scored between 0 and 12 using the 12 items of the Eysenck Personality Questionnaire, Revised Short Form (EPQ-R-S)⁴² with high reliability and concurrent validity⁴².

In the deCODE sample, NEO-FFI personality trait scores^{37,38} were adjusted for sex and age at measurement and were then normalized to a standard normal distribution using quantile normalization.

Regional association and annotation plot. The regional plot of chromosome 8p (**Fig. 2**) was constructed by a web-interface tool, LocusZoom⁴³. The bottom panel displays gene symbol and location within the region derived from UCSC Genome Browser human hg19 assembly. The regional and annotation plots for other significant SNPs are also shown in **Supplementary Figure 4**.

Distributions and correlations for personality scores in the 23andMe sample. Quantile–quantile (QQ) plots of covariate-adjusted personality scores to examine normality are shown in **Supplementary Figure 5**. The distributions at the top tail deviate from normality owing to the limited range of the scores, and those at the bottom tail deviate due to the limited range (for neuroticism and extraversion) and/or extreme values. This violation of the normality assumption can be influential for genetic variants with very low minor allele frequencies (e.g., rare variants)⁴⁴. However, this did not affect our results because our GWAS and LD Score regression⁹ include only common variants.

Pearson correlations, unadjusted and after adjusting for the covariates (age, sex and top five principal components (PCs) for population structure correction⁴⁵), were used to assess phenotypic correlations among the five traits (**Supplementary Table 3**).

Genotyping and imputation. In the 23andMe sample, DNA extraction and genotyping were performed on saliva samples by National Genetics Institute (NGI), a CLIA-licensed clinical laboratory and a subsidiary of Laboratory Corporation of America. Samples were genotyped on one of four genotyping platforms. The V1 and V2 platforms were variants of the Illumina HumanHap550+ BeadChip, including about 25,000 custom SNPs selected by 23andMe, with a total of about 560,000 SNPs. The V3 platform was based on the Illumina OmniExpress+ BeadChip, with custom content to improve the overlap with 23andMe's V2 array, with a total of about 950,000 SNPs. The 23andMe's V4 platform in current use is a fully custom array, including a lower redundancy subset of V2 and V3 SNPs with additional coverage of lower-frequency coding variation, and about 570,000 SNPs. Samples that failed to reach a 98.5% call rate were reanalyzed. As part of 23andMe standard practice, individuals whose analyses failed repeatedly were contacted and asked to provide a new sample.

23andMe participant genotype data were imputed using the 1000 Genomes Project phase 1 version 3 reference panel⁴⁶. The phasing and imputation for each genotyping platform were separated. First, chromosomal segments of no more than 10,000 genotyped SNPs, with overlaps of 200 SNPs, were phased using Beagle (version 3.3.1)⁴⁷. Then, each phased segment was imputed against all-ethnicity 1000 Genomes Project haplotypes (excluding monomorphic and singleton sites) using a high-performance version of Minimac⁴⁸ for 5 rounds and 200 states to estimate parameters. SNPs were filtered by procedures including Hardy–Weinberg equilibrium $P < 10^{-20}$ (stringent threshold for large sample size), call rate < 95% and allele frequencies apparently different from European 1000 Genomes Project reference data. A total of 13,341,935 SNPs was retained after filtering and excluding chromosome X, Y and mitochondria. We focused on autosomal SNPs, which are available for 23andMe, GPC and UK Biobank samples.

Genotyping in cohorts of GPC-1 (ref. 6) and GPC-2 (refs. 7,35) was conducted on Illumina or Affymetrix platforms. Quality control of genotype data was examined in each cohort independently, including checks for European ancestry, sex inconsistencies, Mendelian errors, high genome-wide homozygosity,

relatedness, minor allele frequencies (MAFs), SNP call rate, sample call rate and Hardy–Weinberg equilibrium^{6,7,35,36}. Genotype data of GPC-1 were then imputed using HapMap phase II CEU (Utah residents with Northern and Western European ancestry from the CEPH collection) as a reference panel including ~2.5 million SNPs⁶ and, alternatively, a reference panel from 1000 Genomes Project phase 1 version 3 was used to impute the genotype data of GPC-2 (refs. 7,35,36). Poorly imputed SNPs ($r^2 < 0.3$ or imputation quality (proper_info) < 0.3 (ref. 6) or 0.4 (refs. 7,35) and low MAF (< 0.01 (ref. 6) or $\sqrt{5/N}$ (refs. 7,35)) were excluded in the meta-analyses, resulting in a total number of 1.1 million–6.6 million SNPs^{7,35} across cohorts of GPC.

In the UK Biobank first release genetic data of 152,729 participants (June 2015), about two-thirds of the sample was genotyped using Affymetrix UK Biobank Axiom array, and the remaining were genotyped using the Affymetrix UK BiLEVE Axiom array³. Outlier, multiallelic and low-MAF ($< 1\%$) SNPs were excluded from phasing and imputation procedures. The reference panel of imputation was based on the 1000 Genomes Phase 3 and UK10K haplotype panels³. Further quality control procedures were applied after imputation, yielding a total of 8,268,322 SNPs for further analyses³.

Genotyping, imputation methods and the association analysis method used in the deCODE sample were previously described⁴⁹. A total of 676,913 autosomal SNPs were typed using Illumina SNP chips⁴⁹. SNPs with low MAF ($< 0.1\%$) and low imputation information (< 0.8) were excluded and 99.5% of SNPs remained after imputation.

Genome-wide association analysis. Association tests were performed by regressing personality traits on imputed dosages of SNPs in the 23andMe sample. Age, sex and the top five PCs⁴⁵ for population structure correction were included as covariates, and P values were computed using likelihood ratio tests. For all five personality traits, the correlation structure of SNPs was determined by an LD matrix of 9,270,523 autosomal SNPs generated from European reference sample in 1000 Genomes Project phase 1 version 3 within 1,000,000 bp (1 Mb)^{50,51} using PLINK 1.07 (ref. 52). The original 13,341,935 SNPs were reduced into 9,270,523 SNPs in our subsequent analyses (e.g., LD correlation structure is used to determine LD-independent SNPs). All SNP positions were mapped to Genome Reference Consortium Human Build 37 (GRCh37) and UCSC Genome Browser human hg19 assembly. We made QQ plots with GWAS summary statistics of the 23andMe sample. The QQ plots lie along the expected null line for large P values ($P > 10^{-3}$), indicating that the GWAS results are not inflated by population stratification or cryptic relatedness. This pattern is consistent with the genomic inflation factors (λ)⁵³ close to 1, as shown in **Supplementary Figure 6**.

In each cohort of GPC-1 (ref. 6) and GPC-2 (refs. 7,35), linear regressions with covariates of sex, age and PCs were conducted for association tests using dosage data. The meta-analyses of GWAS results of cohorts for GPC-1 and GPC-2 were performed by the inverse-variance method using METAL⁵⁴. SNPs available in one cohort only were excluded. The totals of 2,305,461, 2,305,682 and 2,305,640 SNPs were available for traits of agreeableness, conscientiousness and openness (respectively) in GPC-1, as well as 6,941,603 SNPs for extraversion and 6,949,614 SNPs for neuroticism in GPC-2. Genomic inflation factors (λ) are 1.01, 1.01, 1.03, 1.02 and 1.02 for agreeableness, conscientiousness, extraversion, neuroticism and openness, respectively.

Meta-analysis of 23andMe and GPC samples. Given improved power for detection of genetic effects with larger sample sizes in GWAS, we performed a combined meta-analysis of 23andMe and GPC samples using METAL⁵⁴ on the basis of the sample-size based method. To assess the quality of meta-analysis, SNPs with heterogeneity $P < 0.05$ were excluded. Eight significant LD-independent SNPs were identified after removing correlated SNPs at LD $r^2 > 0.05$ that are within 1 Mb of the top SNP. In **Table 1**, the percentage of variance explained by each SNP is calculated using equation: $(z^2/(n-k-1+z^2)) \times 100$, where z is the z value for each SNP controlling for covariates, n is the sample size for each SNP and k is the number of covariates in the regression model ($k = 7$ for age, sex, and top five PCs)^{55,56}.

Conditional analysis within 1-Mb region of significant SNPs. We performed a conditional analysis⁵⁷ within the 1-Mb genomic region of each of the six

LD-independent SNPs. In our study, we used 1000 Genomes Project reference panel of European ancestry to estimate LD correlations (r^2) and excluded SNPs correlated at LD $r^2 > 0.9$ with the top associated SNP within a 1-Mb window. We did not detect additional significant SNPs conditional on the top SNPs under the stringent GWAS threshold. However, for the significant loci in 8p, several SNPs still showed substantial association signals ($P \sim 10^{-7}$) conditioning on the top SNPs, rs6981523 or rs2164273.

Genetic correlation analysis. We used the LD Score regression method to examine the pattern of genetic correlations (r_g)^{9,58} across personality traits within and between 23andMe and GPC samples (**Fig. 3a**, **Supplementary Fig. 1** and **Supplementary Table 4**) on the basis of GWAS summary statistics. The LD Score for each SNP measures the amount of pairwise LD r^2 with other SNPs within 1-cM windows from 1000 Genomes Project reference panel of European ancestry. All SNPs were filtered by LD Score regression built-in procedures, including imputation quality (INFO) > 0.9 and MAF > 0.1 , and merged to SNPs in HapMap 3 reference panel. Approximately 0.8 million–1.1 million SNPs (**Supplementary Table 2**) were retained to estimate genetic correlations.

We also examined genetic correlations among the five traits, which have been estimated previously using a twin design^{59,60}, and unrelated individuals' SNP data from a relatively smaller sample, in which many estimates did not converge¹⁹. Our LD Score regression analysis based on a large sample provided additional contribution to this effort.

We further quantified genetic correlations between personality traits and psychiatric disorders, including schizophrenia⁶¹, bipolar disorder⁶², major depressive disorder⁶³, ADHD⁶¹, autism spectrum disorder⁶¹ and anorexia nervosa⁶⁴.

Query for eQTL database. We queried eQTL evidence for our significant SNPs from the Brain eQTL Almanac (Braineac)^{65,66}. The results are listed in **Supplementary Table 1**. We display the brain region with the lowest P value for each SNP among all 10 brain regions. To check the rank of eQTL P values of six LD-independent SNPs in the Braineac database, we randomly selected 50,000 SNPs and queried the database to extract the lowest P value for each SNP, resulting in a total of 36,190 SNPs with eQTL results. To match allele frequencies and distances to transcription start site (TSS) with the significant SNPs, the randomly selected SNPs were stratified into four groups: (i) within transcript, (ii) downstream 0–200 kb, (iii) upstream 0–200 kb and (iv) upstream 200–400 kb. SNPs that fell outside these ranges were removed. The SNPs in the 'within transcript' group were further stratified into three subgroups according to allele frequencies. This procedure resulted in six distributions of eQTL P values that matched the significant SNPs in terms of allele frequencies and TSS, and these were used to determine the ranking of eQTL associations (**Supplementary Tables 1** and **5**). Two SNPs were ranked highly for their significance as eQTL compared to randomly sampled eQTL markers with matched allele frequencies and distances to TTS from the Braineac database (top 10–20% ranking, rs6981523; top 20–30% ranking, rs9611519; **Supplementary Table 5**).

Colocalization analysis between GWAS and eQTL. To investigate whether GWAS-significant SNPs and their eQTLs were colocalized with a shared candidate causal variant, we performed a colocalization analysis, COLOC, that uses Bayesian posterior probability to assess colocalization¹⁸. The SNP-associated locus was defined as within a 1-Mb window¹⁸ for each of the six SNPs (**Table 1**). The prior probabilities that the locus is associated with only trait 1 (i.e., personality traits), only trait 2 (i.e., eQTL) and both are 10^{-5} , 10^{-4} and 10^{-6} , respectively. The posterior probabilities (PPO, PP1, PP2, PP3 and PP4) for five hypotheses (H_0 , no association with either trait; H_1 , association with trait 1, not with trait 2; H_2 , association with trait 2, not with trait 1; H_3 , independent association with two traits, two independent SNPs; H_4 , association with both traits, one shared SNP)¹⁸ were calculated to determine which hypothesis is supported by the data. A limitation of this analysis is the potentially low power in the small eQTL sample ($N = 134$).

SNP-concordant test for the top GWAS signals. To investigate concordance of SNP effects between personality traits and psychiatric disorders, we followed a procedure similar to one described previously^{67,68} by counting the number

of same-direction effect sizes for the LD-independent top SNPs ($P < 10^{-4}$) in the pairwise phenotype data and calculated the proportion of the same-direction effects in the total number of LD-independent top SNPs. The one-sided P value for the proportion of pairwise phenotypes was computed using a binomial test to examine the deviation from 0.5 for the proportion. In **Supplementary Figure 2**, a heat map of the proportions of the same-direction effect for pairwise phenotypes shows a similar pattern with a heat map of genetic correlations in **Figure 3a**.

Hierarchical clustering analysis. We performed hierarchical clustering analysis using dissimilarity measures (1-genetic correlation) implemented in hclust function of R to investigate and display relationships between personality traits and psychiatric disorders. On the basis of genetic correlations, the more highly correlated phenotypes were grouped in the same clusters and displayed by a dendrogram (**Supplementary Fig. 3**), showing an agreement with classifications of the loading plot (**Fig. 3b**).

Data availability. GPC-1 and GPC-2 summary statistics are available at <http://www.tweelingenregister.org/GPC/>; Psychiatric Genomics Consortium (PGC) summary statistics (schizophrenia, bipolar disorder, major depressive disorder, ADHD, autism spectrum disorder and anorexia nervosa) are available at <https://www.med.unc.edu/pgc/results-and-downloads>. The top 10,000 SNPs for five personality traits from the 23andMe discovery data set are available in **Supplementary Data Sets 1–5**. The full GWAS summary statistics for the 23andMe discovery data set will be made available through 23andMe to qualified researchers under an agreement with 23andMe that protects the privacy of the 23andMe participants. Please contact D.A.H. (dhinds@23andme.com) for more information and to apply for data access. All other data reported in the paper are included in the paper and **Supplementary Materials**.

31. Hu, Y. *et al.* GWAS of 89,283 individuals identifies genetic variants associated with self-reporting of being a morning person. *Nat. Commun.* **7**, 10448 (2016).
32. Pickrell, J.K. *et al.* Detection and interpretation of shared genetic influences on 42 human traits. *Nat. Genet.* **48**, 709–717 (2016).
33. Falush, D., Stephens, M. & Pritchard, J.K. Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* **164**, 1567–1587 (2003).
34. Henn, B.M. *et al.* Cryptic distant relatives are common in both isolated and cosmopolitan genetic samples. *PLoS One* **7**, e34267 (2012).
35. van den Berg, S.M. *et al.* Meta-analysis of genome-wide association studies for extraversion: findings from the Genetics of Personality Consortium. *Behav. Genet.* **46**, 170–182 (2016).
36. van den Berg, S.M. *et al.* Harmonization of neuroticism and extraversion phenotypes across inventories and cohorts in the Genetics of Personality Consortium: an application of Item Response Theory. *Behav. Genet.* **44**, 295–313 (2014).
37. Costa, P.T. & McCrae, R.R. *Revised NEO Personality Inventory (NEO PI-RTM) and NEO Five-Factor Inventory (NEO-FFI): Professional Manual* (Psychological Assessment Resources, 1992).
38. Bjornsdottir, G. *et al.* Psychometric properties of the Icelandic NEO-FFI in a general population sample compared to a sample recruited for a study on the genetics of addiction. *Pers. Individ. Dif.* **58**, 71–75 (2014).
39. Gulcher, J.R., Kristjánsson, K., Gudbjartsson, H. & Stefánsson, K. Protection of privacy by third-party encryption in genetic research in Iceland. *Eur. J. Hum. Genet.* **8**, 739–742 (2000).
40. John, O.P.D., E., M. & Kentle, R.L. *The Big Five Inventory—versions 4a and 5a* (University of California, Berkeley Institute of Personality and Social Research, 1991).
41. Soto, C.J. & John, O.P. Ten facet scales for the Big Five Inventory: convergence with NEO PI-R facets, self-peer agreement, and discriminant validity. *J. Res. Pers.* **43**, 84–90 (2009).
42. Eysenck, S.B.G., Eysenck, H.J. & Barrett, P. A revised version of the psychoticism scale. *Pers. Individ. Dif.* **6**, 21–29 (1985).
43. Pruim, R.J. *et al.* LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics* **26**, 2336–2337 (2010).
44. Búžková, P. Linear regression in genetic association studies. *PLoS One* **8**, e56976 (2013).
45. Lehoucq, R.B., Sorensen, D.C. & Yang, C. Society for Industrial and Applied Mathematics. *ARPACK Users' Guide: Solution of Large-scale Eigenvalue Problems with Implicitly Restarted Arnoldi Methods* (Society for Industrial and Applied Mathematics, 1998).
46. 1000 Genomes Project Consortium. *et al.* A map of human genome variation from population-scale sequencing. *Nature* **467**, 1061–1073 (2010).
47. Browning, S.R. & Browning, B.L. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *Am. J. Human Genet.* **81**, 1084–1097 (2007).
48. Howie, B., Fuchsberger, C., Stephens, M., Marchini, J. & Abecasis, G.R. Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nat. Genet.* **44**, 955–959 (2012).
49. Gudbjartsson, D.F. *et al.* Large-scale whole-genome sequencing of the Icelandic population. *Nat. Genet.* **47**, 435–444 (2015).
50. Thompson, W.K. *et al.* An empirical Bayes mixture model for effect size distributions in genome-wide association studies. *PLoS Genet.* **11**, e1005717 (2015).
51. Wang, Y. *et al.* Leveraging genomic annotations and pleiotropic enrichment for improved replication rates in schizophrenia GWAS. *PLoS Genet.* **12**, e1005803 (2016).
52. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).
53. Devlin, B. & Roeder, K. Genomic control for association studies. *Biometrics* **55**, 997–1004 (1999).
54. Willer, C.J., Li, Y. & Abecasis, G.R. METAL: fast and efficient meta-analysis of genome-wide association scans. *Bioinformatics* **26**, 2190–2191 (2010).
55. Hibar, D.P. *et al.* Common genetic variants influence human subcortical brain structures. *Nature* **520**, 224–229 (2015).
56. So, H.C., Li, M. & Sham, P.C. Uncovering the total heritability explained by all true susceptibility variants in a genome-wide association study. *Genet. Epidemiol.* **35**, 447–456 (2011).
57. Yang, J. *et al.* Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nat. Genet.* **44**, 369–375 (2012).
58. Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases and traits. *Nat. Genet.* **47**, 1236–1241 (2015).
59. Distel, M.A. *et al.* The five-factor model of personality and borderline personality disorder: a genetic analysis of comorbidity. *Biol. Psychiatry* **66**, 1131–1138 (2009).
60. Ono, Y. *et al.* Genetic structure of the five-factor model of personality in a Japanese twin population. *Keio J. Med.* **49**, 152–158 (2000).
61. Cross-Disorder Group of the Psychiatric Genomics Consortium. Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet* **381**, 1371–1379 (2013).
62. Psychiatric GWAS Consortium Bipolar Disorder Working Group. Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near *ODZ4*. *Nat. Genet.* **43**, 977–983 (2011).
63. Ripke, S. *et al.* A mega-analysis of genome-wide association studies for major depressive disorder. *Mol. Psychiatry* **18**, 497–511 (2013).
64. Boraska, V. *et al.* A genome-wide association study of anorexia nervosa. *Mol. Psychiatry* **19**, 1085–1094 (2014).
65. Ramasamy, A. *et al.* Genetic variability in the regulation of gene expression in ten regions of the human brain. *Nat. Neurosci.* **17**, 1418–1428 (2014).
66. Trabzuni, D. *et al.* Quality control parameters on a large dataset of regionally dissected human control brains for whole genome expression studies. *J. Neurochem.* **119**, 275–282 (2011).
67. Franke, B. *et al.* Genetic influences on schizophrenia and subcortical brain volumes: large-scale proof of concept. *Nat. Neurosci.* **19**, 420–431 (2016).
68. Nyholt, D.R. SECA: SNP effect concordance analysis using genome-wide association summary results. *Bioinformatics* **30**, 2086–2088 (2014).