

Figure 2

High performance liquid chromatogram of anthraquinones; conditions as in experimental.

1) Methanol; 2) dichloromethane; 3) rhein; 4) aloe emodin; 5) emodin; 6) physcion; 7) chrysophanol.

Flow rate: 2 ml/min.

Preparation of standard solutions: 0.05% W/V solutions of pure chrysophanol, physcion, emodin, aloe-emodin, and rhein in dichloromethane.

#### Acknowledgment

The authors are grateful to Mr. T. D. Turner, Head of the Pharmacognosy section, Welsh School of Pharmacy, UWIST for providing the facilities for this work.

#### References

- [1] M. Takido, Pharm. Bull. Jap. **4** (1956) 45.
- [2] J. W. Fairbairn, T. J. Betts, and V. K. Mital, J. Pharm. Pharmacol. **10** (1958) 436.
- [3] L. Kraus, Planta Med. **7** (1959) 427.
- [4] P. P. Rai, T. D. Turner, and S. L. Greensmith, J. Pharm. Pharmacol. **26** (1974) 722.
- [5] P. P. Rai and M. Shok, Chromatographia **14** (1981) 599.
- [6] K. Savonius, Farm. Aikak **81** (1972) 85.
- [7] T. Furuya, S. Shibata, and J. Iizuka, J. Chromatogr. **21** (1966) 116.
- [8] V. Castagnola, G. Pettinaro, and G. A. de Vries, Dt. Apothztg., **115** (1975) 818.
- [9] P. P. Rai, T. D. Turner, and S. A. Matlin, J. Chromatogr. **110** (1975) 401.
- [10] P. P. Rai, Lloydia **41** (1978) 114.

MS received: January 25, 1983

## Studies on the Essential Oil of *Valeriana celtica* L.

# 10394

C. Bicchi\*

Laboratorio RMN e Spettroscopie applicate alta Tossicologia, Facoltà die Farmacia, Università di Torino-Torino (Italy)

P. Sandra, M. Schelfaut, and M. Verzele

Laboratory of Organic Chemistry, State University of Ghent, Krijgslaan, 281 (S. 4), B-9000 Gent (Belgium)

### Key Words:

Gas chromatography

Capillary column

HPLC sample work-up

*Valeriana celtica* L.

Essential oil

Nepetalactone, isovaleric acid

Feline attractant

### 1 Introduction

*Valeriana celtica* L. [1-3] is a rare, wild-growing Alpine plant also known as "golden nard". It has an interesting history as a traditional household remedy against adverse effects related to the nervous system [4]. Soaps prepared from its roots show excellent nerve-strengthening properties [5].

It also protects linen and wool against insects. Moreover, *Valeriana celtica* L. exerts a strong attractant action on domestic cats (*Felix domestica*).

*Valeriana celtica* L. of the Valerianaicae with a chromosome number of  $2n = c48$  [2] or  $2n : c72, c96$  [3] is a typical Alpine plant. It grows in the Alps at altitudes between 1800 and 2800 m on flat humus-rich, non-lime containing soils, and sometimes on stony ground; the plant flowers in mid-summer (July-August). Very little is known about the composition of the essential oil of *Valeriana celtica* L. The present study deals with this aspect, in particular concerning the presence of feline attractants. The essential oil was obtained by steam distillation of the leaves of wild specimens of *Valeriana celtica* L. harvested in 1981 in Aosta Valley (Cogne) Italy at 2300 m above sea level. The leaves were collected after blooming of the plant because, according to the local inhabitants, the essential oil then possesses better properties.

A voucher has been deposited at the "Giardino botanico alpino Paradisia" Valnontey, Aosta (Italy).

The essential oil was analyzed as such by capillary gas chromatography-mass spectrometry and by the same technique after fractionation of the essential oil by straight phase HPLC facilitating identification of the components [6].

## 2 Experimental

### 2.1 Preparation of the Oil

100 g of dried leaves was pulverized and subjected to steam distillation in a modified Marcusson apparatus [7], yielding 0.22 g of a yellowish essential oil (0.22%).

### 2.2 Fractionation of the Oil

HPLC pre-separation of 10 mg of the essential oil was carried out on a silica column 25 cm × 0.46 cm packed with 5 µm spherical silica ROSiL (Alltech-Europe) installed in a Varian 5000 liquid chromatograph equipped with a Varichrom UV 50 detector operated at 220 nm. Four main fractions originating from a step elution program with 20 ml of hexane and 15 ml of methanol at a flow rate of 1 ml min<sup>-1</sup> were collected in 10 ml test tubes provided with a stopper.

### 2.3 Gas Chromatographic Analysis

Capillary gas chromatography was performed on a HTS-OV-1 column (20 m × 0.3 mm ID, df: 0.3 µm) [8] installed in a Carlo Erba 4160 equipped with a cold on-column injector. Operating conditions were: injection temperature 30°C, column temperature ballistically programmed to 50°C and then to 250°C at 4° min<sup>-1</sup>, detector FID at 300°C, carrier gas (hydrogen) flow rate 2 ml min<sup>-1</sup>. Quantitation was carried out with a Varian CDS 111 integrator.




Figure 1

Chromatogram of the total essential oil of *Valeriana celtica* L.


### 2.4 Gas Chromatography-Mass Spectrometry

GC-MS were obtained on a Finnigan 4000 system equipped with a Data General Nova 3 computer. The same column and conditions as reported for the GC analysis were used to obtain comparable results, except that helium was used as carrier gas (flow rate 2 ml min<sup>-1</sup>). The column was directly connected to the ion source via 1 m OV-1 coated fused silica tubing connected to the glass column with a polyimide seal [9].

Table 1

Identified compounds in the essential oil of *Valeriana celtica* L.

Peak no.	Compound	%[2]
1	Furfural	
2	Toluene	
3	Ethyl valerate	
4	Isovaleric acid	15.2
5	Benzaldehyde	
6	Valeric acid	1.0
7	4-Methylvaleric acid	
8	p-Cymene	
9	Ocimene	
10	Angelica lactone (t)	
11	Fenchyl alcohol (t)	
12	Fenchone	
13	Unknown	
14	Tujone	
15	Camphor	
16	Propyl isovalerate	
17	α-Terpineol	
18	Chrysanthenyl acetate	
19	Sabinyl acetate (t)	19
20	Nepetalactone	20
21	Geranyl acetate	1.3
22	Benzyl isovalerate	22
23	α-Ionone	23
24	β-Patchoulene	26
25	Sesquiterpene hydrocarbon	1.60
26	β-Elementene	26
27	Cedradiene	27
28	Sesquiterpene hydrocarbon	28
29	β-Caryophyllene	
30	β-Gurjunene	204
31	Seychellene	204
32	α-Guaiene	5.9
33	α-Patchoulene	204
34	Sesquiterpene hydrocarbon	204
35	β-Ionone	192
36	Cadinene	206
37	Alloaromadendrene	204
38	γ-Patchoulene	204
39	α-Murolene	204
40	Selina-4(14),7(11)-diene	204
41	Unknown	218
42	Sesquiterpene alcohol	220
43	Sesquiterpene alcohol	228
44	Sesquiterpene hydrocarbon	204
45	Patchoulane (t)	206
46	Sesquiterpene alcohol	220
47	Sesquiterpene alcohol	220
48	Sesquiterpene alcohol	222
49	Sesquiterpene alcohol	220
50	Unknown	218
51	Daucalene	198
52	Isodaucalene	198
53	Sesquiterpene alcohol	222
54	Patchouli alcohol	222
55	Unknown	m.w. 236
56	Unknown	218



**Figure 2**  
EI and CI ( $\text{CH}_4$ ) mass spectrum of unidentified sesquiterpenoid accounting for 6% of the essential oil of *Valeriana celtica* L.

### 3 Results and Discussion

**Figure 1** shows the chromatogram of the total essential oil. Isovaleric acid (peak 4), seychellene (peak 31), patchouli alcohol (peak 54), and an oxygenated sesquiterpene with molecular weight 236 (peak 55) are the main constituents comprising 53.4% of the oil. The total essential oil was fractionated by straight phase HPLC in four fractions. Each fraction was analyzed by capillary gas chromatography-mass spectrometry. The first fraction contained the sesquiterpene hydrocarbons. 17 of them were isolated and 13 identified. Seychellene (5.9% in total oil), cedradiene (1.8%), selina-4-(14),7-(11)diene (1.7%),  $\beta$ -patchoulene (1.6%), and  $\alpha$ -muurolene (1.0%) were the main constituents.

The fraction 2 (**Fig. 2**) was mainly composed of ketones and esters. The principal constituents are: fenchone, tujone, chrysanthenyl acetate, geranyl acetate, and two sesquiterpenoids.

Fraction  $F_3$  presented as main constituents 2 sesquiterpene compounds (MW 206 and 218 respectively) and a 236 molecular weight compound representing 6.0% of the total oil. The EI and CI mass spectra of this unknown compound are shown in Figure 2. The last fraction contained the principal components of the oil: isovaleric acid (15.2% in total oil) and patchouli alcohol (26.2%). Also nepetalactone (1.3%) was identified in this fraction.

The off-line combined technique of HPLC/CGC offers several advantages for essential oil analysis, e.g. fast pre-separation of small sample amounts (10 mg); no losses of volatiles by evaporation of the mobile phase; in on-column injection, the eluates can be injected as such; reduction of artifacts or alteration of the oil. As a result of this investigation 55 compounds have been classified by their mass spectral pattern; 37 were identified and 4 assigned tentatively.

The data are listed in **Table 1** together with the percentages for the compounds accounting for more than 1%. Two compounds of the essential oil of *Valeriana celtica* L. are known to possess feline attractant properties, namely isovaleric acid and nepetalactone. Nepetalactone is the main component of many *Nepeta* species, particularly in *Nepeta cataria* or Catnip. Its action on the olfactory sense of domestic cats was proven by *Waller et al.* [10].

### Acknowledgments

The authors thank Mr. *Silvio Stefanelli* curator of Giardino Botanico Alpino Paradisia, for botanical support and the supply of plant material.

*G. Bicchi* wishes to express his gratitude to Prof. *G. M. Nano* and Prof. *D. Sacco* for their interest and advice.

### References

- [1] *A. Fiori*, Nuova Flora Analitica d'Italia, **2**, 513, 2nd Ed., Bologna 1969. Ed. Agricole.
- [2] *E. Landolt* and *V. H. Hirzel*, Flora der Schweiz, **3**, 324, Basel and Stuttgart 1972. Hans Ernst Hen Birkhäuser Verlag.
- [3] *T. G. Tutin* and *V. H. Heywood*, Flora Europea **4**, 55, Cambridge University Press, Cambridge (1976).
- [4] *H. Schwarz*, Seifensieder Ztg. **68** (1941) 61.
- [5] *W. Obst*, Allgem. Oel- u. Fett-Ztg. **37** (1940) 411.
- [6] *P. Sandra*, *C. Bicchi*, *M. Mashaly*, and *M. Verzele*, in preparation.
- [7] *H. Kaiser* and *W. Lang*, Deut. Apotheker Ztg. **91** (1951) 163.
- [8] *M. Godefroot*, *M. Van Roelenbosch*, *M. Verstappe*, *P. Sandra*, and *M. Verzele*, HRC & CC **3** (1980) 337.
- [9] *P. Sandra*, *M. Schelfaut*, and *M. Verzele*, HRC & CC **5** (1982) 50.
- [10] *G. R. Waller*, *G. H. Price*, and *E. D. Mitchell*, Science **164** (1969) 1282.

MS received: January 10, 1983