

This article was downloaded by: [Ams/Girona*barri Lib]

On: 14 October 2014, At: 01:08

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Essential Oil Research

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tjeo20>

Chemical Composition of the Essential Oil Isolated from Wild Catnip *Nepeta cataria* L. cv. *citriodora* from the Drôme Region of France

Jean-Claude Chalchat ^a & Jacques Lamy ^b

^a Laboratoire de Chimie des Huiles Essentielles, Ecole Nationale Supérieure de Chimie et Université Blaise Pascal de Clermont-Fd, 63177, Aubière Cedex, France

^b Chambre d'Agriculture de la Drôme, Route de Venteroles, Nyons, France

Published online: 09 Dec 2011.

To cite this article: Jean-Claude Chalchat & Jacques Lamy (1997) Chemical Composition of the Essential Oil Isolated from Wild Catnip *Nepeta cataria* L. cv. *citriodora* from the Drôme Region of France, *Journal of Essential Oil Research*, 9:5, 527-532, DOI: [10.1080/10412905.1997.9700770](https://doi.org/10.1080/10412905.1997.9700770)

To link to this article: <http://dx.doi.org/10.1080/10412905.1997.9700770>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

RESEARCH REPORT

Chemical Composition of the Essential Oil Isolated from Wild Catnip *Nepeta cataria* L. cv. *citriodora* from the Drôme Region of France

Jean-Claude Chalchat*

Laboratoire de Chimie des Huiles Essentielles
Ecole Nationale Supérieure de Chimie et Université Blaise Pascal de Clermont-Fd
63177 Aubière Cedex, France

Jacques Lamy

Chambre d'Agriculture de la Drôme, Route de Venteroles, Nyons, France

Abstract

Nepeta cataria L. cv. *citriodora* growing wild in the Drôme region of France was brought into cultivation. Oils produced from cultivated plants harvested throughout the growing season were analyzed by GC and GC/MS. Although 42 components were identified, the oil composition did not depend on the time of harvesting or storage of the plant material prior to distillation. The oil was found to comprise mainly of citronellol (11.44-16.73%), nerol (19.95-30.70%), geraniol (25.13-31.00%) and geranial (4.93-11.05%). The highest oil yield was found to be at the time of full flowering.

Key Word Index

Nepeta cataria L. cv. *citriodora*, Labiatae, essential oil composition, citronellol, citronellyl acetate, geranial, nerol, geraniol.

Introduction

The genus *Nepeta* is represented by either annual or perennial herbs or shrubs most often found in temperate Eurasia and North Africa. Though not native to North America, three species have been found (1): *Nepeta cataria* L., including the garden escape cv. *citriodora*, *N. grandiflora* Bieb. and *N. racemosa* Lam. (Syn. *N. mustinii* Spreng. ex Henckel). Six other species (2) are in collections, in a Southern Ontario experimental garden [*Nepeta distans* Royle, *N. nepetella* L., *N. nuda* L. ssp. *nuda* (syn. *N. pannonica* L.) and *N. podostachys* Benth.], and in an experimental garden in North Carolina (*N. nervosa* Royle ex Benth. and *N. teydea* Webb. et Benth.).

N. cataria, usually called catnip, is native to Europe and Asia but is well-acclimatized in North America where it occurs in abundance near the Great Lakes. Its essential oil (2-7) is characterized by the presence of terpenes and isomers of nepetalactone, the most abundant of which can account for 99% of the total (2-7). An oil from *N. cataria* cv. *citriodora* (sometimes incorrectly named *Nepeta citriodora*)

*Address for correspondence

Received: February 1996
Revised: September 1996

Table I. Chemical composition of essential oil


Year of harvest		1989	1989	1989	1989	1989	1989	1989
Batch		1	2	3	4	5	6	7
No.	Name							
3	α -Pinene	0.65	0.30	0.09	0.42	0.32	0.12	0.19
4	α -Thujene	-	-	t	t	t	t	t
5	α -Pinene	1.54	2.88	0.72	2.38	0.79	0.50	0.99
6	Sabinene	-	-	0.19	-	0.23	t	0.23
7	Myrcene	t	0.11	t	0.08	-	t	t
8	Limonene	0.49	0.22	t	0.16	t	t	t
9	1,8-Cineole	-	-	t	t	t	t	t
11	γ -Terpinene	0.19	0.18	t	0.18	t	t	0.10
12	(E)- β -Ocimene	0.34	0.94	t	0.82	-	t	t
13	p-Cymene	0.20	t	t	0.08	-	t	t
14	Terpinolene	0.15	0.11	t	0.09	0.05	t	t
15	6-Methyl-5-hepten-2-one	0.14	0.10	t	0.08	0.08	0.07	0.19
16	trans-Rose oxide	t	0.12	0.14	0.08	0.08	0.32	0.17
17	cis-Rose oxide	-	t	t	t	t	t	t
18	Photocitral B	-	-	0.15	0.05	0.28	0.19	0.36
19	Photocitral A	-	-	t	-	t	t	t
20	Perillene	-	-	-	t	-	-	t
21	Artemisia ketone	-	-	0.10	0.06	0.09	0.08	0.10
22	Nerol oxide	t	0.11	t	0.09	0.10	t	0.11
23	cis,cis-Photocitral	0.23	0.12	0.25	0.11	0.14	0.19	0.28
24	Citronellal	-	-	1.28	-	0.90	0.69	1.03
25	trans,trans-Photocitral	0.34	0.28	0.56	0.28	0.98	0.58	0.77
26	Linalool	1.42	0.55	1.74	0.50	0.33	0.29	0.21
27	Linalyl acetate	0.20	0.12	10.75	0.15	0.37	0.24	0.13
28	β -Caryophyllene	3.24	4.89	3.39	4.81	3.27	2.74	2.62
29	Geranyl formate	-	-	-	-	t	t	t
30	Citronellyl formate	t	-	0.14	0.23	t	0.10	0.10
31	α -Humulene	0.28	0.41	0.26	0.37	0.29	0.25	0.24
32	Neral	3.07	3.31	7.79	3.17	5.81	6.10	6.68
33	Isoborneol	-	-	-	-	0.06	0.10	0.12
34	Neryl acetate	1.18	0.28	0.13	0.29	0.06	0.12	0.10
35	Piperitone	-	-	0.30	-	0.08	0.13	0.14
36	Methyl chavicol	0.44	0.44	0.47	0.42	0.57	0.60	0.71
37	Geranial	5.12	5.12	11.05	4.99	8.15	8.93	9.37
38	p-Menth-3-en-9-ol	0.56	0.64	0.29	0.63	0.62	0.60	0.83
39	Citronellol	15.10	16.73	12.02	16.47	14.36	13.66	14.62
40	Nerol	28.19	30.53	19.95	30.21	30.34	29.36	30.13
41	Geraniol	27.64	25.96	23.50	25.97	27.48	28.05	25.13
42	2-Phenylethyl hexanoate	-	-	-	-	t	0.05	-
43	iso-Caryophyllene oxide	-	-	0.19	-	0.21	0.26	0.43
44	Caryophyllene oxide	-	t	0.74	0.08	0.86	1.35	1.65
45	Humulene oxide*	-	-	0.09	-	0.09	0.11	0.15

*Correct isomer not given

of Nepeta cataria cv. citriodora oil

1989 8	1989 9	1989 10	1989 11	1989 12	1989 13	1989 14	1989 15	1989-I 16	1989-I 17
0.37	0.50	0.53	0.26	0.24	0.31	-	-	0.15	0.31
t	-	-	0.07	-	1.17	2.05	t	t	
2.43	3.16	3.04	1.83	1.92	2.32	0.34	0.48	0.89	1.56
0.67	0.92	0.79	0.51	0.46	0.51	-	0.05	0.19	0.37
t	0.07	-	-	0.02	-	0.07	0.08	t	-
0.12	0.06	-	0.05	0.09	0.08	0.11	0.07	t	-
0.19	0.06	-	t	0.04	0.11	-	-	t	-
0.28	0.30	-	0.09	-	-	-	-	t	-
-	-	-	-	-	-	-	-	t	-
-	-	-	-	-	-	-	-	0.05	-
-	-	-	-	-	-	-	-	t	-
-	-	-	0.16	-	-	-	-	0.26	0.76
-	0.02	-	0.29	0.06	-	-	0.05	0.09	-
-	0.09	-	0.05	0.09	-	-	t	0.04	-
-	0.06	-	-	0.04	-	-	t	0.34	0.15
-	0.03	-	0.06	0.14	0.19	-	0.16	t	t
-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	0.10	t
-	0.04	-	0.05	0.05	-	-	-	0.04	t
-	-	-	-	-	-	-	-	0.14	0.44
0.12	0.30	0.28	0.36	0.18	0.21	0.20	0.20	0.39	1.09
1.18	1.56	1.17	1.19	0.68	0.82	0.61	0.67	0.94	0.76
0.32	0.75	0.84	0.74	0.34	0.62	0.38	0.57	0.35	t
-	0.16	-	0.16	0.85	0.35	0.45	-	1.05	-
-	0.08	-	0.47	1.28	0.40	-	0.19	3.09	2.22
3.64	4.73	3.79	6.53	4.56	4.73	5.46	5.30	0.06	-
-	-	-	-	0.08	-	0.13	0.03	0.07	t
0.24	0.38	0.38	1.10	0.04	0.31	0.52	0.42	0.23	0.13
7.14	6.53	6.78	5.70	0.37	4.80	3.44	4.57	5.73	7.11
-	-	-	-	4.97	-	-	-	0.07	-
-	-	-	0.21	0.04	-	-	0.18	0.16	-
-	-	-	-	0.07	-	-	-	0.11	-
0.22	0.26	0.39	0.29	0.54	0.49	0.46	0.41	0.58	0.50
9.65	8.96	9.32	7.70	6.70	6.37	4.93	6.26	7.49	9.88
0.22	0.32	0.35	0.36	0.42	0.55	0.39	0.48	0.68	0.50
13.12	11.76	11.44	11.57	12.98	12.78	12.52	12.25	13.93	13.42
28.70	26.05	27.87	26.62	26.65	30.57	28.20	30.70	30.03	29.28
29.84	28.56	31.00	28.08	27.17	30.42	30.80	30.26	26.58	30.09
-	-	-	-	-	-	-	-	t	-
-	0.07	-	0.14	0.65	0.22	0.08	0.17	0.23	t
0.19	0.37	0.46	0.60	3.25	1.41	0.42	1.33	0.94	0.23
-	-	-	0.05	0.18	-	-	-	0.08	t

Downloaded by [Ams/Girona*barri Lib] at 01:08 14 October 2014

Figure 1. Chromatogram of *Nepeta cataria* L. cv. *citriodora* oilTable II. Physical and chemical constants of essential oils of *Nepeta cataria* cv. *citriodora*

Year	Batch	n_D^{20}	$[\alpha]_D^{20}$	d_{20}^{20}
1989	3	1.4842	-2.54	0.906
1989	4	1.4820	-2.43	0.896
1989	6	1.4840	-1.71	0.895
1989	7	1.4831	-2.19	0.891
1991	12	1.4793	-3.70	0.904
1991	13	1.4863	-2.63	0.892
1991	15	1.4839	-2.25	0.892

Table III. Oil yields obtained from *Nepeta cataria* cv. *citriodora* harvested at different stages of maturity.

Batch	Harvest date	Distillation Date	Mass distilled (kg)	Total oil obtained (mL)	Yield of oil (mL/kg)	Vegetative stage		
						Buds	Flowers	Faded flowers
1	Jul 19, '89	Jul 19	29.0	25.0	0.86	50	50	
2*	Jul 19, '89	Jul 19	23.0	18.0	0.78	50	50	
3**	Jul 24, '89	Jul 24	5.2	3.6	0.70			
4	Jul 28, '89	Jul 29	4.4	5.0	1.13	30	50	20
5	Aug 2, '89	Aug 4	5.9	7.0	1.18	20	30	50
6***	Aug 2, '89	Aug 4	4.9	7.0	1.40	20	30	50
7	Aug 16, '89	Aug 18	16.9	24.0	1.40	end of flowering		
8	Jun 28, '91	Jun 26	5.2	2.2	0.42			
11	Jul 14, '91	Jul 14	7.0	7.9	1.10			
12***	Jul 14, '91	Jul 31	5.2	7.0	1.20			
13***	Jul 14, '91	Aug 21	6.4	11.0	1.70			
14***	Jul 14, '91	Aug 28	6.6	7.2	1.10			

*Very green plants; **Yellow plants; ***Dry distilled batches

which was analyzed by Regnier (3), was found to comprise mainly of citronellol, geraniol, neral and geranial with nepetalactones making up at most about 10% of the mixture.

In 1980, Zamureenko (13) found the same components in oil from the former Soviet Union, but no nepetalactone was determined. In 1981, Dmitriev et al. determined the variations in oil accumulation and chemical composition during the growth cycle and storage of the harvested catnip plants that they referred to as *N. cataria* var. *citriodora* Balb. (14).

N. cataria cv. *citriodora* grows wild in the Drôme region of France. The results of initial work on wild plants have induced local herb growers to cultivate this species and develop pilot-scale industrial production of the oil. We report here the first analytical results obtained in 1989 and 1991 of the oil composition carried out on cultivated plants transplanted from the wild.

Experimental

Plant material: *N. cataria* cv. *citriodora* was harvested in July 1989 and 1991 in the Drôme region when the plants were in full flower. Except for batches 12, 13, 14 and 15, which were produced from dried plant material harvested on 14 July 1991 and distilled 31 July, 21 and 28 August and 3 September, all other oils were isolated from freshly harvested plant material. Oil isolation was performed by steam distillation (2-litre/h) on a stainless steel experimental pilot apparatus fitted with two 10-litre vessels custom built at the laboratories of the Chamber of Agriculture at Nyons, France. Pilot tests were carried out on cultivated plant material distilled in a 2000-litre industrial still at the farm.

Analysis: Physical and chemical characteristics of the oils such as refractive index, density and optical rotation, were determined by standard methods.

Analysis by GC was carried out on a DELSI 121C chromatograph fitted with a 25 m x 0.25 mm CP Wax 52CB capillary column with temperature programming from 50°C (5 min) to 220°C at 2°C/min. Injector and detector temperatures were set at 240°C and 255°C and the split ratio was 1/60.

GC/MS coupling was achieved using a SIGMA 300 chromatograph coupled to an HP 5970 mass spectrometer fitted with a 50 m x 0.3 mm CP Wax column which was temperature programmed from 60°-240°C at 2°C/min. Ionization voltage was 70 eV.

Components were identified by comparison of mass spectra with those reported in the literature (19), calculation of retention indices and also comparing them with our own as well as published data, and by the co-injection with standard compounds.

Results and Discussion

Forty-five components were identified in the oil (Table I, Figure 1). The oil of catnip studied, which was similar to that described elsewhere (3,14,15-17), was composed mostly of citronellal, neral, geranial, citronellol, nerol and geraniol (85 to 95%). In addition, small amounts of other oxygenated monoterpenes were also identified. Monoterpenes were also present, while β -caryophyllene and α -humulene together with their oxides were the only sesquiterpenes found. This may be explained by the variation of chemical composition during the hydrodistillation (18). It was surprising to us that no nepetalactone were found in the oils produced. As a result, we believe that this is a distinguishing feature of this cultivar of *N. cataria*.

As can be seen from Table I the chemical composition of the oil of *N. cataria* cv. *citriodora* varies little during its life cycle (batches 1 to 7). It is apparently not affected by drying the plant material before distillation, or by storage before distillation (batches 13 to 15). Also, large scale distillation (batches 16 and 17) yielded an oil similar to that obtained from lab-prepared oils. Given the consistent chemical composition, it is our belief that production could be optimized by choosing the time of harvest that afforded the highest yield (mid-August). Yields were much lower if plants were harvested earlier in the season (July) (batches 8, 11 and 12) (Table II). The physical and chemical characteristics (refractive index, density and optical rotation) of the batches analyzed were remarkably stable and did not vary with time of production (Table III).

The specific chemical composition of this oil gives it a potentially useful olfactive profile. Its consistent physical and chemical characteristics are not appreciably influenced by production conditions, which remain to be optimized. These features make it a potential complement or replacement for melissa (*Melissa officinalis*) and "verveine citronnée" (*Verbena citriodora*) oils, *Aloysia triphylla* (L'Herit.) Britton (syn. *A. citriodora* Palau) oils used in perfumery and flavorings.

References

1. B. M. Lawrence, *Essential oils, 1979-1980*. pp 76-79, Allured Publ. Corp., Carol Stream, IL (1981).
2. B. M. Lawrence, *Essential oils, 1988-1991*. pp 98-199, Allured Publ. Corp., Carol Stream, IL (1992).
3. F. E. Regnier, E. J. Eisenbraun and G. R. Waller, *Nepetalactones and Eptnepetalactone of Nepeta cataria L.* Phytochemistry, **6**, 1271-1280 (1967); F. E. Regnier, E. J. Eisenbraun and G. R. Waller, *Studies on the composition of the essential oils of three Nepeta species*. Phytochemistry, **6**, 1281-1289 (1967).
4. D. L. Nelson, *Methylcyclopentane monoterpenes of Nepeta cataria and Actindia polygama*. PhD thesis, Purdue University (1968).
5. S. D. Sastry, W. R. Springstube and G. R. Waller, *Identification of 5,9-dehydronepetalactone, a new monoterpene from Nepeta cataria*. Phytochemistry, **11**, 453-455 (1972).
6. S. M. McElvain, R. D. Bright and P. R. Johnson, *Constituents of the volatile oil of catnip. I. Nepetalic acid, nepetalactone and related compounds*. J. Amer. Chem. Soc., **63**, 1558-1563 (1941).
7. S. M. McElvain, P. M. Walters and R. D. Bright, *Constituents of the volatile oil of catnip. II. The neutral components of Nepetalic anhydride*. J. Amer. Chem. Soc., **64**, 1828-1831 (1942).
8. F. Bellesia, R. Grandi, U. M. Pagnoni, A. Pinetti and R. Trave, *Biosynthesis of Nepetalactone in Nepeta cataria*. Phytochemistry, **23**, 83-87 (1984).
9. A. O. Tucker and S. S. Tucker, *Catnip and the Catnip Response*. Economic Botany, **42**, 214-231 (1988).
10. K. Sakurai, K. Ikeda and K. Mori, *Both (4aS, 7S, 7aR)-(+)-Nepetalactone and its antipode are powerful attractants for cats*. Agric. Biol. Chem., **52**, 2369-2371 (1988).
11. A. Velasco-Negueruela, M. Mata Rico, P. Bermejo Benito and M. J. Pérez-Alonso, *Composición de los aceites esenciales de nepeta nepetella subsp. aragonensis, Nepeta coerulea subsp. coerulea y Nepeta cataria*. Giorn. Bot. Ital., **122**, 295-302 (1988).
12. S. S. Mishurova and Sh. R. Mamedova, *Morphological characteristics and essential oil composition of Nepeta cataria L.* Izv. Akad. Nauk Az. SSR, Ser. Biol. Nauk, **3**, 11-17 (1989).
13. V. A. Zamureenko, N. A. Klyuev, M. G. Mumladze, L. B. Dmitriev and I. I. Grandberg, *Identification of components of Nepeta cataria var. citriodora Balb. essential oil*. Izv. Timiryazevsk. S-kh. Akad., **5**, 167-169 (1980).
14. L. B. Dmitriev, M. G. Mumladze, G. A. Esvandzhiva, I. I. Grandberg and I. Z. Yakubashvili, *Dynamics of the accumulation and composition of the essential oil of catnip (Nepeta cataria var. citriodora Balb.) during vegetation and storage of raw material*. Izv. Timiryazevsk. S-kh. Akad., **3**, 75-81 (1981).
15. I. G. Kapelev, *Introduction and cultivation of new species of essential-oil-bearing plants*. Maslo-Zhir. Prom-st, **8**, 25-27 (1982).
16. G. Tittel, H. Wagner and R. Bos, *Chemical composition of Melissa oil*. Planta Med., **46**(2), 91-98 (1982).
17. I. I. Lyashch and S. I. Vasil'kevich, *Chemical composition of spice plants and their infusions*. Vestsi Akad. Navuk BSSR, Ser. Biyal. Navuk, **3**, 30-32 (1987).
18. C. Bourrel, F. Perineau, G. Michel and J. M. Bessière, *Catnip (Nepeta cataria L.) essential oil: Analysis of chemical constituents, bacteriostatic and fungistatic properties*. J. Essent. Oil Res., **5**, 159-167 (1993).
19. F. W. McLafferty and D. B. Stauffer, *The Wiley/NBS Registry of Mass Spectral Data*. Vol. 1, Wiley and Sons, New York (1988).