

12 Physiological and Pathological Causes of Behavioural Change

Introduction

Overt behaviour is the consequence of a cat perceiving some change in its environment, evaluating this change, deciding on an appropriate response and the response being generated through the motor systems of the brain to the elements of the skeletal system that control activity. Hence, although behaviour occurs as a consequence of changes in the external environment, the responses generated also depend on internal variations in the processing of information. These processes are susceptible to alteration due not only to normal physiological variations but also to pathological changes. Interpretation of behaviour, therefore, requires an understanding of how the generation of behaviour is modulated by factors influencing the internal state, as well as how responses are generated to events in the external environment.

Pathological changes can be the sole cause of behavioural change – indeed, behavioural signs such as lameness are common first indicators of disease in veterinary medicine. In some cases complex behavioural signs, such as aggressive behaviour towards an owner, can occur entirely as a consequence of pathological events, such as focal seizures. Although such events are rare, their characteristics need to be distinguished from behaviours generated in response to external stimuli when investigating the cause of undesired behaviours. However, physiological or pathological changes more commonly modify cats' behaviour rather than solely cause it. This is through alterations in one or more of the following: (i) perception of external events; (ii) the motivation to show a response; (iii) the threshold at which a response is shown; and (iv) the manner in which a response is generated. Understanding the way that behaviour develops in individual cats, therefore, requires both an understanding of how behaviour is modulated through learning (see [Chapter 3](#), this volume) and how disease can modify such processes. The physical examination of cases by a veterinary surgeon is therefore important before investigating individual cases of undesired behaviour (Fatjó and Bowen, 2009).

In addition to disease processes being important factors in the development of undesired behaviours, cats' responses to their environment can also influence the onset or development of disease. For example, stress can influence susceptibility to infectious disease, or the shedding of infectious agents from animals with carrier status (e.g. Addie *et al.*, 2009). Stress caused by environmental factors also influences immune functioning and the onset of bouts of chronic diseases such as feline idiopathic cystitis (Seawright *et al.*, 2008).

Since it is impractical to consider all possible medical causes of behavioural change here, we will use a functional approach to consider how disease processes may alter the motivation for behavioural responses in the cat, modify their occurrence or generate new responses. First, we consider how physiological or pathological changes may alter an individual cat's relative motivation to show behaviour, hence changing its frequency or timing. Next, we examine how normal behavioural responses to external events can be modified by internal factors, and where behaviours are generated by disease states. Finally, we give examples of how individual responses to the environment can influence the onset of disease in the cat.

Effects on Motivation to Show Normal Behaviour

Increased or decreased motivation to show normal responses, or the initiation of new behaviours, can all occur in association with physiological and pathological changes. Normal physiological variations that may influence the initiation of behaviour include the fluctuation of female reproductive hormones over the oestrous cycle in entire females. For example, profound changes in the behaviour of entire queens occur with the oestrous cycle, notably related to mate seeking (see [Chapter 8](#), this volume), and this may or may not be seasonal depending on regional photoperiods (Faya *et al.*, 2011).

Disease processes can also modify the initiation of behaviour by altering underlying motivation. For example, a cat's motivation to acquire food may be increased by endocrine disorders such as hyperthyroidism (Salisbury, 1991). In such cases, behaviours aimed at obtaining food, such as vocalizing or rubbing around the owner, occur more frequently or in different spatial or temporal contexts. Similarly, conditions decreasing appetite may reduce food-soliciting behaviours towards the owner. Excessive eating or drinking caused by a medical condition will inevitably lead to a higher frequency of toileting behaviour. Hence a cat with a condition causing polydipsia (excessive drinking) may potentially

show signs of inappropriate urination, because access to toileting sites becomes limiting with the increased need to eliminate. Similarly in conditions affecting metabolic rate or body temperature, cats may show altered motivation to seek sources of heat (e.g. in hyperthyroidism) or cool locations (e.g. fever). In some cases such changes can lead to the onset of undesired behaviours, perhaps in combination with situational factors. For example, a cat in a household with others that are not socially compatible may only be able to access a litter box intermittently, but concurrent disease-induced polyuria may bring about a change to an 'inappropriate' toileting site due to insufficient access for toileting needs.

Pain is one of the most common reasons for cats to alter their motivation to show behaviours. For example, cats may have a reduced desire to play or interact when suffering from joint pain, and this may be the first sign of which owners are aware in the development of osteoarthritis in older cats (Lascelles and Robertson, 2010). Pain may also lead to context-specific responses: for example, a cat with a bite abscess on the tail base may avoid being stroked by its owners, perhaps by moving away or even showing aggression to stop owner contact. Pain is processed through a network of structures known as the 'pain matrix' (Jones *et al.*, 2003). The system has two parallel pathways, one of which (the medial pathway) is involved in the emotional component (i.e. sensation) of pain; the other (lateral) pathway is involved in discriminating the particular details of the stimulus (i.e. type of sensation, location, intensity and duration). Both pathways involve the thalamus and pass through to the insular and somatosensory cortices, respectively (Kulkarni *et al.*, 2005). On contact with a painful stimulus, the areas of the brain associated with fear and anxiety are activated (Rainville, 2002); this is obviously adaptive as it enables the animal to learn the salient aspects of an environment that lead to a paineliciting outcome.

Altered motivation to groom areas of the body is also a common result of conditions causing irritation to the skin, and thus activation of mechanoreceptors (see [Chapter 2](#), this volume). A range of conditions including parasite infestation and atopic, infectious or autoimmune-mediated skin disease can result in a cat showing increased grooming, biting or other skin-directed behaviour.

In cases of general illness or infection, animals commonly show what are termed 'sickness behaviours' (Johnson, 2002), which have a range of functions such as conserving energy or promoting defensive mechanisms in the body such as fever (Broom, 2006). Response to infection is mediated by the release of cytokines, which have a direct effect on the brain, leading to increased sleep and reduced activity. Cytokines are also suggested to influence activation of the

hypothalamo–pituitary–adrenal axis and hence modulate behavioural responses (McCann *et al.*, 2000). Reduced motivation to be active or interact socially may also occur with sensory deficits, such as loss of sight (Turner, 2004).

Modulation of Existing Responses to External Events

In many cases, changes in internal state through physiological variation or disease do not lead to the initiation of a behaviour, but modify existing behavioural responses to external events. These effects can occur through the input of sensory information to the brain, the evaluation of information or in the generation of a motor response.

Modulation of sensory information inputting the brain

Information entering the brain is generated in the sensory organs such as the retinal cells in the eye, the nasal epithelium or the sensory receptors in the skin (see [Chapter 2](#), this volume). Conditions that influence the relative activation of these organs by external stimuli, or the passage of impulses along the sensory nerves to the sensory cortices of the brain, will influence the extent to which cats respond to external events. For example, Siamese cats have an abnormal development of sensory input from the retina to the lateral geniculate nucleus, which means that their capacity to use stereoscopic vision to determine depth (see [Chapter 2](#), this volume) is limited, and it is believed that the development of a squint in some individuals is to enable some overlap of visual fields (Hubel and Wiesel, 1971; [Chapter 2](#), this volume). While the plasticity of the sensory cortex enables most Siamese to respond apparently normally to external events, reduced or absent binocular capability is likely to modify when and how individual cats respond to events, and potentially influence their hunting ability. Perception of olfactory information is dependent on the integrity of the nasal epithelium, and chronic damage from infectious upper respiratory tract disease can influence the relative perception of scents in the environment (Scherk, 2010), influencing a range of scent-related behaviours such as toileting and social interaction (see [Chapters 5 and 8](#), this volume).

Modulation of sensory information from the integument is another example of where abnormalities can influence the extent to which cats perceive external stimuli. Changes in the relative activity of peripheral mechanoreceptors can result in either altered sensation (dysaesthesia) or exaggerated response to

stimulation (hyperaesthesia) (Rizzo *et al.*, 1996). These changes in relative activation of sensory neurons arise as a result of changes in neuronal conduction properties with variation in sub-type of sodium ion channels in the nerve membrane (Waxman *et al.*, 2000). Damage to a peripheral sensory nerve may result, for example, in relative hyperpolarization, such that the threshold of activation by touch sensation is reduced (Matzner and Devor, 1992). These changes in peripheral nerve activation thresholds following damage are one potential underlying cause for behavioural changes in cats sometimes described as 'hyperaesthesia syndrome' or 'hyperkinesis' (Shell, 1994), where cats present with twitching or rippling skin and may jump around to groom intensely, as if responding to severe irritation, following only a mild touch sensation.

Factors influencing processing and evaluation of sensory information

Behaviour arising as a consequence of external changes can also be modulated by factors that influence how information is centrally processed or appraised. Normal physiological changes can influence these processes, such as fluctuations in reproductive hormones, since sex steroids act as modulatory neurotransmitters across many regions of the brain (Rupprecht and Holsboer, 1999). For example, evidence from other species suggests that changes in oestrogen may have profound effects on the activity of a range of different neurotransmitters at the cellular level, resulting in an increased chance of behavioural responses to external events and a reduced threshold of response to painful stimuli (Aloisi and Bonifazi, 2006).

Pathologies can also influence how sensory information is evaluated, and change either the threshold for the behaviour or the degree of behavioural response. Endocrine abnormalities can have a profound effect on responses to external events, as well as a range of physical signs. In Cushing's disease, for example, overproduction of glucocorticoids may ultimately reduce the production of corticotrophin-releasing factor in the hypothalamus due to negative feedback, resulting in an animal that is unresponsive to external events and lethargic. Corticosteroids are also commonly used in the treatment of various medical conditions in cats, and evidence from human patients suggests that iatrogenic steroids may also have effects on mood and behaviour (Brown and Chandler, 2001).

Diseases that influence the breakdown and metabolism of dietary components can also have an influence on the threshold of response generation in the brain. The most common examples are cases where either hepatic (liver) or renal

(kidney) function is compromised, resulting in a reduced ability to remove by-products of digestion from the circulation and excrete these via the urine. The behavioural and neurological consequences of such diseases are commonly termed 'encephalopathies'. The effects on brain function can arise through the build-up of compounds that are neurotoxic, or alternatively can be caused by the relative unavailability of amino acids necessary for neurotransmitter turnover. For example, increased ammonium salts in the circulation as a result of hepatic insufficiency cannot be metabolized to urea in the central nervous system, and are instead converted to glutamine, which ultimately leads to an excess of this amino acid in the brain (Albrecht *et al.*, 2007). Since glutamine is the precursor for the excitatory neurotransmitter glutamate, this may explain transient decreases in the threshold of responding (for example, increased aggression) associated with encephalopathies. In addition, chronic low-grade hyperammonaemia has been associated with memory deficits and difficulty in adapting to new environments in both humans and rats (Apelqvist *et al.*, 1999). Because increased levels of ammonia, and hence increased glutamine and glutamate, occur as a protein meal is digested and the by-products circulate in the bloodstream, behavioural signs associated with hyperammonaemia will often occur temporarily, associated with feeding. Ultimately, increasing levels of glutamine result in cerebral oedema (Albrecht *et al.*, 2007) and more obvious neurological symptoms, such as loss of motor coordination, depression, hysteria, pacing, circling, seizures and, ultimately, coma and death.

Alterations of normal behaviour can arise as a consequence of disorders of the central nervous system. The specific effects of lesions on behaviour will relate to the specific area of the central nervous system affected. For example, lesions affecting the pineal gland may influence the sleep-wake cycle (Uz *et al.*, 2003). In older cats, behavioural changes in response to external events are commonly caused by a range of clinical conditions including osteoarthritis, systemic hypertension (often secondary to chronic kidney disease or hyperthyroidism), hyperthyroidism or sensory deficits (Gunn-Moore, 2011). In some cases such changes may result from reduced motivation to show new behaviours, for example due to pain, as discussed earlier. There is also a tendency across species for cognitive ability to decline with increasing age. For example, in both humans (Cherry and Park, 1993) and dogs (Christie *et al.*, 2005) the ability to perform allocentric tasks (i.e. requiring reference to external landmarks) declines with age. However, in a proportion of cats more profound changes in cognitive ability and behaviour are associated with specific pathological changes analogous to Alzheimer's disease in people, known as

‘cognitive dysfunction’ (Landsberg *et al.*, 2010). Associated alterations in response to external events might include reduced social interaction with owners or, alternatively, increased dependency on owner attention and presence, loss of previously learnt associations, altered or ‘mixed-up’ responses to previously learnt cues (e.g. change in toileting substrate preference) or general disorientation.

Influences on motor outputs and initiation of behaviour

Disease processes may also modify behavioural responses to external events by altering the output of motor information from the brain and the activation of peripheral muscles via neuromuscular junctions. Changes in this element of the pathway may influence the cat’s ability to show a desired behaviour, or alter the form of the behaviour. A cat’s ability to show a response may be influenced by diseases affecting the motor cortex of the brain or spinal cord. For example, spinal damage can influence the motor control of elimination, potentially resulting in urination or defecation in undesired locations. Control of toileting can also be influenced by conditions affecting the gastrointestinal or urinary tract. Diseases causing inflammation in the bowel, or those affecting absorption, will potentially influence both the frequency and urgency of defecation.

Feline idiopathic cystitis (FIC), also known as idiopathic feline lower urinary tract disease (iFLUTD), is the most common medical cause of abnormal urination in the cat, and hence is an important differential diagnosis to consider when investigating cats presenting with inappropriate elimination (Buffington *et al.*, 1997). FIC is termed ‘idiopathic’ when there is no obvious physical cause to account for the condition, and is diagnosed by excluding other causes of lower urinary tract inflammation (such as urinary tract infection, urethral strictures, neoplasia or urolithiasis) (Kalkstein *et al.*, 1999). Inappropriate elimination may be the first presenting sign of FIC, and is thought to occur because the cat associates the pain of urination with the specific location in which it has previously urinated. In addition, the condition causes increased frequency and urgency of urination. Because the signs of the condition are commonly present during bouts of 3 or 4 days, affected cats may show repeated changes in the location of urination for several days in a row. Pain on urination can also cause the cat to appear distressed and vocalize before and during urination, and abdominal pain may make cats reluctant to be handled, either moving away or showing aggression as an avoidance response. Male cats may also change their posture from a squat to standing up, as squatting, which bends the urethra, may

cause further discomfort (Seawright *et al.*, 2008).

The ability of cats to show desired behavioural responses can also be influenced by diseases affecting movement, such as degenerative joint disease (e.g. osteoarthritis), neuropathies of motor nerves (e.g. diabetic polyneuropathy), disorders of the neuromuscular junction (e.g. myasthenia gravis) or problems with muscular functions (e.g. myopathies associated with feline leukaemia virus).

Behaviours Occurring Entirely as a Result of Disease Processes

In some cases behavioural signs in cats are generated entirely as a result of a disease process, and occur unrelated to external events (Reisner, 1991). In general these types of behaviour are less common than where pathology acts to modify behavioural responses to external events. Abnormalities that can generate a response may occur in any part of the process by which a normal response is generated (i.e. sensory input to the brain, the processing of information in the central nervous system or the output of information from the motor cortex).

The generation of abnormal sensory information will create similar behavioural responses to sensory inputs based on actual external stimuli. For example, in addition to dysaesthetic and hyperaesthetic responses in peripheral sensory nerves (described previously), paraesthetic responses can occur, whereby spontaneous activity is generated in the nerve without any mechanical stimulation of the corresponding receptors. This abnormal or ectopic impulse generation is also commonly called sensory neuropathy. Because these abnormal inputs are processed as if they arise in response to normal external events, the reaction of the cat is appropriate to irritation or pain arising from the source (i.e. grooming, scratching or attempting to bite at the affected part of the body). In some cases these responses can be extreme and cause both cat and owner considerable distress. The tail is a common target for such behaviour, possibly because the sensory nerves in the tail are more susceptible to damage (e.g. through bite injuries or the tail being caught in a doorway). A cat may repeatedly and persistently attack its tail to the extent that it becomes damaged and requires repeated medical or surgical intervention. Depending on the origin of the sensory damage, and the extent to which sensitization of the response has occurred, the behaviour can even continue following amputation of part or all of the tail. An

abnormal pain sensation in the face and mouth region has also been described, particularly in oriental breeds. Although there may be multiple possible factors involved in the development of this condition, damage to the facial nerves, such as the trigeminal nerve, is likely to be an important factor. Cats suffering from this condition show signs varying in severity; in the most extreme cases cats can cause considerable self-damage by clawing at their own face in an apparent attempt to alleviate the pain (Rusbridge *et al.*, 2010).

Spontaneous activity through ectopic activity in nerve cells also occurs in the brain. Spontaneous depolarization shifts or repetitive discharge from hyperexcitable collections of neurons in the brain are expressed as epileptic seizures. In the typical 'grand mal' seizure, this activity spreads across the whole brain. However, the spread of electrical activity can also be localized, influencing only one part of the brain. The behavioural response arising from these focal seizures will depend on the part of the brain that is affected and the extent to which it spreads (Dahl, 1999). Hence, a focal seizure in the visual cortex is likely to result in a cat responding as if it has seen something (i.e. a 'visual hallucination'). The response will often be identical to that as would occur if the animal had seen a 'real' stimulus, since activation of this brain region will lead to the generation of the same behavioural response as if the activation was due to inputs via the optic nerve. Hence, cats may appear to pounce on or chase objects which are not there. Seizure activity localized to the limbic part of the brain, responsible for the generation of emotional responses, can result in sudden unprovoked behavioural responses indicative of extreme emotional disturbance. For example, cats may suddenly run off and hide, or show extreme aggression to the nearest person, object or other animal. Where the focus of focal seizures is the motor cortex, resulting behaviours are often rather more fixed in form, rather than the more 'goal-oriented' and variable behaviours associated with sensory focal seizures.

Seizure activity develops over time through the process of 'kindling', whereby spontaneous activity in one area gradually results in the 'recruitment' of neighbouring cells (Bertram, 2007). In the early stages, seizure activity may therefore result in relatively mild signs which are often disregarded. For example, humans with seizures in the limbic part of the brain may initially report mild 'feelings' occurring spontaneously, which develop over time into more obvious seizures with clear behavioural indicators (Bertram, 2007). The plasticity of the brain, and susceptibility of cells to repeated electrical stimulation, means that behavioural changes may be used to identify early seizure activity (Shihab *et al.*, 2011). The other factor to consider with epilepsy

is the effect of seizures on behaviour occurring between the episodes. Because the action of the seizures themselves is to partially 'remodel' areas of the brain, the response of the animal to normal stimuli in the interictal (between seizure) period can also be affected (Adamec, 2003).

Persistence of Behavioural Outcomes of Disease

When evaluating the behaviour of an individual animal, it is important not only to consider currently occurring physiological or pathological impacts on behaviour, but to investigate the potential impact of historical effects. In many cases, behavioural changes can arise as a consequence of a previous condition that has been resolved, but where the behavioural response has been retained through learning. For example, male cats with FIC may stand to urinate in order to facilitate the passage of inflammatory material, or due to urethral spasm. However, where these individuals learn that this posture eases the pain associated with elimination, they may continue to show this posture even after resolution of the disease. Similarly, avoidance responses learnt when a cat was in pain may be retained after resolution of the painful lesion. Where aggression to an owner stroking the caudal dorsum effectively prevents contact on the tail base, a cat may continue to respond aggressively after resolution of a tail base abscess because it has not had the opportunity to learn that contact in this area is no longer painful. Behaviours originating for medical reasons can also become reinforced and established for other reasons. For example, grooming behaviour initiated because of an irritating lesion could become reinforced by the owner paying the cat attention when the behaviour is shown. This type of reinforcement is particularly likely where cats highly value owner attention, as response from the owner will be a valued outcome.

Environmental Stressors in Disease Aetiology

It is increasingly recognized in human medicine that psychological stressors impact extensively on somatic disease (Nater *et al.*, 2006). Similarly, in veterinary medicine, the influence on disease processes of environment, and individual responses to environmental change, have gained in recognition (Casey, 2010). However, much still needs to be investigated to elucidate the relationship between stress exposure and disease susceptibility. It is unclear, for

example, why some individuals appear to be more susceptible to somatic disease than others exposed to similar environments. In addition, it is not known why different individuals exposed to the same stressors develop conditions involving different body systems. However, it is likely that some individuals are predisposed to vulnerability, such that co-occurrence with extreme environments leads to dysregulated stress responses, resulting in disease. Further research is needed to investigate the nature of this vulnerability, and how it interacts with stressors that lead to such consequences.

In cats, FIC is the condition that is most widely recognized as associated with exposure to stress (Seawright *et al.*, 2008). Indeed, differentiating cases of FIC from inappropriate elimination can be complicated, as exposure to events that cats find aversive is an important 'flare factor' in the multifactorial aetiology of this condition (Cameron *et al.*, 2004), and hence similar factors can lead to both inappropriate elimination and FIC. Other chronic conditions in which stress exposure is considered a factor in humans include irritable bowel syndrome (Murray *et al.*, 2004) and chronic skin disease (Kimyai-Asadi and Usman, 2001), both of which are anecdotally also relevant in cats. The hyperaesthetic sensory responses, as described earlier, are also postulated to have environmental stress as an aetiological factor.

In addition to chronic disease conditions, where clinical signs appear to be exacerbated or precipitated by stressful events, acute stress can have important impacts on susceptibility to infectious disease. For example, the high level of stress experienced by many cats when entering a boarding or rescue cattery, associated with loss of predictability and change in environmental cues, leads to an increased susceptibility to infectious disease, at the same time as they are brought into the proximity of other cats, thus increasing the probability of contact with infectious agents.

Conclusions

Important welfare implications arise from the appropriate recognition of the influences of disease on behaviour. Behavioural signs may be the first indicators of disease and enable the early treatment of potentially serious medical conditions. In addition, the recognition of confounding disease processes will ensure that behavioural interventions are considered that take these into account, and better inform prognosis when treating undesired behaviours. Furthermore, where stress is likely to impact on disease risk, interventions to reduce exposure to stressors can have important health impacts.

to stressors can have important health impacts.

In considering the relationship between pathological disease or physiological variation and behavioural signs it is important to consider not only those behaviours that are directly caused by pathologies, but also those where changes in internal state may modify existing responses. In addition, it is imperative to adopt a rational, functional approach and consider those aspects of normal response generation where pathology could be having an effect. For example, where a cat is observed attacking its tail, the important factors to investigate and differentiate from behavioural causes will range from conditions that cause irritation of the skin right through to those that might be generating an abnormal motor response.

References

- Abrahams, V.C., Hodgkins, M. and Downey, D. (1987) Morphology, distribution and density of sensory receptors in the glabrous skin of the cat rhinarium. *Journal of Morphology* 191, 109–114.
- Adamec, R. (2003) Kindling induced lasting interictal alterations of affective behavior. *Annals of the New York Academy of Sciences* 985, 495–497.
- Adamec, R.E. (1976) The interaction of hunger and preying in the domestic cat (*Felis catus*): an adaptive hierarchy? *Behavioural Biology* 18, 263–272.
- Adamec, R.E., Stark-Adamec, C. and Livingstone, K.E. (1980) The development of predatory aggression and defense in the domestic cat (*Felis catus*). *Behavioural and Neural Biology* 30, 389–409.
- Adamec, R.E., Stark-Adamec, C. and Livingstone, K.E. (1983) The expression of an early developmentally emergent defensive bias in the adult domestic cat (*Felis catus*) in non-predatory situations. *Applied Animal Ethology* 10, 89–108.
- Adamelli, S., Marinelli, L., Normando, S. and Bono, G. (2005) Owner and cat features influence the quality of life of the cat. *Applied Animal Behaviour Science* 94, 89–98.
- Addie, D., Belák, S., Bourcraut-Baralon, C., Egberin, H., Frymus, T., Gruffydd-Jones, T. *et al.* (2009) Feline infectious peritonitis: ABCD guidelines on prevention and management. *Journal of Feline Medicine and Surgery* 11, 594–604.
- Albrecht, J., Sonnewald, U., Waagepetersen, H.S. and Schousboe, A. (2007) Glutamine in the central nervous system: function and dysfunction. *Frontiers in Bioscience* 12, 332–343.
- Aldis, O. (1975) *Play Fighting*. Academic Press, New York.
- Aloisi, A.M. and Bonifazi, M. (2006) Sex hormones, central nervous system and pain. *Hormones and Behaviour* 50, 1–7.
- APBC (2005) Annual review of cases 2005. Available at: http://www.apbc.org.uk/sites/default/files/review_2005.pdf (accessed 11 September 2012).

- Apelqvist, G., Hindfelt, B., Andersson, G. and Bengtsson, F. (1999) Altered adaptive behaviour expressed in an open-field paradigm in experimental hepatic encephalopathy. *Behavioural Brain Research* 106, 165–173.
- Appleby, M.C. (1993) How animals perceive a hierarchy: reactions to Freeman *et al.* *Animal Behaviour* 46, 1232–1233.
- Archer, J. (2011) Pet keeping: a case study in maladaptive behaviour. In: Salmon, C.A. and Shackelford, T.K. (eds) *The Oxford Handbook of Evolutionary Family Psychology*. Oxford University Press, New York, pp. 281–296.
- Bacon, B.A., Lepore, F. and Guillemot, J.-P. (1999) Binocular interactions and spatial disparity sensitivity in the superior colliculus of the Siamese cat. *Experimental Brain Research* 124, 181–192.
- Bang, A., Deshpande, S., Sumana, A. and Gadagkar, R. (2010) Choosing an appropriate index to construct dominance hierarchies in animal societies: a comparison of three indices. *Animal Behaviour* 79, 631–636.
- Bard, P. and Macht, M.B. (1958) The behaviour of chronically decerebrate cats. In: Wolstenholme, G.E.W. and O'Connor, C.M. (eds) *CIBA Foundation Symposium on the Neurological Basis of Behaviour*. J. & A. Churchill, Ltd, London, pp. 55–71.
- Barratt, D. (1997) Home range size, habitat utilization and movement patterns of suburban and farm cats (*Felis catus*). *Ecography* 20, 271–280.
- Barrett, P. and Bateson, P. (1978) The development of play in cats. *Behaviour* 66, 106–120.
- Barry, K. and Crowell-Davis, S. (1999) Gender differences in the social behavior of the neutered indoor-only domestic cat. *Applied Animal Behaviour Science* 64, 193–211.
- Bateson, P. (1979) How do sensitive periods arise and what are they for? *Animal Behaviour* 27, 470–486.
- Bateson, P. (2000) Behavioural development in the cat. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*, 2nd Edn. Cambridge University Press, Cambridge, pp. 9–22.
- Bateson, P. and Bateson, M. (2002) Post-weaning feeding problems in young domestic cats – a new hypothesis. *The Veterinary Journal* 163, 113–114.
- Bateson, P. and Turner, D.C. (1988) Questions about cats. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*. Cambridge University Press, Cambridge, pp. 193–201.

- Bateson, P., Mendl, M. and Feaver, J. (1990) Play in the domestic cat is enhanced by rationing of the mother during lactation. *Animal Behaviour* 40, 514–525.
- Belkin, M., Yinon, U., Rose, L. and Reisert, I. (1977) Effect of visual environment on refractive error of cats. *Documenta Ophthalmologica* 42, 433–437.
- Berkley, M.A. (1976) Cat visual psychophysics: neural correlates and comparisons with man. *Progress in Psychobiology and Physiological Psychology* 6, 63–119.
- Bernstein, I.L. (1999) Taste aversion learning: a contemporary perspective. *Nutrition* 15, 229–234.
- Bernstein, P.L. (2007) The human–cat relationship. In: Rochlitz, I. (ed) *The Welfare of Cats*. Springer Press, Dordrecht, The Netherlands, pp. 47–89.
- Bernstein, P.L. and Strack, M. (1996) A game of cat and house: spatial patterns and behavior of 14 domestic cats (*Felis catus*) in the home. *Anthrozoos* 9, 25–39.
- Bertram, E. (2007) The relevance of kindling for human epilepsy. *Epilepsia* 48, 65–74.
- Biben, M. (1979) Predation and predatory play behaviour of domestic cats. *Animal Behaviour* 27, 81–94.
- Blakemore, C. and Van Sluyters, R.C. (1975) Innate and environmental factors in the development of the kitten's visual cortex. *Journal of Physiology* 248, 663–716.
- Bonnaud, E., Medina, F.M., Vidal, E., Nogales, M., Tershy, B., Zavaleta, E.S. *et al.* (2011) The diet of feral cats on islands: a review and a call for more studies. *Biological Invasions* 13, 581–603.
- Borchelt, P.L. and Voith, V.L. (1982) Classification of animal behaviour problems. *Veterinary Clinics of North America: Small Animal Practice* 12, 571–585.
- Boudreau, J.C. (1989) Neurophysiology and stimulus chemistry of mammalian taste systems. In: Teranishi, R., Buttery, R.G. and Shahidi, F. (eds) *Flavor Chemistry: Trends and Developments*, ACS Symposium Series 388, pp. 122–137.
- Bradshaw, J.W.S. (1986) Mere exposure reduces cats' neophobia to unfamiliar food.

Animal Behaviour 34, 613–614.

Bradshaw, J.W.S. (1991) Sensory and experiential factors in the design of foods for domestic dogs and cats. *Proceedings of the Nutrition Society* 50, 99–106.

Bradshaw, J.W.S. (2006) The evolutionary basis for the feeding behavior of domestic dogs (*Canis familiaris*) and cats (*Felis catus*). *Journal of Nutrition* 136, 1927S–1931S. Bradshaw, J.W.S. and Cameron-Beaumont, C. (2000) The signalling repertoire of the domestic cat and its undomesticated relatives. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*, 2nd Edn. Cambridge University Press, Cambridge, pp. 68–93.

Bradshaw, J.W.S. and Casey, R.A. (2007) Anthropomorphism and anthropocentrism as influences in the quality of life of companion animals. *Animal Welfare* 16(S), 149–154.

Bradshaw, J.W.S. and Cook, S. (1996) Patterns of pet cat behaviour at feeding occasions. *Applied Animal Behavior Science* 47, 61–74.

Bradshaw, J.W.S. and Hall, S.L. (1999) Affiliative behaviour of related and unrelated pairs of cats in catteries: a preliminary report. *Applied Animal Behaviour Science* 63, 251–255.

Bradshaw, J.W.S. and Limond, J. (1997) Attachment to cats and its relationship with emotional support: a cross cultural study. In: *Proceedings of the ISAZ Conference*, Boston, Massachusetts, 24–25 July.

Bradshaw, J.W.S. and Lovett, R.E. (2003) Dominance hierarchies in domestic cats: useful construct or bad habit? *Proceedings of the British Society of Animal Science Conference 2003*, 16.

Bradshaw, J.W.S., Horsfield, G.F., Allen, J.A. and Robinson, I.H. (1999) Feral cats: their role in the population dynamics of *Felis catus*. *Applied Animal Behaviour Science* 65, 273–283.

Bradshaw, J.W.S., Casey, R.A. and MacDonald, J.M. (2000a) The occurrence of unwanted behaviour in the pet cat population. *Proceedings of the Companion Animal Behaviour Therapy Study Group Study Day*, Birmingham, UK, pp. 41–42.

Bradshaw, J.W.S., Healey, L.M., Thorne, C.J., Macdonald, D.W. and Arden-Clark, C. (2000b) Differences in food preferences between individuals and populations of domestic cats *Felis silvestris catus*. *Applied Animal Behaviour Science* 68, 257–268.

- Bradshaw, J.W.S., Blackwell, E.J. and Casey, R.A. (2009) Dominance in domestic dogs – useful construct or bad habit? *Journal of Veterinary Behavior* 4, 135–144.
- Bravo, M., Blake, R. and Morrison, S. (1988) Cats see subjective contours. *Vision Research* 18, 861–865.
- Broom, D.M. (1988) The scientific assessment of animal welfare. *Applied Animal Behaviour Science* 20, 5–19.
- Broom, D.M. (2006) Behaviour and welfare in relation to pathology. *Applied Animal Behaviour Science* 97, 73–83.
- Brown, E.S. and Chandler, P.A. (2001) Mood and cognitive changes during systemic corticosteroid therapy. *Primary Care Companion. Journal of Clinical Psychiatry* 3, 17–21.
- Brown, K.A., Buchwald, J.S., Johnson, J.R. and Mikolich, D.J. (1978) Vocalization in the cat and kitten. *Developmental Psychobiology* 11, 559–570.
- Brown, S.L. (1993) The social behaviour of neutered domestic cats (*Felis catus*). PhD thesis, University of Southampton, Southampton, UK.
- Brown, S.L. and Bradshaw, J.W.S. (1996) Social behaviour in a small colony of feral cats. *Journal of the Feline Advisory Bureau* 34, 35–37.
- Buesching, C.D., Stopka, P. and Macdonald, D.W. (2003) The social function of allo-marking in the European badger (*Meles meles*). *Behaviour* 140, 965–980.
- Buffington, C.A., Chew, D.J., Kendall, M.S., Scrivani, P.V., Thompson, S.B., Blaisdell, J.L. *et al.* (1997) Clinical evaluation of cats with nonobstructive urinary tract diseases. *Journal of American Veterinary Medical Association* 210, 46–50.
- Burgess, P.R. and Perl, E.R. (1973) Cutaneous mechanoreceptors and nociceptors. In: Iggo, A. (ed.) *Handbook of Sensory Physiology*, Vol. II, The Somatosensory System. Springer-Verlag, New York, pp. 29–78.
- Cafazzo, S. and Natoli, E. (2009) The social function of tail up in the domestic cat (*Felis sylvestris catus*). *Behavioural Processes* 80, 60–66.
- Cameron, M.E., Casey, R.A., Bradshaw, J.W.S., Waran, N.K. and Gunn-Moore, D.A. (2004) A study of environmental and behavioural factors that may be associated with feline idiopathic cystitis. *Journal of Small Animal Practice* 45, 144–147.

- Cameron-Beaumont, C.L. (1997) Visual and tactile communication in the domestic cat (*Felis silvestris catus*) and undomesticated small felids. PhD thesis, University of Southampton, Southampton, UK.
- Cameron-Beaumont, C., Lowe, S.E. and Bradshaw, J.W.S. (2002) Evidence suggesting preadaptation to domestication throughout the small Felidae. *Biological Journal of the Linnean Society* 75, 361–366.
- Caro, T.M. (1980) Effects of the mother, object play, and adult experience on predation in cats. *Behavioural and Neural Biology* 29, 29–51.
- Caro, T.M. (1981) Predatory behaviour and social play in kittens. *Behaviour* 76, 1–24.
- Caro, T.M. (1989) Determinants of asociality in felids. In: Standen, V. and Foley, R.A. (eds) *Comparative Socioecology: the Behavioural Ecology of Humans and Other Mammals*. Blackwell Scientific Publications, Oxford, UK, pp. 41–74.
- Carpenter, J.A. (1956) Species differences in taste preferences. *Journal of Comparative and Physiological Psychology* 49, 139–144.
- Casey, R. (2008) Undesirable behaviours in the domestic cat: development, consequences and treatment. PhD thesis, University of Bristol, Bristol, UK.
- Casey, R.A. (2010) Fear, anxiety and conflict in companion animals. In: Lindley, S. and Watson, P. (eds) *BSAVA Manual of Canine and Feline Rehabilitation, Supportive and Palliative Care*. British Small Animal Veterinary Association, Quedgeley, UK, pp. 31–41.
- Casey, R.A. and Bradshaw, J.W.S. (2001) A comparison of referred feline clinical behaviour cases with general population prevalence data. *Scientific Proceedings of the British Small Animal Veterinary Association Congress*, Birmingham, UK, p. 529.
- Casey, R.A. and Bradshaw, J.W.S. (2005) The assessment of welfare. In: Rochlitz, I. (ed.) *The Welfare of Cats*. Springer, Dordrecht, The Netherlands, pp. 23–46.
- Casey, R.A. and Bradshaw, J.W.S. (2008) The effects of additional socialisation for kittens in a rescue centre on their behaviour and suitability as a pet. *Applied Animal Behaviour Science* 114, 196–205.
- Casey, R.A., Vandenbussche, S., Bradshaw, J.W.S. and Roberts, M.A. (2009) Reasons for relinquishment and return of domestic cats (*Felis silvestris catus*) to rescue shelters in the UK. *Anthrozoos* 22, 347–358.
- Chaadaeva, E.V. and Sokolova, N.N. (2005) Development of vocal repertoire in

- kittens of *Felis lybica* and *F.catus*. *Zoologicheskii Zhu* 84, 1402–1415.
- Cherry, K.E. and Park, D.C. (1993) Individual difference and contextual variables influence spatial memory in younger and older adults. *Psychology and Aging* 8, 517–526.
- Chesler, P. (1969) Maternal influence in learning by observation in kittens. *Science* 166, 901–903.
- Christie, L.-A., Studzinski, C.M., Araujo, J.A., Leung, C.S.K., Ikeda-Douglas, C.J., Head, E. *et al.* (2005) A comparison of egocentric and allocentric age-dependent spatial learning in the beagle dog. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 29, 361–369.
- Church, S.C., Allen, J.A. and Bradshaw, J.W.S. (1996) Frequency-dependent food selection by domestic cats: a comparative study. *Ethology* 102, 495–509.
- Clark, C.C.A., Gruffydd-Jones, T. and Murray, J.K. (2012) Number of cats and dogs in UK welfare organisations. *Veterinary Record* 170, 493.
- Clark, J.M. (1975) The effects of selection and human preference on coat colour gene frequencies in urban cats. *Heredity* 35, 195–210.
- Clifford, C.W.G. and Ibbotson, M.R. (2003) Fundamental mechanisms of visual motion detection: models, cells and functions. *Progress in Neurobiology* 68, 409–437.
- Clutton-Brock, J. (1987) *A Natural History of Domesticated Mammals*. Cambridge University Press, Cambridge, and the British Museum (Natural History), London.
- Colgan, P. (1989) *Animal Motivation*. Chapman & Hall, London.
- Collard, R.R. (1967) Fear of strangers and play behavior in kittens with varied social experience. *Child Development* 38, 877–891.
- Collier, G., Johnson, D.F. and Morgan, C. (1997) Meal patterns of cats encountering variable food procurement cost. *Journal of the Experimental Analysis of Behavior* 67, 303–310.
- Cooper, L.L. (1997) Feline inappropriate elimination. *Veterinary Clinics of North America: Small Animal Practice* 27, 569–600.
- Costalupes, J.A. (1983) Temporal integration of pure tones in the cat. *Hearing Research* 9, 43–54.
- Crouch, J.E. (1969) *Text Atlas of Cat Anatomy*. Lea and Febiger, Philadelphia, Pennsylvania.

- Crowell-Davis, S.L., Barry, K. and Wolfe, R. (1997) Social behaviour and aggressive problems of cats. *Veterinary Clinics of North America: Small Animal Practice* 27, 549–568.
- Crowell-Davis, S.L., Curtis, T. and Knowles, R. (2004) Social organization in the cat: a modern understanding. *Journal of Feline Medicine and Surgery* 6, 19–28.
- Curtis, T., Knowles, R. and Crowell-Davis, S. (2003) Influence of familiarity and relatedness on proximity and allogrooming in domestic cats (*Felis catus*). *American Journal of Veterinary Research* 64, 1151–1154.
- Dahl, J.C. (1999) A behaviour medicine approach to epilepsy – time for a paradigm shift? *Scandinavian Journal of Behaviour Therapy* 28, 97–114.
- Dantas-Divers, L.M.S., Crowell-Davis, S.L., Alford, K., Genaro, G., D’Almeida, J.M. and Paixao, R.L. (2011) Agonistic behavior and environmental enrichment of cats communally housed in a shelter. *Journal of the American Veterinary Medical Association* 239, 796–802.
- Dards, J.L. (1983) The behaviour of dockyard cats: interactions of adult males. *Applied Animal Ethology* 10, 133–153.
- Davey, G. (1989) *Ecological Learning Theory*. Routledge, London.
- de Boer, J.N. (1977a) Dominance relations in pairs of domestic cats. *Behavioural Processes* 2, 227–242.
- de Boer, J.N. (1977b) The age of olfactory cues functioning in chemocommunication among male domestic cats. *Behavioural Processes* 2, 209–225.
- Deag, J.M., Manning, A. and Lawrence, C.E. (1988) Factors influencing the mother–kitten relationship. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*. Cambridge University Press, Cambridge, pp. 23–39.
- DeAngelis, G.C. (2000) Seeing in three dimensions: the neurophysiology of stereopsis. *Trends in Cognitive Sciences* 4, 80–90.
- Deliagina, T.G., Orlovsky, G.N., Zelenin, P.V. and Beloozerova, I.N. (2006) Neural bases of postural control. *Physiology* 21, 216–225.
- Devillard, S., Say, L. and Pontier, D. (2003) Dispersal pattern of domestic cats (*Felis catus*) in a promiscuous urban population: do females disperse or die? *Journal of Animal Ecology* 72, 203–211.
- Dickman, C.R. (2009) House cats as predators in the Australian environment:

- impacts and management. *Human–Wildlife Conflicts* 3, 41–48.
- Drews, C. (1993) The concept and definition of dominance in animal behaviour. *Behaviour* 125, 283–313.
- Driscoll, C.A., Menotti-Raymond, M., Roca, A.L., Hupe, K., Johnson, W.E., Geffen, E. *et al.* (2007) The Near Eastern origin of cat domestication. *Science* 317, 519–523.
- Dukas, R. (2002) Behavioural and ecological consequences of limited attention. *Philosophical Transactions of the Royal Society of London B* 357, 1539–1547.
- Dumas, C. (1992) Object permanence in cats (*Felis catus*): an ecological approach to the study of invisible displacements. *Journal of Comparative Psychology* 106, 404–410.
- Dumas, C. (2000) Flexible search behavior in domestic cats (*Felis catus*): a case study of predator–prey interaction. *Journal of Comparative Psychology* 114, 232–238.
- Dumas, C. and Dore, F.Y. (1991) Cognitive development in kittens (*Felis catus*): an observational study of object permanence and sensorimotor intelligence. *Journal of Comparative Psychology* 105, 357–365.
- Durr, R. and Smith, C. (1997) Individual differences and their relation to social structure in domestic cats. *Journal of Comparative Psychology* 111, 412–418.
- Dybdall, K., Strasser, R. and Katz, T. (2007) Behavioral differences between owner surrender and stray domestic cats after entering an animal shelter. *Applied Animal Behaviour Science* 104, 85–94.
- Eckstein, R.A. and Hart, B.L. (2000) The organisation and control of grooming in cats. *Applied Animal Behaviour Science* 68, 131–140.
- Edney, A.T.B. (ed.) (1988) *The Waltham Book of Dog and Cat Nutrition*, 2nd Edn. Pergamon Press, Oxford, UK.
- Eisert, R. (2011) Hypercarnivory and the brain: protein requirements of cats reconsidered. *Journal of Comparative Physiology B* 181, 1–17.
- Ellis, S. (2009) Environmental enrichment: practical strategies for improving animal welfare. *Journal of Feline Medicine and Surgery* 11, 901–912.
- Elul, R. and Marchiafava, P.L. (1964) Accommodation of the eye as related to behaviour in the cat. *Archives Italienne de Biologie* 102, 616–644.
- Everett, G.M. (1944) Observations on the behavior and neurophysiology of

- acute thiamin deficient cats. *American Journal of Physiology* 141, 439–448.
- Evinger, C. and Fuchs, A.F. (1978) Saccadic, smooth pursuit and optokinetic eye movements of the trained cat. *Journal of Physiology* 285, 209–229.
- Ewer, R.F. (1973) *The Carnivores*. Weidenfield and Nicolson, London.
- Fatjó, J. and Bowen, J. (2009) Medical and metabolic influences on behavioural disorders. In: Horwitz, D.F. and Mills, D.S. (eds) *BSAVA Manual of Canine and Feline Behavioural Medicine*, 2nd Edn. British Small Animal Veterinary Association, Quedgeley, UK, pp. 1–9.
- Faure, E. and Kitchener, A.C. (2009) An archaeological and historical review of the relationship between felids and people. *Anthrozoös* 22, 221–238.
- Fay, R.R. (1988) Comparative psychoacoustics. *Hearing Research* 34, 295–306.
- Faya, M., Carranza, A., Priotto, M., Abeya, M., Diaz, J.D. and Gobello, C. (2011) Domestic queens under natural temperate photoperiod do not manifest seasonal anestrus. *Animal Reproduction Science* 129, 78–81.
- Feaver, J., Mendl, M. and Bateson, P. (1986) A method for rating the individual distinctiveness of domestic cats. *Animal Behaviour* 34, 1016–1025.
- Feldman, H. (1993) Maternal care and differences in the use of nests in the domestic cat. *Animal Behaviour* 45, 13–23.
- Feldman, H. (1994a) Domestic cats and passive submission. *Animal Behaviour* 47, 457–459.
- Feldman, H. (1994b) Methods of scent marking in the domestic cat. *Canadian Journal of Zoology* 72, 1093–1099.
- Finkler, H. and Terkel, J. (2010) Cortisol levels and aggression in neutered and intact free-roaming female cats living in urban social groups. *Physiology and Behavior* 99, 343–347.
- Finkler, H., Gunther, I. and Terkel, J. (2011) Behavioral differences between urban feeding groups of neutered and sexually intact free-roaming cats following a trap–neuter–return procedure. *Journal of the American Veterinary Medical Association* 238, 1141–1148.
- Firestein, S. (2001) How the olfactory system makes sense of scents. *Nature* 413, 211–218.
- Fiset, S. and Doré, F.Y. (1996) Spatial encoding in domestic cats (*Felis catus*). *Journal of Experimental Psychology: Animal Behavior Processes* 22, 420–437.
- Fiset, S. and Doré, F.Y. (2006) Duration of cats' (*Felis catus*) working memory

- for disappearing objects. *Animal Cognition* 9, 62–70.
- Fowler, G.A. and Sherk, H. (2003) Gaze during visually-guided locomotion in cats. *Behavioural Brain Research* 139, 83–96.
- Frank, M.C. (2011) Sleep and developmental plasticity are not just for kids. *Progress in Brain Research* 193, 221–232.
- Fraser, D. (2009) Assessing animal welfare: different philosophies, different scientific approaches. *Zoo Biology* 28, 507–518.
- Frazer Sissom, D.E., Rice, D.A. and Peters, G. (1991) How cats purr. *Journal of Zoology*, London 223, 67–78.
- Gálvez-López, E., Maes, L.D. and Abourachid, A. (2011) The search for stability on narrow supports: an experimental study in cats and dogs. *Zoology* 114, 224–232.
- Geigy, C.A., Heid, S., Steffen, F., Danielson, K., Jaggy, A. and Gaillard, C. (2007) Does a pleiotropic gene explain deafness and blue irises in white cats? *Veterinary Journal* 173, 548–553.
- German, A.J. (2006) The growing problem of obesity in dogs and cats. *Journal of Nutrition* 136, 1940S–1946S.
- Gittleman, J.L. (1991) Carnivore olfactory bulb size: allometry, phylogeny and ecology. *Journal of Zoology*, London 225, 253–272.
- Gordon, G. and Jukes, M.G.M. (1964) Dual organisation of the exteroceptive components of the cat's gracile nucleus. *Journal of Physiology* 139, 385–399.
- Gordon, J.K., Matthaei, C. and van Heezik, Y. (2010) Belled collars reduce catch of domestic cats in New Zealand by half. *Wildlife Research* 37, 372–378.
- Goulet, S., Dore, F.Y. and Rousseau, R. (1994) Object permanence and working memory in cats (*Felis catus*). *Journal of Experimental Psychology: Animal Behavior Processes* 20, 347–365.
- Grastyan, E. and Vereczkei, L. (1974) Effects of spatial separation of the conditioned signal from the reinforcement: a demonstration of the conditioned character of the orienting response or the orientational character of conditioning. *Behavioural Biology* 10, 121–146.
- Gray, J.A.B. (1966) The representation of information about rapid changes in a population of receptor units signalling mechanical events. In: de Reuck, A.V.S. and Knight, J. (eds) *Touch, Heat and Pain*. CIBA Foundation,

London, pp. 299–315.

- Grillner, S., Wallén, P., Saitoh, K., Kozlov, A. and Robertson, B. (2008) Neural bases of goal-directed locomotion in vertebrates—an overview. *Brain Research Reviews* 57, 2–12.
- Gunn-Moore, D.A. (2011) Cognitive dysfunction in cats: clinical assessment and management. *Topics in Companion Animal Medicine* 26, 17–24.
- Gunter, R. (1951) Visual size constancy in the cat. *British Journal of Psychology* 42, 288–293.
- Gunther, I., Finkler, H. and Terkel, J. (2011) Demographic differences between urban feeding groups of neutered and sexually intact free-roaming cats following a trap–neuter–return procedure. *Journal of the American Veterinary Medical Association* 238, 1134–1140.
- Guyot, G.W., Cross, H.A. and Bennett, T.L. (1980) Early social isolation of the domestic cat: responses to separation from social and nonsocial rearing stimuli. *Developmental Psychobiology* 13, 309–315.
- Hall, S.L. (1998) Object play by adult animals. In: Bekoff, M. and Byers, J.A. (eds) *Animal Play: Evolutionary, Comparative and Ecological Perspectives*. Cambridge University Press, Cambridge, pp. 45–60.
- Hall, S.L. and Bradshaw, J.W.S. (1998) The influence of hunger on object play by adult domestic cats. *Applied Animal Behaviour Science* 58, 143–150.
- Hall, S.L., Bradshaw, J.W.S. and Robinson, I.H. (2002) Object play in adult domestic cats: the roles of habituation and disinhibition. *Applied Animal Behaviour Science* 79, 263–271.
- Hart, B.L. (1974) Normal behavior and behavioral problems associated with sexual function, urination and defecation. *Veterinary Clinics of North America* 4, 589–606.
- Hart, B.L. (1979) Breed-specific behaviour. *Feline Practice* 9, 10–13.
- Hart, B.L. and Barrett, R.E. (1973) Effects of castration on fighting, roaming and urine spraying in adult male cats. *Journal of the American Veterinary Medical Association* 163, 290–292.
- Hart, B.L. and Leedy, M.G. (1987) Stimulus and hormonal determinants of flehmen behaviour in cats. *Hormones and Behaviour* 21, 44–52.
- Haskins, R. (1977) Effect of kitten vocalizations on maternal behavior. *Journal of Comparative and Physiological Psychology* 91, 830–838.
- Haskins, R. (1979) A causal analysis of kitten vocalization: an observational and

- experimental study. *Animal Behaviour* 27, 726–736.
- Heath, S.E. (2005) Behaviour problems and welfare. In: Rochlitz, I. (ed.) *The Welfare of Cats*. Springer, Dordrecht, The Netherlands, pp. 91–118.
- Heffner, H.E. (1998) Auditory awareness. *Applied Animal Behaviour Science* 57, 259–268.
- Heffner, R.S. and Heffner, H.E. (1985) Hearing range of the domestic cat. *Hearing Research* 19, 85–88.
- Hein, A. and Held, R. (1967) Dissociation of the visual placing response into elicited and guided components. *Science* 158, 390–391.
- Held, S. and Spinka, M. (2011) Animal play and animal welfare. *Animal Behaviour* 81, 891–899.
- Hendriks, W.H., Moughan, P.J., Tarttelin, M.F. and Woolhouse, A.D. (1995) Felinine: a urinary amino acid of Felidae. *Comparative Biochemistry and Physiology B* 112, 581–588.
- Herman, M.D., Denlinger, S.L., Patarca, R., Katz, L. and Hobson, J.A. (1991) Developmental phases of sleep and motor behaviour in a cat mother–infant system: a time-lapse video approach. *Canadian Journal of Psychology* 45, 101–114.
- Herron, M.E. (2010) Advances in understanding and treatment of feline inappropriate elimination. *Topics in Companion Animal Medicine* 25, 195–202.
- Hewson-Hughes, A.K., Hewson-Hughes, V.L., Miller, A.T., Hall, S.R., Simpson, S.J. and Raubenheimer, D. (2011) Geometric analysis of macronutrient selection in the adult domestic cat, *Felis catus*. *Journal of Experimental Biology* 214, 1039–1051.
- Hobhouse, L.T. (1915) *Mind in Evolution*, 2nd Edn. MacMillan, London.
- Horn, J., Mateus-Pinilla, N., Warner, R.E. and Heske, E. (2011) Home range, habitat use, and activity patterns of free-roaming domestic cats. *The Journal of Wildlife Management* 75, 1177–1185.
- Houpt, K.A. (2011) *Domestic Animal Behavior for Veterinarians and Animal Scientists*. Wiley-Blackwell, Ames, Iowa.
- Houpt, W.J. and Wolski, T.R. (1982) *Domestic Animal Behaviour for Veterinarians and Animal Scientists*. Iowa State University Press, Ames, Iowa.
- Howland, D., Bregman, B. and Glodberger, M. (1995) The development of

- quadrupedal locomotion in the kitten. *Experimental Neurobiology* 135, 93–107.
- Huang, A.Y. and May, B.J. (1996) Sound orientation behavior in cats. II. Mid-frequency spectral cues for sound localization. *Journal of the Acoustical Society of America* 100, 1070–1080.
- Huang, G.T., Rosowski, J.J. and Peake, W.T. (2000) Relating middle-ear acoustic performance to body size in the cat family: measurements and models. *Journal of Comparative Physiology A* 186, 447–465.
- Hubel, D.H. and Wiesel, T.N. (1971) Aberrant visual projections in the Siamese cat. *Journal of Physiology* 218, 33–62.
- Hudson, R., Raihani, G., Gonzalez, D., Bautista, A. and Distel, H. (2009) Nipple preference and contests in suckling kittens of the domestic cat are unrelated to presumed nipple quality. *Developmental Psychobiology* 51, 322–332.
- Hudson, R., Guarneros, M., Rodriguez, A. and Raihani, G. (2010) Emergence of personality in kittens: a qualitative approach to detecting individual behavioural differences among littermates during the first postnatal month (abstract). *Developmental Psychobiology* 52, 703.
- Hughes, A. (1972) Vergence in the cat. *Vision Research*, 12, 1961–1964.
- Hughes, A. (1977) The topography of vision in mammals of contrasting life style: comparative optics and retinal organisation. In: Crescitelli, F. (ed.) *The Visual System in Vertebrates*. Springer-Verlag, New York, pp. 613–756.
- Iggo, A. (1966) Cutaneous receptors with a high sensitivity to mechanical displacement. In: de Reuck, A.V.S. and Knight, J. (eds) *Touch, Heat and Pain*. CIBA Foundation, London, pp. 237–256.
- Iggo, A. (1982) Cutaneous sensory mechanisms. In: Barlow, H.B. and Mollon, J.D. (eds) *The Senses*. Cambridge University Press, Cambridge, pp. 369–408.
- Ishida, Y. and Shimizu, M. (1998) Influence of social rank on defecating behaviors in feral cats. *Journal of Ethology* 16, 15–21.
- Ishida, Y., Yahara, T., Kasuya, E. and Yamane, A. (2001) Female control of paternity during copulation: inbreeding avoidance in feral cats. *Behaviour* 138, 235–250.
- Jalowiec, J.E., Panksepp, J., Shabselowitz, H., Zolovick, A.J., Stern, W. and Morgane, P.J. (1973) Suppression of feeding in cats following 2-deoxy-D-glucose. *Physiology and Behavior* 10, 805–807.

- Jerison, H.J. (1985) Animal intelligence as encephalisation. *Philosophical Transactions of the Royal Society of London B* 308, 21–35.
- Jessup, D.A. (2004) The welfare of feral cats and wildlife. *Journal of the American Veterinary Medical Association* 225, 1377–1383.
- John, E.R., Chesler, P., Bartlett, F. and Victor, I. (1968) Observational learning in cats. *Science* 159, 1489–1491.
- Johnson, R.F., Randall, S. and Randall, W. (1983) Freerunning and entrained circadian rhythms in activity, eating and drinking in the cat. *Journal of Interdisciplinary Cycle Research* 14, 315–327.
- Johnson, R.W. (2002) The concept of sickness behaviour: a brief chronological account of four key discoveries. *Veterinary Immunology and Immunopathology* 87, 443–450.
- Jones, A.K., Kulkarni, B. and Derbyshire, S.W. (2003) Pain mechanisms and their disorders. *British Medical Bulletin* 65, 83–93.
- Jongman, E.C. (2007) Adaptation of domestic cats to confinement. *Journal of Veterinary Behavior* 2, 193–196.
- Kalkstein, T.S., Kruger, J.M. and Osborne, C.A. (1999) Feline lower urinary tract disease. Part I. Clinical manifestations. *Compendium of Continuing Education for Practicing Vets* 21, 15–26.
- Kane, E. (1989) Feeding behaviour of the cat. In: Burger, I.H. and Rivers, J.P.W. (eds) *Nutrition of the Dog and Cat*. Cambridge University Press, Cambridge, pp. 147–158.
- Karsh, E.B. (1983) The effects of early and late handling on the attachment of cats to people. In: Anderson, R.K., Hart, B.L. and Hart, L.A. (eds) *The Pet Connection, Conference Proceedings*. Globe Press, St Paul, Minnesota.
- Karsh, E.B. and Turner, D.C. (1988) The human–cat relationship. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*. Cambridge University Press, Cambridge, pp. 159–177.
- Kays, R. and DeWan, A. (2004) Ecological impact of inside/outside house cats around a suburban nature preserve. *Animal Conservation* 7, 273–283.
- Kerby, G. and Macdonald, D.W. (1988) Cat society and the consequences of colony size. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*. Cambridge University Press, Cambridge, pp. 67–81.
- Kessler, M.R. and Turner, D.C. (1997) Stress and adaptation of cats (*Felis*

- silvestris catus*) housed singly, in pairs and in groups in boarding catteries. *Animal Welfare* 6, 243–254.
- Kessler, M.R. and Turner, D.C. (1999) Socialization and stress in cats (*Felis silvestris catus*) housed singly and in groups in animal shelters. *Animal Welfare* 8, 15–26.
- Kiley-Worthington, M. (1976) The tail movements of ungulates, canids and felids with particular reference to their causation and function as displays. *Behaviour* 56, 69–115.
- Kiley-Worthington, M. (1984) Animal language? Vocal communication of some ungulates, canids and felids. *Acta Zoologica Fennica* 171, 83–88.
- Kimyai-Asadi, A. and Usman, A. (2001) The role of psychological stress in skin disease. *Journal of Cutaneous Medicine and Surgery* 5, 140–145.
- Kitchener, A. (1991) *The Natural History of the Wild Cats*. Christopher Helm, London.
- Knowles, R.J., Curtis, T.M. and Crowell-Davis, S.L. (2004) Correlation of dominance as determined by agonistic interactions with feeding order in cats. *American Journal of Veterinary Research* 65, 1548–1556.
- Kolb, B. and Nonneman, A.J. (1975) The development of social responsiveness in kittens. *Animal Behaviour* 23, 368–374.
- Kry, K. and Casey, R. (2007) The effect of hiding enrichment on stress levels and behaviour of domestic cats (*Felis silvestris catus*) in a shelter setting and the implications for adoption potential. *Animal Welfare* 16, 375–383.
- Kulkarni, B., Bentley, D.E., Elliot, R., Youell, P., Watson, A. and Derbyshire, S.W. (2005) Attention to pain localisation and unpleasantness discriminates the functions of the medial and lateral pain systems. *European Journal of Neuroscience* 21, 3133–3142.
- Kuo, Z.Y. (1960) Studies on the basic factors in animal fighting. VII. Interspecies coexistence in mammals. *Journal of Genetic Psychology* 97, 211–225.
- Landsberg, G., Denenberg, S. and Araujo, J. (2010) Cognitive dysfunction in cats. A syndrome we used to dismiss as ‘old age’. *Journal of Feline Medicine and Surgery* 12, 837–848.
- Lascelles, D. and Robertson, S. (2010) DJD-associated pain in cats: what can we do to promote patient comfort? *Journal of Feline Medicine and Surgery* 12, 200–212.

- Levine, M.S., Lloyd, R.L., Fisher, R.S., Hull, C.D. and Buchwald, N.A. (1987) Sensory, motor and cognitive alterations in aged cats. *Neurobiology of Aging* 8, 253–263.
- Leyhausen, P. (1979) *Cat Behavior: The Predatory and Social Behavior of Domestic and Wild Cats*. Garland STPM Press, New York.
- Li, X., Li, W., Wang, H., Cao, J., Maehashi, K., Huang, L. *et al.* (2005) Pseudogenization of a sweet-receptor gene accounts for cats' indifference toward sugar. *PLoS Genetics* 1, e3.
- Liberg, O. and Sandell, M. (1988) Spatial organisation and reproductive tactics in the domestic cat and other felids. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*. Cambridge University Press, Cambridge, pp. 83–98. Liberg, O., Sandell, M., Pontier, D. and Natoli, E. (2000) Density, spatial organization and reproductive tactics in the domestic cat and other felids. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*, 2nd Edn. Cambridge University Press, Cambridge, pp. 120–147.
- Lilith, M., Calver, M. and Garkaklis, M. (2008) Roaming habits of pet cats on the suburban fringe in Perth, Western Australia – what size buffer zone is needed to protect wildlife in reserves? *Conference Proceedings: Too Close for Comfort: Contentious Issues in Human–Wildlife Encounters*, pp. 65–72.
- Lilith, M., Calver, M. and Garkaklis, M. (2010) Do cat restrictions lead to increased species diversity or abundance of small and medium-sized mammals in remnant urban bushland? *Pacific Conservation Biology* 16, 162–172.
- Lipinski, M.J., Froenicke, L., Baysac, K.C., Billings, N.C., Leutenegger, C.M., Levy, A.M. *et al.* (2008) The ascent of cat breeds: genetic evaluations of breeds and worldwide random-bred populations. *Genomics* 91, 12–21.
- Loffler, G. (2008) Perception of contours and shapes: low and intermediate stage mechanisms. *Vision Research* 48, 2106–2127.
- Longcore, T., Rich, C. and Sullivan, L.M. (2009) Critical assessment of claims regarding management of feral cats by trap–neuter–return. *Conservation Biology* 23, 887–894.
- Loop, M.S. and Frey, T.J. (1981) Critical flicker fusion in Siamese cats. *Experimental Brain Research* 43, 65–68.
- Loop, M.S., Millican, C.L. and Thomas, S.R. (1987) Photopic spectral sensitivity of the cat. *Journal of Physiology* 382, 537–553.

- Lowe, S. and Bradshaw, J.W.S. (2001) Ontogeny of individuality in the domestic cat in the home environment. *Animal Behaviour* 61, 231–237.
- Lowe, S. and Bradshaw, J.W.S. (2002) Responses of pet cats to being held by an unfamiliar person. *Anthrozoos* 15 69–69.
- Macdonald, D.W. (1983) The ecology of carnivore social behaviour. *Nature* 301, 379–389.
- Macdonald, D.W. and Loveridge, A.J. (eds) (2010) *Biology and Conservation of Wild Felids*. Oxford University Press, Oxford.
- Macdonald, D.W., Apps, P.J., Carr, G.M. and Kerby, G. (1987) Social dynamics, nursing coalitions and infanticide among farm cats, *Felis catus*. *Advances in Ethology (supplement to Ethology)* 28, 1–64.
- Macdonald, D.W., Yamaguchi, N. and Kerby, G. (2000) Group-living in the domestic cat: its sociobiology and epidemiology. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*, 2nd Edn. Cambridge University Press, Cambridge, pp. 95–118.
- MacDonald, M.L., Rogers, Q.R. and Morris, J.G. (1985) Aversion of the cat to dietary medium-chain triglycerides and caprylic acid. *Physiology and Behavior* 35, 371–375.
- MacDonnell, M.F. and Flynn, J.P. (1966) Control of sensory fields by stimulation of hypothalamus. *Science* 152, 1406–1408.
- Malmström, T. and Kröger, R.H.H. (2006) Pupil shapes and lens optics in the eyes of terrestrial vertebrates. *Journal of Experimental Biology* 209, 18–25.
- Marchei, P., Divero, S., Falocci, Fatjo, J., Ruiz-de-la-Torre, J.L. and Manteca, X. (2009) Breed differences in behavioural development in kittens. *Physiology and Behavior* 96, 522–531.
- Martin, P. (1984) The (four) whys and wherefores of play in cats: a review of functional, evolutionary, developmental and causal issues. In: Smith, P.K. (ed.) *Play in Animals and Humans*. Blackwell, Oxford, UK, pp. 71–94.
- Martin, P. and Bateson, P. (1985) The ontogeny of locomotor play behaviour in the domestic cat. *Animal Behaviour* 33, 502–510.
- Martin, P. and Bateson, P. (1988) Behavioural development in the cat. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*. Cambridge University Press, Cambridge, pp. 9–22.
- Martin, R.L. and Webster, W.R. (1987) The auditory spatial acuity of the domestic cat in the interaural horizontal and median vertical planes.

- Hearing Research* 30, 239–252.
- Martin, R.L. and Webster, W.R. (1989) Interaural sound pressure level differences associated with sound-source locations in the frontal hemifield of the domestic cat. *Hearing Research* 38, 289–302.
- Matzner, O. and Devor, M. (1992) Na⁺ conductance and the threshold for repetitive neuronal firing. *Brain Research* 597, 92–98.
- McCann, S.M., Minura, M., Karanth, S., Yu, W.H., Mastronardi, C.A. and Rettori, V. (2000) The mechanism of action of cytokines to control the release of hypothalamic and pituitary hormones in infection. *Annals of the New York Academy of Sciences* 917, 4–18.
- McCobb, E.C., Patronek, G.J., Marder, A., Dinnage, J.D. and Stone, M.S. (2005) Assessment of stress levels among cats in four animal shelters. *Journal of the American Veterinary Medical Association* 226, 548–555.
- McComb, K., Taylor, A.M., Wilson, C. and Charlton, B.D. (2009) The cry embedded within the purr. *Current Biology* 19, R507–R508.
- McCune, S. (1992) Temperament and the welfare of caged acts. PhD thesis, University of Cambridge, Cambridge, UK.
- McCune, S. (1995) The impact of paternity and early socialization on the development of cats' behaviour to people and novel objects. *Applied Animal Behaviour Science* 45, 109–124.
- McCune, S., McPherson, J. and Bradshaw, J. (1995) Avoiding problems: the importance of socialization. In: *The Waltham Book of Human–Animal Interaction: Benefits and Responsibilities of Pet Ownership*. Pergamon/Elsevier Science, Ltd, Oxford, UK.
- McCune, S., Stevenson, J., Fretwell, L., Thompson, A. and Mills, D.S. (2008) Ageing does not significantly affect performance in a spatial learning task in the domestic cat (*Felis sylvestris catus*). *Applied Animal Behaviour Science* 112, 345–356.
- McEllistrem, J.E. (2004) Affective and predatory violence: a bimodal classification system of human aggression and violence. *Aggression and Violent Behavior* 10, 1–30.
- McFarland, D. (1985) *Animal Behaviour: Psychobiology, Ethology and Evolution*. Longman, Harlow, UK.
- McGreevy, P.D. and Boakes, R.A. (2007) *Carrots and Sticks: Principles of Animal Training*. Cambridge University Press, Cambridge, p. 318.

- McVea, D.A. and Pearson, K.G. (2007) Stepping of the forelegs over obstacles establishes longlasting memories in cats. *Current Biology* 17, R621–R623.
- Medina, F.M., Bonnaud, E., Vidal, E., Tershy B., Zavaleta, E.S., Donlan, C.J. *et al.* (2011) A global review of the impacts of invasive cats on island endangered vertebrates. *Global Change Biology* 17, 3503–3510.
- Meek, P. (2003) Home range of house cats (*Felis catus*) living within a national park. *Australian Mammalogy* 25, 51–60.
- Meier, M. and Turner, D.C. (1985) Reactions of home cats during encounters with a strange person: evidence for two personality types. *Journal of the Delta Society (later Antrozoos)* 2, 45–53.
- Mellen, J.D. (1988) The effects of hand-raising on sexual behavior of captive small felids using domestic cats as a models. *Annual Proceedings of the American Association of Zoological Parks and Aquariums*, pp. 253–259.
- Mellen, J.D. (1993) A comparative analysis of scent-marking, social and reproductive behaviour in 20 species of small cats (*Felis*). *American Zoologist* 33, 151–166.
- Mendl, M. (1988) The effects of litter-size variation on the development of play behaviour in the domestic cat: litters of one and two. *Animal Behaviour* 36, 20–34.
- Mendl, M. and Harcourt, R. (1988) Individuality in the domestic cat. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*. Cambridge University Press, Cambridge, pp. 41–54.
- Mendl, M., Brooks, J., Basse, C., Burman, O., Paul, E., Blackwell, E. *et al.* (2010) Dogs showing separation-related behaviour exhibit a ‘pessimistic’ cognitive bias. *Current Biology* 20, 839–840.
- Mendoza, D.L. and Ramirez, J.M. (1987) Play in kittens (*Felis domesticus*) and its association with cohesion and aggression. *Bulletin of the Psychonomic Society* 25, 27–30.
- Menotti-Raymond, M.A. and O’Brien, S. (1995) Evolutionary conservation of ten micro-satellite loci in four species of Felidae. *Journal of Heredity* 86, 319–322.
- Mertens, C. (1991) Human–cat interactions in the human setting. *Anthrozoos* 4, 214–231.
- Mertens, C. and Schär, R. (1988) Practical aspects of research on cats. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: the Biology of its Behaviour*. Cambridge University Press, Cambridge, pp. 179–190.

- Mertens, C. and Turner, D.C. (1988) Experimental analysis of human–cat interactions during first encounters. *Anthrozoos* 2, 83–97.
- Metsers, E., Seddon, P. and van Heezik, Y. (2010) Cat-exclusion zones in rural and urban-fringe landscapes: how large would they have to be? *Wildlife Research* 37, 47–56.
- Michael, R.P. (1961) Observations upon the sexual behaviour of the domestic cat (*Felis catus* L.) under laboratory conditions. *Behaviour* 18, 1–24.
- Miklósi, A. (2007) Human–animal interactions and social cognition in dogs. In: Jensen, P. (ed.) *The Behavioural Biology of Dogs*. CAB International, Wallingford, UK, pp. 207–222.
- Miklósi, A., Pongracz, P., Lakatos, G., Topal, J. and Csanyi, V. (2005) A comparative study of the use of visual communicative signals in interactions between dogs and humans and cats and humans. *Journal of Comparative Psychology* 119, 179–186.
- Miles, R.C. (1958) Learning in kittens with manipulatory, exploratory and food incentives. *Journal of Comparative and Physiological Psychology* 51, 39–42.
- Miyazaki, M., Yamashita, T., Taira, H. and Suzuki, A. (2008) The biological function of cauxin, a major urinary protein of the domestic cat (*Felis catus*). In: Hurst, J.L., Beynon, R.J., Roberts, S.C. and Wyatt, T. (eds) *Chemical Signals in Vertebrates II*. Springer, New York, pp. 51–60.
- Moelk, M. (1944) Vocalizing in the house-cat; a phonetic and functional study. *American Journal of Psychology* 57, 184–205.
- Morgan, S.A., Hansen, C.M., Ross, J.G., Hickling, G.J. and Ogilvie, S.C. (2009) Urban cat (*Felis catus*) movement and predation activity associated with a wetland reserve in New Zealand. *Wildlife Research* 36, 574–580.
- Morris, J.G. (2002) Idiosyncratic nutrient requirements of cats appear to be diet-induced evolutionary adaptations. *Nutrition Research Reviews* 15, 153–168.
- Mugford, R.A. (1977) External influences on the feeding of carnivores. In: Kare, M.R. and Maller, O. (eds) *The Chemical Senses and Nutrition*. Academic Press, New York, pp. 25–50.
- Mumma, R. and Warren, J.M. (1968) Two-cue discrimination learning by cats. *Journal of Comparative and Physiological Psychology* 66, 116–121.
- Murray, C.D., Flynn, J., Ratcliffe, L., Jacyna, M.R., Kamm, M.A. and Emmanuel, A.V. (2004) Acute physical and psychological stress and its

- influence on autonomic outflow to the gut in irritable bowel syndrome. *Gut* 53(Suppl. 3), A29.
- Murray, J.K., Browne, W.J., Roberts, M.A., Whitmarsh, A. and Gruffydd-Jones, T.J. (2010) Number and ownership profiles of cats and dogs in the UK. *Veterinary Record* 166, 163–168.
- Nater, U.M., Gaab, J., Rief, W. and Ehlert, U. (2006) Recent trends in behavioral medicine. *Current Opinion in Psychiatry* 19, 180–183.
- Natoli, E. (1985) Behavioural responses of urban feral cats to different types of urine marks. *Behaviour* 94, 234–243.
- Natoli, E. (1990) Mating strategies in cats: a comparison of the role and importance of infanticide in domestic cats, *Felis catus* L., and lions, *Panthera leo* L. *Animal Behaviour* 40, 183–186.
- Natoli, E. and De Vito, E. (1991) Agonistic behaviour, dominance rank and copulatory success in a large multi-male feral cat colony (*Felis catus*) in central Rome. *Animal Behaviour* 42, 227–241.
- Natoli, E., De Vito, E. and Pontier, D. (2000) Mate choice in the domestic cat (*F. catus*) *Aggressive Behavior* 26, 455–465.
- Natoli, E., Baggio, B. and Pontier, D. (2001) Male and female agonistic and affiliative relationships in a social group of farm cats (*Felis catus*). *Behavioural Processes* 53, 137–143.
- Natoli, E., Schmid, M., Say, L. and Pontier, D. (2007) Male reproductive success in a social group of urban feral cats (*Felis catus* L.). *Ethology* 113, 283–289.
- Nelson, S.H., Evans, A.D. and Bradbury, R.B. (2005) The efficacy of collar-mounted devices in reducing the rate of predation of wildlife by domestic cats. *Applied Animal Behaviour Science* 94, 273–285.
- Nicastro, N. (2004) Perceptual and acoustic evidence for species-level differences in meow vocalizations by domestic cats (*Felis catus*) and African wild cats (*Felis sylvestrus lybica*). *Journal of Comparative Psychology* 118, 287–296.
- Nicastro, N. and Owren, M.J. (2003) Classification of domestic cat (*Felis catus*) vocalizations by naive and experienced human listeners. *Journal of Comparative Psychology* 117, 44–52.
- Nieder, A. (2002) Seeing more than meets the eye: processing of illusory contours in animals. *Journal of Comparative Physiology A* 188, 249–260.

- Nogales, M., Martin, A., Tershy, B.R., Donlan, J., Veitch, D., Puerta, N. *et al.* (2004) A review of feral cat eradication on islands. *Conservation Biology* 18, 310–319.
- Norton, T.T. (1974) Receptive-field properties of superior colliculus cells and development of visual behaviour in kittens. *Journal of Neurophysiology* 37, 674–690.
- O'Brien, S.J. and Johnson, W.E. (2007) The evolution of cats. Genomic paw prints in the DNA of the world's wild cats have clarified the cat family tree and uncovered several remarkable migrations in their past. *Scientific American* 297, 68–75.
- O'Brien, S.J., Johnson, W., Driscoll, C., Pontius, J., Pecon-Slattery, J. and Menotti-Raymond, M. (2008) State of cat genomics. *Trends in Genetics* 24, 268–279.
- O'Haire, M. (2010) Companion animals and human health: benefits, challenges, and the road ahead. *Journal of Veterinary Behavior* 5, 226–234.
- Olmstead, C.E. and Villablanca, J.R. (1980) Development of behavioural audition in the kitten. *Physiology and Behaviour* 24, 705–712.
- Oswald, I. (1962) *Sleeping and Waking*. Elsevier, Amsterdam.
- Ottway, D.S. and Hawkins, D.M. (2003) Cat housing in rescue shelters: a welfare comparison between communal and discrete-unit housing. *Animal Welfare* 12, 173–177.
- Pageat, P. and Gaultier, E. (2003) Current research in canine and feline pheromones. *Veterinary Clinics of North America: Small Animal Practice* 33, 187–211.
- Palen, G.F. and Goddard, G.V. (1966) Catnip and oestrous behaviour in the cat. *Animal Behaviour* 14, 372–377.
- Passanisi, W.C. and Macdonald, D.W. (1990) Group discrimination on the basis of urine in a farm cat colony. In: Macdonald, D.W., Muller-Schwarze, D. and Natynczuk, S.E. (eds) *Chemical Signals in Vertebrates* 5. Oxford University Press, Oxford, pp. 336–345.
- Pasternak, T. and Merigan, W.H. (1980) Movement detection by cats: invariance with direction and target configuration. *Journal of Comparative and Physiological Psychology* 94, 943–952.
- Perfiliev, S., Pettersson, L.G. and Lundberg, A. (1998) Control of claw movements in cats. *Neuroscience Research* 31, 337–342.
- PETA (2011) *Inside the Fur Industry: Factory Farms*. People for the Ethical

-
- Treatment of Animals, Norfolk, Virginia. Available at: <http://www.peta.org/issues/Animals-Used-for-Clothing/inside-the-fur-industry-factory-farms.aspx>. (accessed 28 August 2012).
- Plantinga, E.A., Bosch, G. and Hendriks, W.H. (2011) Estimation of the dietary nutrient profile of free-roaming feral cats: possible implications for nutrition of domestic cats. *British Journal of Nutrition* 106, S35–S48.
- Podberscek, A.L. (2009) Good to pet and eat: the keeping and consuming of dogs and cats in South Korea. *Journal of Social Issues* 65, 615–632.
- Pontier, D. and Natoli, E. (1996) Reproductive success of male domestic cats (*Felis catus*): a case history. *Behavioural Processes* 37, 85–88.
- Pontier, D. and Natoli, E. (1999) Infanticide in rural male cats (*Felis catus*) as a reproductive mating tactic. *Aggressive Behaviour* 25, 445–449.
- Poucet, B. (1985) Spatial behaviour of cats in cue-controlled environments. *Quarterly Journal of Experimental Psychology* 37B, 155–179.
- Pozza, M.E., Stella, J.L., Chappuis-Gagnon, A.-C., Wagner, S.O. and Buffington, C.A.T. (2008) Pinch-induced behavioral inhibition ('clipnosis') in domestic cats. *Journal of Feline Medicine and Surgery* 10, 82–87.
- Pryor, P.A., Hart, B.L., Bain, M.J. and Cliff, K.D. (2001) Causes of urine marking in cats and effects of environmental management on the frequency of marking. *Journal of the American Veterinary Medical Association* 219, 1709–1713.
- Radinsky, L. (1975) Evolution of the felid brain. *Brain Behaviour & Evolution* 11, 214–254.
- Raihani, G., Gonzalez, D., Arteaga, L. and Hudson, R. (2009) Olfactory guidance of nipple attachment and suckling in kittens of the domestic cat: inborn and learned responses. *Developmental Psychobiology* 51, 662–671.
- Rainville, P. (2002) Brain mechanisms of pain affect and pain modulation. *Current Opinion in Neurobiology* 12, 195–204.
- Randall, W. and Parsons, V. (1987) Three views of the annual phase map of the domestic cat, *Felis catus* L. *Journal of Interdisciplinary Cycle Research* 18, 17–28.
- Reed, D.R., Tanaka, T. and McDaniel, A.H. (2006) Diverse tastes: genetics of sweet and bitter perception. *Physiology & Behavior* 88, 215–226.
- Reis, P.M., Jung, S., Aristoff, J.M. and Stocker, R. (2010) How cats lap: water uptake by *Felis catus*. *Science* 330, 1231–1234.

- Reisner, I. (1991) The pathophysiologic basis of behavior problems. *Veterinary Clinics of North America: Small Animal Practice* 21, 207–224.
- Reisner, I., Houpt, K., Erb, H. and Quimby, F. (1994) Friendliness to humans and defensive aggression in cats: the influence of handling and paternity. *Physiology and Behaviour* 55, 1119–1124.
- Rizzo, M.A., Kocsis, J.D. and Waxman, S.G. (1996) Mechanisms of paresthesiae, dysesthesiae and hyperesthesiae: role of Na⁺ channel heterogeneity. *European Neurology* 36, 3–12.
- Robertson, S.A. (2008) A review of feral cat control. *Journal of Feline Medicine and Surgery* 10, 366–375.
- Robertson, S.A. and Lascelles, B.D.X. (2010) Long-term pain in cats: how much do we know about this important welfare issue? *Journal of Feline Medicine and Surgery* 12, 188–199.
- Rochlitz, I. (ed.) (2005a) *The Welfare of Cats*. Springer, Dordrecht, The Netherlands. Rochlitz, I. (2005b) Housing and welfare. In: Rochlitz, I. (ed.) *The Welfare of Cats*. Springer, Dordrecht, The Netherlands, pp. 177–203.
- Romand, R. and Ehret, G. (1984) Development of sound production in normal, isolated, and deafened kittens during the first postnatal months. *Developmental Psychobiology* 17, 629–649.
- Rosenblatt, J.S. (1972) Learning in newborn kittens. *Scientific American* 227, 18–25.
- Rossignol, R., Dubuc, R. and Gossard J.-P. (2006) Dynamic sensorimotor interactions in locomotion. *Physiological Reviews* 86, 89–154.
- Rowan, A.N. and Williams, J. (1987) The success of companion animal management programs: a review. *Anthrozoos* 1, 110–122.
- Rozin, P. (1976) The selection of foods by rats, humans, and other animals. *Advances in the Study of Behaviour* 6, 21–76.
- Rupprecht, R. and Holsboer, F. (1999) Neuroactive steroids: mechanisms of action and neuropsychopharmacological perspectives. *Trends in Neurosciences* 22, 410–416.
- Rusbridge, C., Heath, S., Gunn-Moore, D.A., Knowler, S.P., Johnson, N. and McFayden, A.K. (2010) Feline orofacial pain syndrome (FOPS): a retrospective study of 113 cases. *Journal of Feline Medicine and Surgery* 12, 498–508.

- Russell, K., Sabin, R., Holt, S., Bradley, R. and Harper, E.J. (2000) Influence of feeding regimen on body condition in the cat. *Journal of Small Animal Practice* 41, 12–17.
- Ruxton, G.D., Thomas, S. and Wright, J.W. (2002) Bells reduce predation of wildlife by domestic cats (*Felis catus*). *Journal of Zoology* 256, 81–83.
- Salazar, I., Quinteiro, P.S., Cifuentes, J.M. and Caballero, T.G. (1996) The vomeronasal organ of the cat. *Journal of Anatomy* 188, 445–454.
- Salisbury, S.K. (1991) Hyperthyroidism in cats. *Compendium of Continuing Veterinary Education (Europe)* 13, 606–614.
- Sandell, M. (1989) The mating tactics and spacing patterns of solitary carnivores. In: Gittleman, J.L. (ed.) *Carnivore Behaviour, Ecology, and Evolution 1*. Cornell University Press, New York, pp. 164–182.
- Say, L. and Pontier, D. (2004) Spacing pattern in a social group of stray cats: effects on male reproductive success. *Animal Behaviour* 68, 175–180.
- Say, L., Pontier, D. and Natoli, E. (1999) High variation in multiple paternity of domestic cats in relation to environmental conditions. *Proceedings of the Royal Society of London Series B* 266, 2071–2074.
- Say, L., Pontier, D. and Natoli, E. (2001) Influence of oestrus synchronization on male reproductive success in the domestic cat (*F. catus*). *Proceedings of the Royal Society of London Series B* 268, 1049–1053.
- Say, L., Devillard, S. and Natoli, E. (2002) The mating system of feral cats in a sub-antarctic environment. *Polar Biology* 25, 838–842.
- Scherk, M. (2010) Snots and snuffles: rational approach to chronic feline upper respiratory syndromes. *Journal of Feline Medicine and Surgery* 12, 548–557.
- Schmidt, P., Lopez, R. and Collier, B. (2007) Survival, fecundity and movements of free-roaming cats. *Journal of Wildlife Management* 71, 915–919.
- Schneirla, T.C. and Rosenblatt, J.S. (1961) Behavioral organisation and genesis of the social bond in insects and mammals. *Journal of Orthopsychiatry* 31, 223–253.
- Seawright, A., Casey, R.A., Kiddie, J., Murray, J., Gruffydd-Jones, T., Harvey, A. *et al.* (2008) A case of recurrent feline idiopathic cystitis: the control of clinical signs with behaviour therapy. *Journal of Veterinary Behavior: Clinical Applications and Research* 3, 32–38.

- Serpell, J. (2000) Domestication and history of the cat. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*, 2nd Edn. Cambridge University Press, Cambridge, pp. 179–192.
- Serpell, J.A. and Paul, E.S. (2011) Pets in the family: an evolutionary perspective. In: Salmon, C.A and Shackelford, T.K. (eds) *The Oxford Handbook of Evolutionary Family Psychology*. Oxford University Press, New York, pp. 297–309.
- Shell, L.G. (1994) Feline hyperaesthesia syndrome. *Feline Practice* 22, 10.
- Sheppard, G. and Mills, D.S. (2003) Construct models in veterinary behavioural medicine: lessons from the human experience. *Veterinary Research Communications* 27, 175–191.
- Shettleworth, S.J. (1984) Natural history and evolution of learning in nonhuman mammals. In: Marler, P. and Terrace, H.S. (eds) *The Biology of Learning*. Springer-Verlag, Berlin, pp. 419–433.
- Shihab, N., Bowen, J. and Volk, H.A. (2011) Behavioural changes in dogs associated with development of idiopathic epilepsy. *Epilepsy & Behavior* 21, 160–167.
- Shimizu, M. (2001) Vocalizations of feral cats: sexual differences in the breeding season. *Mammal Study* 26, 85–92.
- Siegel, A. and Pott, C.B. (1988) Neural substrates of aggression and flight in the cat. *Progress in Neurobiology* 31, 261–283.
- Siegel, A. and Shaikh, M.B. (1997) The neural bases of aggression and rage in the cat. *Aggression and Violent Behavior* 2, 241–271.
- Siegford, J.M., Wlshaw, S.O., Brunner, P. and Zanella, A.J. (2003) Validation of a temperament test for domestic cats. *Anthrozoös* 16, 332–351.
- Silva-Rodriguez, E.A. and Sieving K.E. (2011) Influence of care of domestic carnivores on their predation on vertebrates. *Conservation Biology* 25, 808–815.
- Sims, V. and Chin, M. (2002) Responsiveness and perceived intelligence as predictors of speech addressed to cats. *Anthrozoös* 15, 166–177.
- Smith, D.F.E., Durman, K.J., Roy, D.B. and Bradshaw, J.W.S. (1994) Behavioural aspects of the welfare of rescued cats. *Journal of the Feline Advisory Bureau* 31, 25–28.
- Smithers, R.H.N. (1968) Cat of the Pharaohs. *Animal Kingdom* 61, 16–23.
- Soennichsen, S. and Chamove, A.S. (2002) Responses of cats to petting by

- humans. *Anthrozoös* 15, 258–265.
- Sokolov, E.N. (1963) Higher nervous functions: the orienting reflex. *Annual Review of Physiology* 25, 545–580.
- Sokolov, V.E., Naidenko, S.V. and Serbenyuk, M.A. (1996) Recognition by the European lynx of the species and sex and age of conspecific, familiar, and unfamiliar individuals according to urinary odors. *Izvestiya Akademii Nauk Seriya Biologicheskaya* 5, 571–577.
- Stein, B.E., Magalhaes-Castro, B. and Kruger, L. (1976) Relationship between visual and tactile representations in cat superior colliculus. *Journal of Neurophysiology* 39, 401–419.
- Sunquist, M. and Sunquist, F. (2002) *Wild Cats of the World*. University of Chicago Press, Chicago, Illinois.
- Takeuchi, Y. and Mori, Y. (2009) Behavioral profiles of feline breeds in Japan. *Journal of Veterinary Medical Science* 71, 1053–1057.
- Tami, G., Martorell, A., Torre, C., Compagnucci, M. and Manteca, X. (2011a) A model to quantify the anticipatory response in cats. *Animal Welfare* 20, 191–200.
- Tami, G., Torre, C., Compagnucci, M. and Manteca, X. (2011b) Interpretation of ambiguous spatial stimuli in cats. *Animal Welfare* 20, 185–189.
- Tamura, T., Nakatani, K. and Yau, K.W. (1989) Light adaptation in cat retinal rods. *Science* 245, 755–758.
- Tanaka, S., Tani, T., Ribot, J., O’Hashi, K. and Imamura, K. (2009) A postnatal critical period for orientation plasticity in the cat visual cortex. *PLoS One* 4, e5380.
- Thorne, C.J. (1982) Feeding behaviour in the cat – recent advances. *Journal of Small Animal Practice* 23, 555–562.
- Todd, N.B. (1977) Cats and commerce. *Scientific American* 237, 100–107.
- Tollin, D.J., McClaine, E.M. and Yin, T.C.T. (2010) Short-latency, goal-directed movements of the pinnae to sounds that produce auditory spatial illusions. *Journal of Neurophysiology* 103, 446–457.
- Tritsch, M.E. (1993) Color choice behavior in cats and the effect of changes in the color of the illuminant. *Naturwissenschaften* 80, 287–288.
- Troy, J.B. and Shou, T. (2002) The receptive fields of cat retinal ganglion cells in physiological and pathological states: where we are after half a century of research. *Progress in Retinal and Eye Research* 21, 263–302.

- Tscanz, B., Hegglin, D., Gloor, S. and Bontadina, F. (2011) Hunters and non-hunters: skewed predation rate by domestic cats in a rural village. *European Journal of Wildlife Research* 57, 597–602.
- Turner, D.C. (1988) Cat behaviour and the human/cat relationship. *Animalis Familiaris* 3, 16–21.
- Turner, D.C. (1991) The ethology of the human–cat relationship. *Swiss Archive for Veterinary Medicine* 133, 63–70.
- Turner, D. (2000) The human–cat relationship. In: Turner, D. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*, 2nd Edn. Cambridge University Press, Cambridge, pp. 193–206.
- Turner, D.C. and Bateson, P. (2000) Why the cat? In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*, 2nd Edn. Cambridge University Press, Cambridge, pp. 3–6.
- Turner, D.C. and Meister, O. (1988) Hunting behaviour of the domestic cat. In: Turner, D.C. and Bateson, P. (eds) *The Domestic Cat: The Biology of its Behaviour*. Cambridge University Press, Cambridge, pp. 111–121.
- Turner, S. (2004) Blindness in pets: seeing solutions. *Veterinary Times* 34, 8–9.
- UK Cat Behaviour Working Group (1995) *An Ethogram for Behavioural Studies of the Domestic Cat (Felis silvestris catus L)*. UFAW, Whearhampstead, UK.
- Uz, T., Akhisaroglu, M., Ahmed, R. and Manev, H. (2003) The pineal gland is critical for circadian Period1 expression in the striatum and for circadian cocaine sensitization in mice. *Neuropsychopharmacology* 28, 2117–2123.
- Van den Bos, R. (1998) The function of allogrooming in domestic cats (*Felis silvestris catus*); a study in a group of cats living in confinement. *Journal of Ethology* 16, 1–13.
- Van den Bos, R., Meijer, M.K. and Spruijt, B.M. (2000) Taste reactivity patterns in domestic cats (*Felis silvestris catus*). *Applied Animal Behaviour Science* 69, 149–168.
- Vazquez-Dominguez, E., Ceballos, G. and Cruzado, J. (2004) Extirpation of an insular subspecies by a single introduced cat: the case of the endemic deer mouse *Peromyscus guardia* on Estanque Island, Mexico. *Oryx* 38, 347–350.
- Verberne, G. and De Boer, J.N. (1976) Chemocommunication among domestic cats mediated by the olfactory and vomeronasal senses. *Zeitschrift für Tierpsychologie* 42, 86–109.

- Vigne, J.-D., Guilaine, J., Debue, K., Haye, L., and Gérard, P. (2004) Early taming of the cat in Cyprus. *Science* 304, 259.
- Villablanca, J.R. and Olmstead, C.E. (1979) Neurological development of kittens. *Developmental Psychobiology* 12, 101–127.
- Warren, J.M. (1960) Oddity learning set in a cat. *Journal of Comparative and Physiological Psychology* 53, 433–434.
- Warren, J.M. (1972) Transfer of responses to open and closed shapes in discrimination learning by cats. *Perception & Psychophysics* 12, 449–452.
- Warren, J.M. (1976) Irrelevant cues and shape discrimination learning by cats. *Animal Learning and Behaviour* 4, 22–24.
- Warren, J.M. and Beck, C.H. (1966) Visual probability learning by cats. *Journal of Comparative and Physiological Psychology* 61, 316–318.
- Watt, D.G.D. (1976) Responses of cats to sudden falls: an otolith originating reflex assisting landing. *Journal of Neurophysiology* 39, 257–265.
- Waxman, S.G., Dib-Hajj, S., Cummins, T.R. and Black, J.A. (2000) Sodium channels and their genes: dynamic expression in the normal nervous system, dysregulation in disease states. *Brain Research* 886, 5–14.
- Wedl, M., Bauer, B., Gracey, D., Grabmayer, C., Spielauer, E., Day, J. *et al.* (2011) Factors affecting the temporal patterns of dyadic behaviours and interactions between domestic cats and their owners. *Behavioural Processes* 86, 58–67.
- Weinstein, L. and Alexander, R. (2010) College students and their cats. *College Student Journal Publisher: Project Innovation (Alabama)* 44.
- Wells, D.L. and Hepper, P.G. (1992) The behaviour of dogs in a rescue shelter. *Animal Welfare* 1, 171–186.
- Wells, D.L. and Millsopp, S. (2009) Lateralised behaviour in the domestic cat, *Felis silvestris catus*. *Animal Behaviour* 78, 537–541.
- West, M.J. (1974) Social play in the domestic cat. *American Zoologist* 14, 427–436.
- West, M.J. (1979) Play in domestic kittens. In: Cairns, R.B. (ed.) *The Analysis of Social Interactions*. Hillsdale, New Jersey.
- Wetzel, M.C., Anderson, R.C., Brady, T.H. and Norgren, K.S. (1977) Kinematics of treadmill galloping by cats. III. Coordination during gait conversions and implications for neural control. *Behavioral Biology* 21, 107–127.

- White, T.D. and Boudreau, J.C. (1975) Taste preferences of the cat for neurophysiology active compounds. *Physiological Psychology* 3, 405–410.
- Whitt, E., Douglas, M., Osthaus, B. and Hocking, I. (2009) Domestic cats (*Felis catus*) do not show causal understanding in a string-pulling task. *Animal Cognition* 12, 739–743.
- Wilkinson, F. (1986) Visual texture segmentation in cats. *Behavioural Brain Research* 19, 71–82.
- Williams, C.M. and Kramer, E.M. (2010) The advantages of a tapered whisker. *PloS One* 5, e8806.
- Wilson, V.J. and Melville Jones, G. (1979) *Mammalian Vestibular Physiology*. Plenum Press, New York and London.
- Winslow, C.N. (1938) Observations of dominance-subordination in cats. *The Journal of Genetic Psychology* 52, 425–428.
- Wolfe, R. (2001) The social organization of the free-ranging domestic cat (*Felis catus*). PhD dissertation, University of Georgia, Athens, Georgia.
- Wolski, D.V.M. (1982) Social behavior of the cat. *Veterinary Clinics of North America: Small Animal Practice* 12, 693–706.
- Yamane, A. (1998) Male reproductive tactics and reproductive success of the group living cat (*Felis catus*). *Behavioural Processes* 43, 239–249.
- Yamane, A. (1999) Male homosexual mounting in the group-living feral cat (*Felis catus*). *Ethology Ecology and Evolution* 11, 399–406.
- Yeon, S.C., Kim, Y.K., Park, S.J., Lee, S.S., Lee, S.Y., Suh, E.H. *et al.* (2011) Differences between vocalization evoked by social stimuli in feral cats and house cats. *Behavioural Processes* 87, 183–189.
- Young, J.M., Massa, H.F., Hsu, L. and Trask, B.J. (2010) Extreme variability among mammalian V1R gene families. *Genome Research* 20, 10–18.
- Yu, S., Rogers, Q.R. and Morris, J.G. (1997) Absence of a salt (NaCl) preference or appetite in sodium-replete or depleted kittens. *Appetite* 29, 1–10.
- Zarzo, M. (2007) The sense of smell: molecular basis of odorant recognition. *Biological Reviews* 82, 455–479.
- Zentall, T.R. (2005) Selective and divided attention in animals. *Behavioural Processes* 69, 1–15.
- Žernicki, B. (1993) Learning deficits in lab-reared cats. *Acta Neurobiologicae Experimentalis* 53, 231–236.

Zoran, D.L. (2002) The carnivore connection to nutrition in cats. *Journal of the American Veterinary Medical Association* 221, 1559–1567.

Zoran, D.L. and Buffington, C.A. (2011) Effects of nutrition choices and lifestyle changes on the well-being of cats, a carnivore that has moved indoors. *Journal of the American Veterinary Medical Association* 239, 596–606.