

The Relationship of Bilingualism Compared to Monolingualism to the Risk of Cognitive Decline or Dementia: A Systematic Review and Meta-Analysis

Naaheed Mukadam*, Andrew Sommerlad and Gill Livingston
Division of Psychiatry, University College London, London, UK

Accepted 19 February 2017

Abstract.

Background: Bilingualism may contribute to cognitive reserve, protect against cognitive decline, and delay the onset of dementia.

Objective: We systematically reviewed evidence about the effect of bilingualism on subsequent cognitive decline or dementia.

Methods: We searched electronic databases and references for longitudinal studies comparing cognitive decline in people who were bilingual with those who were monolingual and evaluated study quality. We conducted meta-analyses using random effects models to calculate pooled odds ratio of incident dementia.

Results: We included 13/1,156 eligible articles. Meta-analysis of prospective studies of the effects of bilingualism on future dementia gave a combined Odds Ratio of dementia of 0.96 (95% CI 0.74–1.23) in bilingual participants ($n = 5,527$) compared to monolinguals. Most retrospective studies found that bilingual people were reported to develop symptoms of cognitive decline at a later age than monolingual participants.

Conclusion: We did not find that bilingualism protects from cognitive decline or dementia from prospective studies. Retrospective studies are more prone to confounding by education, or cultural differences in presentation to dementia services and are therefore not suited to establishing causative links between risk factors and outcomes.

Keywords: Bilingualism, cognitive decline, dementia, prospective cohort studies

INTRODUCTION

As the number of people with dementia continues to rise worldwide, with the accompanying social and healthcare burden [1], there is growing interest in factors that may delay or prevent the onset of cognitive decline and dementia [2]. It is recommended that people should learn multiple languages to delay the onset of dementia [3].

Cognitive reserve, defined as resilience to neuropathological damage [4], has been shown to delay

dementia onset, possibly by enhancing neural networks [5] or improving specific cognitive strategies [6]. Being fluent in two or more languages may contribute to cognitive reserve [7], and this may be a specific effect, rather than the general effect of more education, because switching languages possibly leads to an enhanced executive function rather than enhancing medial temporal memory circuitry [8]. Bilingualism, however, is complex and heterogeneous and is linked to factors, such as education, that can also affect risk of dementia [9].

Retrospective studies have found that bilingualism delays the onset of dementia by around four years [7, 10, 11]. Some prospective studies have similarly found a protective effect of learning additional

*Correspondence to: Naaheed Mukadam, MSc, Division of Psychiatry, University College London, Maple House, 149 Tottenham Court Road, London W1T 7NF, UK. Tel.: +44 02076799251; E-mail: n.mukadam@ucl.ac.uk.

languages [12–14], while others have not [15–17]. We therefore systematically reviewed the literature in this field to clarify the link between bilingualism and cognitive decline or dementia.

METHODS

Search strategy

We searched PubMed (from 1946) and Web of Knowledge (from 1900) until 23 November 2016, using search terms “bilingual*” OR “language” AND “dementia”, “AD”, “Alzheimer*” OR “cognit*”. We placed no limits on language or date of publication. We combined the search results and removed duplicates. We searched the references of included papers for further papers of interest.

Inclusion criteria

We included primary research published in peer-reviewed publications in any language which fulfilled the following three criteria:

- included people who spoke more than one language and a comparison group who did not.
- reported on cognitive function in participants not diagnosed with pre-existing neurological disorders.
- reported either a quantitative cognitive outcome measure on a validated cognitive test or incident dementia or incident mild cognitive impairment.

Exclusion criteria

- Meeting abstracts and letters.
- Comparisons between multilinguals and bilinguals with no monolingual group.

Searches and inclusion of papers

One of the authors (NM) conducted the searches and read all titles and abstracts. She read papers of studies with abstracts or titles that met inclusion criteria in full to decide whether they met inclusion criteria and discussed those which there were any questions about with the other authors.

Quality assessment

Two of the authors (NM and AS) independently read included papers and assessed their quality using

an eight-point checklist from the Newcastle-Ottawa scale for non-randomized studies [18] (see Supplementary Table 1). The questions were: Was the cohort representative of a defined population? Was the exposure (language status) accurately defined and measured? Was outcome clearly defined and measured? Have the authors adjusted for all important confounding factors? Was follow-up complete (>70%) (including death as follow-up)? Was follow-up long enough (>5 years)?

We pre-specified that we would categorize as higher quality studies those with a definition of or assessment of bilingualism, with reliable and valid cognitive outcome measures and adjusted for important confounders known to be associated with cognitive outcomes such as age, sex, education, vascular risk factors, and other potential confounders such as immigration and socio-economic status. This was to ensure that higher quality studies had valid measures of the exposure and outcome and the findings could not be accounted for by known confounders. We contacted authors for further information regarding their studies if this was not clear, in order to be able to assess quality accurately.

Analysis

If studies had multiple waves of data collection, we examined data from after the five-year follow-up. We planned to combine data from three or more studies where possible using a meta-analysis. We extracted raw data of numbers of people diagnosed with dementia in the respective bilingual and non-bilingual groups and combined unadjusted odds ratios from included studies, to calculate an overall unadjusted risk of developing dementia in bilinguals versus non-bilinguals using random effects models meta-analyses [19] with RevMan version 5.3 software. This approach is suitable for combining studies from heterogeneous populations and when different binary outcome measures are reported as it accounts for between-study variance [20].

RESULTS

The PRISMA diagram in Fig. 1 shows our search strategy results. We included 13 of 1,154 articles, reporting 13 separate studies fulfilling our criteria. Four studies were excluded after the full paper was retrieved—one because there was no record of whether or not participants spoke more than one




Fig. 1. PRISMA diagram showing search results and included studies.

134 language, [14], another because it did not include a
 135 monolingual comparator group [21]. and two because
 136 they compared bilingual participants with multilin-
 137 gual rather than monolingual participants [12, 13].
 138 Of the included studies, five were prospective and
 139 reported in Table 1 and eight were retrospective
 140 or cross-sectional and are reported in Table 2. We
 141 contacted and obtained additional information on
 142 follow-up rates and outcomes from authors of two
 143 included papers. Quality scores for each item in all
 144 studies are given in Supplementary Table 2.

145 *Prospective studies (see Table 1)*

146 The prospective studies all recruited a random
 147 sample of community-dwelling participants with-
 148 out baseline cognitive impairment [15–17, 22, 23].

Bilingualism was defined as the self-reported ability
 to communicate in two languages. One study vali-
 dated reports of bilingualism with a reading test [17]
 but used self-defined proficiency in primary analyses.
 The outcome, measured at least five years later, was
 either cognitive testing or formal diagnostic assess-
 ments. One study interviewed people in English and
 defined them as native English speakers or non-native
 English speakers with the latter group being asked
 if they spoke another language and how often they
 spoke it [23]. This study may therefore have included
 some bilingual native English speakers in the ‘mono-
 lingual’ group.

Four higher quality studies used dementia diag-
 nosis as the outcome [15–17, 23]. All of these
 studies made the diagnosis by using cognitive screen-
 ing tests then further cognitive assessment if scores

149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165

Table 1
Prospective studies

Study	Quality score	Participants and Setting; Country	N	Number of years follow-up	Follow-up rate (%)	Definition of bilingualism; comparator groups	Procedure	Baseline differences	What controlled for	Outcome
Bak et al. [22]	5	Healthy general population of people aged 11 years in 1947 – from Scottish Mental Survey; Scotland	853	50	78.1	Bilingualism self-defined as learning another language well enough to communicate in it; monolingual	Childhood intelligence at age 11 then cognitive tests 50 years later	Not stated	Childhood intelligence, age, sex, participant and father's social class	Passive/active bilingualism ↑ scores on g-factor (estimate 0.23/0.29, $p = 0.01/0.03$)
Lawton et al. [15]	5	Community dwelling Hispanic people identified from census; USA	1789	10	99.3	Bilingualism self-defined as speaking more than one language at least "very often"; monolingual	Cognitive screen. If scores low further neuropsychology testing and specialist adjudication	Bilingual participants had significantly more years of education	Immigrant status	Mean age of dementia diagnosis of bilingual participants (79.31 years) not significantly different from monolingual participants (81.10), $F(1, 77) = 1.27$, $p = 0.26$
Sanders et al. [23]	7	Community based longitudinal study of aging. Medicare recipients or registered voters sampled; USA	1779	7	91.6	Non-native English speakers (NNES) (bilingual); native monolingual English speakers (NES)	Neuro-psychological assessment at baseline; then annually. Dementia diagnosis by specialist consensus	NNES older, more likely to be white, married and immigrant, less educated, less hypertension than NES	Sex, race, years of education, immigration marital status, self-reported hypertension, diabetes, myocardial infarction, and stroke	No association between NNES status and incident dementia (HR 1.26, 95% CI 0.76–2.09; $p = 0.36$)
Yeung et al. [16]	5	Longitudinal study of Community dwelling elders, randomly selected from health care register; Canada	1468	5	67.4	Self-described: Monolinguals (56%) versus English as second language (38%) versus English bilinguals (5%)	Cognitive screening. Specialist examination & diagnosis if scored below cut off	No significant differences in age or education across all groups.	Age, sex, education, subjective memory loss at baseline	No association between language status and dementia: Adjusted OR 0.99 (95% CI 0.61, 1.59) in bilinguals versus monolinguals
Zahodne et al. [17]	7	Longitudinal aging study, from Medicare registry. No baseline cognitive impairment; USA	1067	23	80.8	All Spanish speakers. Bilinguals reported speaking English 'well' or 'very well'. Subgroup validated with reading test	Cognitive tests administered at each visit. Diagnosis by specialist consensus	Bilinguals younger, more education, more females, younger age of immigration	Age, sex, education, proportion of life spent in the U.S., country of origin, and recruitment wave	No difference in adjusted rate of dementia conversion in Cox regression: HR = 1.18 (95% CI: 0.96 – 1.46)

Table 2
Retrospective studies. MMSE, Mini-Mental State Examination; MCI, mild cognitive impairment

Study	Quality score	Setting and participants; Country	N	Definition of bilingualism; comparator group	Procedure	Baseline differences	What controlled for	Outcome
Alladi et al. [10]	3	People in memory clinic diagnosed with dementia; India	648	Self-defined ability to communicate in more than one language; monolinguals	Family members of people with dementia asked when first symptoms noticed	Bilinguals more likely to be male, have received more education, be urban dwellers	Literacy, years of education, sex, family history, vascular risk	Bilinguals onset of symptoms 4.5 years later than monolinguals unadjusted $p < 0.0001$ Adjusted analyses $F_{1,1458} = 4.89, p = 0.027$
Bialystok et al. [11]	2	People in memory clinic diagnosed with dementia; Canada	184	Most of adult life using two languages, judged by specialists based on notes; monolinguals	People with dementia and their family members asked when first symptoms noticed	Bilinguals older, less educated, lower MMSE, lower occupation, more likely to be immigrants	Age, education and occupation	Bilinguals onset of symptoms 4 years later than monolinguals, $p < 0.003$, No difference in rate of cognitive decline.
Bialystok et al. [27]	3	People in memory clinic diagnosed with dementia or MCI & no other neurological condition; Canada	149	Majority of adult life using two languages, judged by specialists; monolinguals	Patients and family members of people asked when first symptoms noticed	Bilinguals significantly less educated, more likely to be migrants, less likely to smoke and drink alcohol	Education and immigration	Bilinguals onset of MCI symptoms 4.7 years than monolinguals and 7.2 years later Alzheimer's dementia $F_{1,145} = 10.75, p = 0.001$
Chertkow et al. [29]	4	Memory clinic patients diagnosed with dementia; Canada	632	Most of adult life using two or more languages; monolinguals	Clinician consensus about age at dementia diagnosis.	No between group differences in age, years of education or initial MMSE	Sex, education, and immigrant status	No significant difference between bilingual and monolinguals' age of diagnosis or MMSE scores
Clare et al. [26]	3	Memory clinic patients or on register diagnosed with dementia and MMSE score $> 18/30$; Wales	86	Self-defined, speaking > 1 language for most of life. Also objective measure of proficiency; monolinguals	Age at time of diagnosis from clinical records.	Bilinguals less highly qualified though years education not significantly different	Education	No significant difference in age of diagnosis $F(1,79) = 2.97, p = 0.089$ or executive function scores

(Continued)

Table 2
(Continued)

Study	Quality score	Setting and participants; Country	N	Definition of bilingualism; comparator group	Procedure	Baseline differences	What controlled for	Outcome
Craik et al. [7]	2	People in memory clinic diagnosed with dementia; Canada	211	Majority of adult life using two languages, judged by specialists based on notes; monolinguals	Patients and family members of people asked when first symptoms noticed	Bilinguals older, less educated, more immigrants, lower employment status	Sex	Bilinguals onset of symptoms 5.1 years than monolinguals. Two way ANOVA ($F(1205) = 16.25$, $p < 0.0001$) with bilingualism and sex
Ossher et al. [25]	2	People referred by physician or advert responders with subjective memory complaints and at least MCI on testing; Canada	111	Majority of adult life using two languages, judged by specialists based on info in notes; monolinguals	Screened by memory tests. Those with objective memory impairment had more cognitive tests for MCI subtype	No significant differences in education or gender. No information on employment status	Nil	Bilinguals onset of amnesic MCI 4.5 years later than monolinguals ($t(66) = 2.46$, $p < 0.02$). No difference in multiple domain MCI or in duration of symptoms based on informant report
Woumans et al. [28]	2	Memory clinic patients with dementia diagnosis; Belgium	134	Self-defined at least "good" on second language and speaking it \geq once a week; monolinguals	Family members of people diagnosed with dementia asked when first symptoms noticed	No statistics given but bilinguals more educated	Sex, occupation and education	Age of onset of symptoms 4.6 years later in bilinguals ($F(1109) = 7.05$, $p = 0.009$)

were low. One of these studies calculated an odds ratio for developing dementia, adjusted for age, sex, education, and subjective memory loss [16]. The others compared mean age of dementia diagnosis [15], hazard ratio for incident dementia [23], and Cox regression on rate of dementia conversion [17], respectively. None of these studies found significant differences between bilingual and monolingual participants.

These outcomes were too heterogeneous to be combined in a meta-analysis but all the papers contained raw data of numbers of people diagnosed with dementia in the respective bilingual and non-bilingual groups. We extracted this data and conducted a meta-analysis of 5,527 participants. The meta-analysis combined unadjusted odds ratios from included studies to calculate an overall unadjusted odds ratio of developing dementia in bilinguals versus non-bilinguals of 0.96 (95% CI 0.74–1.23) (see Fig. 2), which indicates no advantage of bilingualism in protecting against dementia compared to monolingualism. In the studies included in the meta-analysis, two reported bilinguals to have received more education, one found no significant difference between education of bilinguals and monolinguals, and one reported that they received less education although reading level and therefore English proficiency, was similar in both groups.

Another, lower quality, study did not control for any confounding factors (e.g., sex, education) [22]. This study used scores on validated tests of different cognitive functions such as verbal fluency and memory. It found that those who were bilingual had higher scores on the tests of premorbid cognitive functioning, National Adult Reading Test [24] and General Fluid-Type Intelligence (G-factor), than monolinguals.

Retrospective studies (see Table 2)

The included retrospective studies were generally set in memory clinics or other specialist centers where people with memory complaints came for assessment. Trained specialists made diagnoses of dementia or mild cognitive impairment (MCI) according to validated diagnostic criteria. Most of the participants in these studies had come seeking help for cognitive complaints. One study recruited participants by advertising to the public and specifically requesting physician referrals of people with memory complaints [25] and participation required subjective memory complaints. Bilingualism was defined either

by self-report of the ability to speak two languages, or as speaking two languages for most of one's adult life. One study in this group also included an objective measure of language proficiency [26].

Five studies asked informants when they had first noticed participants' symptoms of cognitive impairment [7, 10, 11, 27, 28]. All of these studies found that bilingual participants' informants noticed symptom onset four to five years later than their monolingual counterparts. In all of these studies, bilingual participants were either more likely to be immigrants or to have had more years of education than monolingual participants.

Three studies used age of diagnosis at the clinic visit at which they were diagnosed with either all-cause dementia or MCI as the outcome [25, 26, 29]. Of these, two found no significant difference in age of diagnosis between monolinguals and bilinguals and no significant differences in years of education between the two groups [26, 29]. The third study found that age of diagnosis of amnesic MCI, was on average 4.5 years later in bilinguals than monolinguals but there was no difference in age of diagnosis for multiple domain MCI (mean difference -2.6 years, $t(41) = 1.11$; $p = 0.27$) [25]. The monolingual and bilingual participants did not differ in years of education but there was no information on their employment or immigrant status.

DISCUSSION

Our systematic review is the first to bring together all published evidence comparing cognitive decline or dementia in people who are bilingual compared to those who are monolingual. We found that, in individual prospective studies, there was no difference between bilingual and monolingual participants in the rate of development of dementia when baseline differences were taken into account. Combining these studies in this new meta-analysis has strengthened this conclusion as we found no reduction in the odds ratio of dementia in those who were bilingual compared to those who were not. By contrast, bilingual participants present around 4.5 years later in retrospective studies, where individuals' participation in the study depended on self-presentation, and time of initial symptoms are self-reported rather than standardized.

Studying the effect of an exposure, in this case bilingualism, on outcome is ideally carried out prospectively in order to reduce recall bias and clarify




Fig. 2. Forest plot showing odds ratio of developing dementia in those defined as bilingual versus those who were not.

the temporal relationship. None of the prospective studies of the development of dementia as an outcome found any protective effect of bilingualism, either individually, when adjusted for confounders, or on meta-analysis. These studies were large, examined all participants for dementia using standard methods, with good follow-up rates, controlled for confounders, had a duration of 5–10 years and measured incidence of dementia, a clinically relevant outcome. As large high-quality prospective studies have not shown an association between bilingualism and dementia, this indicates that bilingualism is not an independent protective factor.

A prospective but lower quality study (which did not control for sex or education) was not included in the meta-analysis as it measured cognitive function rather than incident dementia. It found bilingualism had a protective effect on cognition. In this study, bilingual people scored more highly on the National Adult Reading Test, which is a measure of premorbid attainment, suggesting higher cognition and education at entry; although there was no information on baseline differences in participants [22]. Thus, differences in the groups' outcomes may be due to educational or social differences rather than to bilingualism itself.

Retrospective studies in this review usually used either informant report about the date of onset of symptoms or the date of presentation to memory clinic as date of onset. This is potentially influenced by many personal and cultural factors. People from minority ethnic backgrounds tend to seek help later for dementia [30] and may define the onset of symptoms differently, potentially explaining findings of later reported symptom onset from retrospective studies that included more immigrants in the bilingual group. Although some of these studies adjusted statistically for baseline differences in education, they cannot account for cultural differences in help-seeking.

Retrospective studies that did not include a greater number of people from immigrant backgrounds in

the bilingual groups, usually included bilingual participants with higher levels of education. Education is protective against cognitive decline [31]. Although these studies have adjusted for education in their analyses, where group assignment is non-random, there is no way of determining whether associations between group membership (bilingual versus non-bilingual) and the dependent variable are due to random error or a true group difference [32]. In addition, years of education completed is not always an indicator of quality of education and the latter could be influenced by other variables such as socioeconomic status.

Two studies which did not qualify for inclusion, compared multilingual participants with bilingual participants. Both of these studies were conducted in countries where speaking multiple languages is common and participants were likely to switch between different languages many times a day. One study had a 14% follow-up rate and did not compare those who dropped out and those who did not [12]. The other was cross-sectional and found being multilingual rather than bilingual was protective, giving an odds ratio of 0.3 for cognitive impairment (95% Confidence Interval 0.10–0.92) after adjustment for education and age [13]. Both studies found that knowing and using more than two languages seems to confer a cognitive advantage and multi-lingualism may differ from bilingualism but there is not enough evidence as yet to draw definitive conclusions.

Strengths and limitations of this review

Our review was systematic and we searched using broad search terms and refined our search strategy to include as many potentially relevant papers as possible. We also hand searched references of relevant papers to identify further papers. We are therefore unlikely to have missed papers matching our inclusion criteria. We also emailed authors for missing information or clarification and this improved the accuracy of our information. Quality rating was completed using a scale which is widely used and

independently derived. The quality was rated by two authors independently. However, only one author screened titles and abstracts for inclusion. We have not carried out a funnel plot to screen for publication bias but as most of the prospective studies found negative results, publication bias is unlikely to positively skew the results. We could only carry out a meta-analysis on unadjusted odds ratios so our estimate is likely to over-estimate the effect of bilingualism.

Conclusion

We did not find evidence that bilingualism, when appropriately adjusted for education, protects from cognitive decline or dementia. Public health policy should therefore remove recommendations regarding bilingualism [3] as a strategy to delay dementia and instead concentrate on more generally reducing cognitive inactivity [33].

ACKNOWLEDGMENTS

We would like to thank Laura Zahodne and Philip Gasquoine who responded to our requests for further information.

Authors' disclosures available online (<http://j-alz.com/manuscript-disclosures/17-0131r1>).

SUPPLEMENTARY MATERIAL

The supplementary material is available in the electronic version of this article: <http://dx.doi.org/10.3233/JAD-170131>.

REFERENCES

- [1] Prince M, Wimo A, Guerchet M, Ali G-C, Wu Y-T, Prina M (2015) *World Alzheimer Report 2015: The Global Impact of Dementia*. Alzheimer's Disease International.
- [2] Norton S, Matthews FE, Barnes DE, Yaffe K, Brayne C (2014) Potential for primary prevention of Alzheimer's disease: An analysis of population-based data. *Lancet Neurol* **13**, 788-794.
- [3] European Union (2009) Study on the contribution of multilingualism to creativity.
- [4] Stern Y (2006) Cognitive reserve and Alzheimer disease. *Alzheimer Dis Assoc Disord* **20**, 112-117.
- [5] Stern Y (2012) Cognitive reserve in ageing and Alzheimer's disease. *Lancet Neurol* **11**, 1006-1012.
- [6] Amieva H, Mokri H, Le Goff M, Meillon C, Jacqmin-Gadda H, Foubert-Samier A, Orgogozo JM, Stern Y, Dartigues JF (2014) Compensatory mechanisms in higher-educated subjects with Alzheimer's disease: A study of 20 years of cognitive decline. *Brain* **137**, 1167-1175.
- [7] Craik FI, Bialystok E, Freedman M (2010) Delaying the onset of Alzheimer disease: Bilingualism as a form of cognitive reserve. *Neurology* **75**, 1726-1729.
- [8] Gold BT (2015) Lifelong bilingualism and neural reserve against Alzheimer's disease: A review of findings and potential mechanisms. *Behav Brain Res* **281**, 9-15.
- [9] Bak TH (2016) The impact of bilingualism on cognitive aging and dementia: Finding a path through a forest of confounding variables. *Linguist Approaches Biling* **6**, 205-226.
- [10] Alladi S, Bak TH, Duggirala V, Surampudi B, Shailaja M, Shukla AK, Chaudhuri JR, Kaul S (2013) Bilingualism delays age at onset of dementia, independent of education and immigration status. *Neurology* **81**, 1938-1944.
- [11] Bialystok E, Craik FI, Freedman M (2007) Bilingualism as a protection against the onset of symptoms of dementia. *Neuropsychologia* **45**, 459-464.
- [12] Kave G, Eyal N, Shorek A, Cohen-Mansfield J (2008) Multilingualism and cognitive state in the oldest old. *Psychol Aging* **23**, 70-78.
- [13] Perquin M, Vaillant M, Schuller A-M, Pastore J, Dartigues J-F, Lair M-L, Diederich N (2013) Lifelong exposure to multilingualism: New evidence to support cognitive reserve hypothesis. *PLoS One* **8**, e62030.
- [14] Wilson RS, Boyle PA, Yang J, James BD, Bennett DA (2015) Early life instruction in foreign language and music and incidence of mild cognitive impairment. *Neuropsychology* **29**, 292-302.
- [15] Lawton DM, Gasquoine PG, Weimer AA (2015) Age of dementia diagnosis in community dwelling bilingual and monolingual Hispanic Americans. *Cortex* **66**, 141-145.
- [16] Yeung CM, St John PD, Menec V, Tyas SL (2014) Is bilingualism associated with a lower risk of dementia in community-living older adults? Cross-sectional and prospective analyses. *Alzheimer Dis Assoc Disord* **28**, 326-332.
- [17] Zahodne LB, Schofield PW, Farrell MT, Stern Y, Manly JJ (2014) Bilingualism does not alter cognitive decline or dementia risk among Spanish-speaking immigrants. *Neuropsychology* **28**, 238-246.
- [18] Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P (2013) The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analyses. Available from http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp
- [19] DerSimonian R, Laird N (1986) Meta-analysis in clinical trials. *Control Clin Trials* **7**, 177-188.
- [20] Fu R, Gartlehner G, Grant M, Shamliyan T, Sedrakyan A, Wilt TJ, Griffith L, Oremus M, Raina P, Ismaila A (2011) Conducting quantitative synthesis when comparing medical interventions: AHRQ and the Effective Health Care Program. *J Clin Epidemiol* **64**, 1187-1197.
- [21] Gollan TH, Salmon DP, Montoya RI, Galasko DR (2011) Degree of bilingualism predicts age of diagnosis of Alzheimer's disease in low-education but not in highly educated Hispanics. *Neuropsychologia* **49**, 3826-3830.
- [22] Bak TH, Nissan JJ, Allerhand MM, Deary IJ (2014) Does bilingualism influence cognitive aging? *Ann Neurol* **75**, 959-963.
- [23] Sanders AE, Hall CB, Katz MJ, Lipton RB (2012) Non-native language use and risk of incident dementia in the elderly. *J Alzheimers Dis* **29**, 99-108.
- [24] Nelson H, Willison J (1991) *The revised National Adult Reading Test - test manual*. NFER-Nelson, Windsor, UK.

- 458 [25] Ossher L, Bialystok E, Craik FI, Murphy KJ, Troyer AK (2012) The effect of bilingualism on amnesic mild cognitive
459 impairment. *J Gerontol B Psychol Sci Soc Sci* **68**, 8-12. 474
- 460 [26] Clare L, Whitaker CJ, Craik FI, Bialystok E, Martyr 475
461 A, Martin-Forbes PA, Bastable AJM, Pye KL, Quinn C, 476
462 Thomas EM, Gathercole VC, Hindle JV (2014) Bilingual- 477
463 ism, executive control, and age at diagnosis among people 478
464 with early-stage Alzheimer's disease in Wales. *J Neuropsychol* **10**, 163-185. 479
- 465 [27] Bialystok E, Craik FIM, Binns MA, Ossher L, Freedman M 480
466 (2014) Effects of bilingualism on the age of onset and pro- 481
467 gression of MCI and AD: Evidence from executive function 482
468 tests. *Neuropsychology* **28**, 290-304. 483
- 469 [28] Woumans E, Santens P, Sieben A, Versijpt J, Stevens M, 484
470 Duyck W (2015) Bilingualism delays clinical manifestation 485
471 of Alzheimer's disease. *Biling Lang Cogn* **18**, 568-574. 486
472 487
- [29] Chertkow H, Whitehead V, Phillips N, Wolfson C, Atherton 474
J, Bergman H (2010) Multilingualism (but not always bilin- 475
gualism) delays the onset of Alzheimer disease: Evidence 476
from a bilingual community. *Alzheimer Dis Assoc Disord* 477
24, 118-125. 478
- [30] Mukadam N, Cooper C, Livingston G (2011) A systematic 479
review of ethnicity and pathways to care in dementia. *Int J 480
Geriatr Psychiatry* **26**, 12-20. 481
- [31] Valenzuela MJ, Sachdev P (2006) Brain reserve and demen- 482
tia: A systematic review. *Psychol Med* **36**, 441-454. 483
- [32] Miller GA, Chapman JP (2001) Misunderstanding analysis 484
of covariance. *J Abnorm Psychol* **110**, 2001-2048. 485
- [33] World Health Organization. Global plan on the public health 486
response to dementia 2017-2025. 487

Uncorrected Author Proof