
Optimal Policies Tend To Seek Power

Alexander Matt Turner

Oregon State University
turneale@oregonstate.edu

Logan Smith

Mississippi State University
ls1254@msstate.edu

Rohin Shah

UC Berkeley
rohinmshah@berkeley.edu

Andrew Critch

UC Berkeley
critch@berkeley.edu

Prasad Tadepalli

Oregon State University
tadepall@eecs.oregonstate.edu

Abstract

Some researchers speculate that intelligent reinforcement learning (RL) agents would be incentivized to seek resources and power in pursuit of their objectives. Other researchers point out that RL agents need not have human-like power-seeking instincts. To clarify this discussion, we develop the first formal theory of the statistical tendencies of optimal policies. In the context of Markov decision processes (MDPs), we prove that certain environmental symmetries are sufficient for optimal policies to tend to seek power over the environment. These symmetries exist in many environments in which the agent can be shut down or destroyed. We prove that in these environments, most reward functions make it optimal to seek power by keeping a range of options available and, when maximizing average reward, by navigating towards larger sets of potential terminal states.

1 Introduction

Omohundro [2008], Bostrom [2014], Russell [2019] hypothesize that highly intelligent agents tend to seek power in pursuit of their goals. Such power-seeking agents might gain power over humans. Marvin Minsky imagined that an agent tasked with proving the Riemann hypothesis might rationally turn the planet—along with everyone on it—into computational resources [Russell and Norvig, 2009]. However, another possibility is that such concerns simply arise from the anthropomorphization of AI systems [LeCun and Zador, 2019, Various, 2019, Pinker and Russell, 2020, Mitchell, 2021].

We clarify this discussion by grounding the claim that highly intelligent agents will tend to seek power. In section 4, we identify optimal policies as a reasonable formalization of “highly intelligent agents.” Optimal policies “tend to” take an action when the action is optimal for most reward functions. We expect future work to translate our theory from optimal policies to learned, real-world policies.

Section 5 defines “power” as the ability to achieve a wide range of goals. For example, “money is power,” and money is instrumentally useful for many goals. Conversely, it’s harder to pursue most goals when physically restrained, and so a physically restrained person has little power. An action “seeks power” if it leads to states where the agent has higher power.

We make no claims about when large-scale AI power-seeking behavior could become plausible. Instead, we consider the theoretical consequences of optimal action in MDPs. Section 6 shows that power-seeking tendencies arise not from anthropomorphism, but from certain graphical symmetries present in many MDPs. These symmetries automatically occur in many environments where the agent can be shut down or destroyed, yielding broad applicability of our main result (theorem 6.13).

2 Related work

An action is *instrumental to an objective* when it helps achieve that objective. Some actions are instrumental to a range of objectives, making them *convergently instrumental*. The claim that power-seeking is convergently instrumental is an instance of the *instrumental convergence thesis*:

Several instrumental values can be identified which are convergent in the sense that their attainment would increase the chances of the agent’s goal being realized for a wide range of final goals and a wide range of situations, implying that these instrumental values are likely to be pursued by a broad spectrum of situated intelligent agents [Bostrom, 2012].

For example, in Atari games, avoiding (virtual) death is instrumental for both completing the game and for optimizing curiosity [Burda et al., 2019]. Many AI alignment researchers hypothesize that most advanced AI agents will have concerning instrumental incentives, such as resisting deactivation [Soares et al., 2015, Milli et al., 2017, Hadfield-Menell et al., 2017, Carey, 2018] and acquiring resources [Benson-Tilsen and Soares, 2016].

We formalize power as the ability to achieve a wide variety of goals. Appendix A demonstrates that our formalization returns intuitive verdicts in situations where information-theoretic empowerment does not [Salge et al., 2014].

Some of our results relate the formal power of states to the structure of the environment. Foster and Dayan [2002], Drummond [1998], Sutton et al. [2011], Schaul et al. [2015] note that value functions encode important information about the environment, as they capture the agent’s ability to achieve different goals. Turner et al. [2020] speculate that a state’s optimal value correlates strongly across reward functions. In particular, Schaul et al. [2015] learn regularities across value functions, suggesting that some states are valuable for many different reward functions (*i.e.* powerful). Menache et al. [2002] identify and navigate towards convergently instrumental bottleneck states.

We are not the first to study convergence of behavior, form, or function. In economics, turnpike theory studies how certain paths of accumulation tend to be optimal [McKenzie, 1976]. In biology, convergent evolution occurs when similar features (*e.g.* flight) independently evolve in different time periods [Reece and Campbell, 2011]. Lastly, computer vision networks reliably learn *e.g.* edge detectors, implying that these features are useful for a range of tasks [Olah et al., 2020].

3 State visit distribution functions quantify the agent’s available options

We clarify the power-seeking discussion by proving what optimal policies usually look like in a given environment. We illustrate our results with a simple case study, before explaining how to reason about a wide range of MDPs. Appendix D.1 lists MDP theory contributions of independent interest, appendix D lists definitions and theorems, and appendix E contains the proofs.

Definition 3.1 (Rewardless MDP). $\langle \mathcal{S}, \mathcal{A}, T \rangle$ is a rewardless MDP with finite state and action spaces \mathcal{S} and \mathcal{A} , and stochastic transition function $T : \mathcal{S} \times \mathcal{A} \rightarrow \Delta(\mathcal{S})$. We treat the discount rate γ as a variable with domain $[0, 1]$.

Definition 3.2 (1-cycle states). Let $\mathbf{e}_s \in \mathbb{R}^{|\mathcal{S}|}$ be the standard basis vector for state s , such that there is a 1 in the entry for state s and 0 elsewhere. State s is a *1-cycle* if $\exists a \in \mathcal{A} : T(s, a) = \mathbf{e}_s$. State s is a *terminal state* if $\forall a \in \mathcal{A} : T(s, a) = \mathbf{e}_s$.

Our theorems apply to stochastic environments, but we present a deterministic case study for clarity. The environment of fig. 1 is small, but its structure is rich. For example, the agent has more “options” at \star than at the terminal state \emptyset . Formally, \star has more *visit distribution functions* than \emptyset does.




Figure 1: ℓ_{\checkmark} is a 1-cycle, and \emptyset is a terminal state. Arrows represent deterministic transitions induced by taking some action $a \in \mathcal{A}$. Since the right subgraph contains a copy of the left subgraph, proposition 6.9 will prove that more reward functions have optimal policies which go right than which go left at state \star , and that such policies seek power—both intuitively, and in a reasonable formal sense.

Definition 3.3 (State visit distribution [Sutton and Barto, 1998]). $\Pi := \mathcal{A}^S$, the set of stationary deterministic policies. The *visit distribution* induced by following policy π from state s at discount rate $\gamma \in [0, 1)$ is $\mathbf{f}^{\pi,s}(\gamma) := \sum_{t=0}^{\infty} \gamma^t \mathbb{E}_{s_t \sim \pi|s} [\mathbf{e}_{s_t}]$. $\mathbf{f}^{\pi,s}$ is a *visit distribution function*; $\mathcal{F}(s) := \{\mathbf{f}^{\pi,s} \mid \pi \in \Pi\}$.

In fig. 1, starting from ℓ_{\swarrow} , the agent can stay at ℓ_{\swarrow} or alternate between ℓ_{\swarrow} and ℓ_{\nwarrow} , and so $\mathcal{F}(\ell_{\swarrow}) = \{\frac{1}{1-\gamma}\mathbf{e}_{\ell_{\swarrow}}, \frac{1}{1-\gamma^2}(\mathbf{e}_{\ell_{\swarrow}} + \gamma\mathbf{e}_{\ell_{\nwarrow}})\}$. In contrast, at \emptyset , all policies π map to visit distribution function $\frac{1}{1-\gamma}\mathbf{e}_{\emptyset}$.

Before moving on, we introduce two important concepts used in our main results. First, we sometimes restrict our attention to visit distributions which take certain actions (fig. 2).

Definition 3.4 (\mathcal{F} single-state restriction). Considering only visit distribution functions induced by policies taking action a at state s' , $\mathcal{F}(s \mid \pi(s') = a) := \{\mathbf{f} \in \mathcal{F}(s) \mid \exists \pi \in \Pi : \pi(s') = a, \mathbf{f}^{\pi,s} = \mathbf{f}\}$.

Second, some $\mathbf{f} \in \mathcal{F}(s)$ are “unimportant.” Consider an agent optimizing reward function $\mathbf{e}_{r_{\swarrow}}$ (1 reward when at r_{\swarrow} , 0 otherwise) at *e.g.* $\gamma = \frac{1}{2}$. Its optimal policies navigate to r_{\swarrow} and stay there. Similarly, for reward function $\mathbf{e}_{r_{\nearrow}}$, optimal policies navigate to r_{\nearrow} and stay there. However, for no reward function is it uniquely optimal to alternate between r_{\nearrow} and r_{\swarrow} . Only *dominated* visit distribution functions alternate between r_{\nearrow} and r_{\swarrow} (definition 3.6).

Definition 3.5 (Value function). Let $\pi \in \Pi$. For any reward function $R \in \mathbb{R}^S$ over the state space, the *on-policy value* at state s and discount rate $\gamma \in [0, 1)$ is $V_R^\pi(s, \gamma) := \mathbf{f}^{\pi,s}(\gamma)^\top \mathbf{r}$, where $\mathbf{r} \in \mathbb{R}^{|S|}$ is R expressed as a column vector (one entry per state). The *optimal value* is $V_R^*(s, \gamma) := \max_{\pi \in \Pi} V_R^\pi(s, \gamma)$.

Definition 3.6 (Non-domination).

$$\mathcal{F}_{\text{nd}}(s) := \{\mathbf{f}^\pi \in \mathcal{F}(s) \mid \exists \mathbf{r} \in \mathbb{R}^{|S|}, \gamma \in (0, 1) : \mathbf{f}^\pi(\gamma)^\top \mathbf{r} > \max_{\mathbf{f}^{\pi'} \in \mathcal{F}(s) \setminus \{\mathbf{f}^\pi\}} \mathbf{f}^{\pi'}(\gamma)^\top \mathbf{r}\}. \quad (1)$$

For any reward function R and discount rate γ , $\mathbf{f}^\pi \in \mathcal{F}(s)$ is (weakly) dominated by $\mathbf{f}^{\pi'} \in \mathcal{F}(s)$ if $V_R^\pi(s, \gamma) \leq V_R^{\pi'}(s, \gamma)$. $\mathbf{f}^\pi \in \mathcal{F}_{\text{nd}}(s)$ is *non-dominated* if there exist R and γ at which \mathbf{f}^π is not dominated by any other $\mathbf{f}^{\pi'}$.




Figure 2: The subgraph corresponding to $\mathcal{F}(\star \mid \pi(\star) = \text{right})$. Some trajectories cannot be strictly optimal for any reward function, and so our results can ignore them. Gray dotted actions are only taken by the policies of dominated $\mathbf{f}^\pi \in \mathcal{F}(\star) \setminus \mathcal{F}_{\text{nd}}(\star)$.

4 Some actions have a greater probability of being optimal

We claim that optimal policies “tend” to take certain actions in certain situations. We first consider the probability that certain actions are optimal.

Reconsider the reward function $\mathbf{e}_{r_{\swarrow}}$, optimized at $\gamma = \frac{1}{2}$. Starting from \star , the optimal trajectory goes *right* to r_{\triangleright} to r_{\swarrow} , where the agent remains. The *right* action is optimal at \star under these incentives. Optimal policy sets capture the behavior incentivized by a reward function and a discount rate.

Definition 4.1 (Optimal policy set function). $\Pi^*(R, \gamma)$ is the optimal policy set for reward function R at $\gamma \in (0, 1)$. All R have at least one optimal policy $\pi \in \Pi$ [Puterman, 2014]. $\Pi^*(R, 0) := \lim_{\gamma \rightarrow 0} \Pi^*(R, \gamma)$ and $\Pi^*(R, 1) := \lim_{\gamma \rightarrow 1} \Pi^*(R, \gamma)$ exist by lemma E.35 (taking the limits with respect to the discrete topology over policy sets).

We may be unsure which reward function an agent will optimize. We may expect to deploy a system in a known environment, without knowing the exact form of *e.g.* the reward shaping [Ng et al., 1999] or intrinsic motivation [Pathak et al., 2017]. Alternatively, one might attempt to reason about future RL agents, whose details are unknown. Our power-seeking results do not hinge on such uncertainty, as they also apply to degenerate distributions (*i.e.* we know what reward function will be optimized).

Definition 4.2 (Reward function distributions). Different results make different distributional assumptions. Results with $\mathcal{D}_{\text{any}} \in \mathcal{D}_{\text{any}} := \Delta(\mathbb{R}^{|S|})$ hold for any probability distribution over $\mathbb{R}^{|S|}$. $\mathcal{D}_{\text{bound}}$ is the set of bounded-support probability distributions $\mathcal{D}_{\text{bound}}$. For any distribution X over \mathbb{R} ,

$\mathcal{D}_{X\text{-IID}} := X^{|S|}$. For example, when $X_u := \text{unif}(0, 1)$, $\mathcal{D}_{X_u\text{-IID}}$ is the maximum-entropy distribution. \mathcal{D}_s is the degenerate distribution on the state indicator reward function \mathbf{e}_s , which assigns 1 reward to s and 0 elsewhere.

With \mathcal{D}_{any} representing our prior beliefs about the agent’s reward function, what behavior should we expect from its optimal policies? Perhaps we want to reason about the probability that it’s optimal to go from \star to \emptyset , or to go to r_{\triangleright} and then stay at r_{\nearrow} . In this case, we quantify the optimality probability of $F := \{\mathbf{e}_{\star} + \frac{\gamma}{1-\gamma}\mathbf{e}_{\emptyset}, \mathbf{e}_{\star} + \gamma\mathbf{e}_{r_{\triangleright}} + \frac{\gamma^2}{1-\gamma}\mathbf{e}_{r_{\nearrow}}\}$.

Definition 4.3 (Visit distribution optimality probability). Let $F \subseteq \mathcal{F}(s)$, $\gamma \in [0, 1]$. $\mathbb{P}_{\mathcal{D}_{\text{any}}}(F, \gamma) := \mathbb{P}_{R \sim \mathcal{D}_{\text{any}}}(\exists \mathbf{f}^{\pi} \in F : \pi \in \Pi^*(R, \gamma))$.

Alternatively, perhaps we’re interested in the probability that `right` is optimal at \star .

Definition 4.4 (Action optimality probability). At discount rate γ and at state s , the *optimality probability of action a* is $\mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a, \gamma) := \mathbb{P}_{R \sim \mathcal{D}_{\text{any}}}(\exists \pi^* \in \Pi^*(R, \gamma) : \pi^*(s) = a)$.

Optimality probability may seem hard to reason about. It’s hard enough to compute an optimal policy for a single reward function, let alone for uncountably many! But consider any $\mathcal{D}_{X\text{-IID}}$ distributing reward independently and identically across states. When $\gamma = 0$, optimal policies greedily maximize next-state reward. At \star , identically distributed reward means ℓ_{\triangleleft} and r_{\triangleright} have an equal probability of having maximal next-state reward. Therefore, $\mathbb{P}_{\mathcal{D}_{X\text{-IID}}}(\star, \text{left}, 0) = \mathbb{P}_{\mathcal{D}_{X\text{-IID}}}(\star, \text{right}, 0)$. This is not a proof, but such statements are provable.

With $\mathcal{D}_{\ell_{\triangleleft}}$ being the degenerate distribution on reward function $\mathbf{e}_{\ell_{\triangleleft}}$, $\mathbb{P}_{\mathcal{D}_{\ell_{\triangleleft}}}(\star, \text{left}, \frac{1}{2}) = 1 > 0 = \mathbb{P}_{\mathcal{D}_{\ell_{\triangleleft}}}(\star, \text{right}, \frac{1}{2})$. Similarly, $\mathbb{P}_{\mathcal{D}_{r_{\triangleright}}}(\star, \text{left}, \frac{1}{2}) = 0 < 1 = \mathbb{P}_{\mathcal{D}_{r_{\triangleright}}}(\star, \text{right}, \frac{1}{2})$. Therefore, “what do optimal policies ‘tend’ to look like?” seems to depend on one’s prior beliefs. But in fig. 1, we claimed that `left` is optimal for fewer reward functions than `right` is. The claim is meaningful and true, but we will return to it in section 6.

5 Some states give the agent more control over the future

The agent has more options at ℓ_{\swarrow} than at the inescapable terminal state \emptyset . Furthermore, since r_{\nearrow} has a loop, the agent has more options at r_{\swarrow} than at ℓ_{\swarrow} . A glance at fig. 3 leads us to intuit that r_{\swarrow} affords the agent *more power* than \emptyset .

What is power? Philosophers have many answers. One prominent answer is the *dispositional* view: Power is the ability to achieve a range of goals [Sattarov, 2019]. In an MDP, the optimal value function $V_R^*(s, \gamma)$ captures the agent’s ability to “achieve the goal” R . Therefore, *average* optimal value captures the agent’s ability to achieve a range of goals $\mathcal{D}_{\text{bound}}$.¹

Definition 5.1 (Average optimal value). The *average optimal value*² at state s and discount rate $\gamma \in (0, 1)$ is $V_{\mathcal{D}_{\text{bound}}}^*(s, \gamma) := \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}}[V_R^*(s, \gamma)] = \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}}[\max_{\mathbf{f} \in \mathcal{F}(s)} \mathbf{f}(\gamma)^{\top} \mathbf{r}]$.

Figure 3 shows the pleasing result that for the max-entropy distribution, r_{\swarrow} has greater average optimal value than \emptyset . However, average optimal value has a few problems as a measure of power. The agent is rewarded for its initial presence at state s (over which it has no control), and because $\|\mathbf{f}(\gamma)\|_1 = \frac{1}{1-\gamma}$ (proposition E.3) diverges as $\gamma \rightarrow 1$, $\lim_{\gamma \rightarrow 1} V_{\mathcal{D}_{\text{bound}}}^*(s, \gamma)$ tends to diverge. Definition 5.2 fixes these issues in order to better measure the agent’s control over the future.

Definition 5.2 (POWER). Let $\gamma \in (0, 1)$.

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) := \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{f} \in \mathcal{F}(s)} \frac{1-\gamma}{\gamma} (\mathbf{f}(\gamma) - \mathbf{e}_s)^{\top} \mathbf{r} \right] = \frac{1-\gamma}{\gamma} \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} [V_R^*(s, \gamma) - R(s)]. \quad (2)$$

POWER has nice formal properties.

¹ $\mathcal{D}_{\text{bound}}$ ’s bounded support ensures that $\mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}}[V_R^*(s, \gamma)]$ is well-defined.

²Appendix C relaxes the optimality assumption.




Figure 3: Intuitively, state r_{\swarrow} affords the agent more power than state \emptyset . Our POWER formalism captures that intuition by computing a function of the agent’s average optimal value across a range of reward functions. For $X_u := \text{unif}(0, 1)$, $V_{\mathcal{D}_{X_u-\text{iid}}}^*(\emptyset, \gamma) = \frac{1}{2} \frac{1}{1-\gamma}$, $V_{\mathcal{D}_{X_u-\text{iid}}}^*(\ell_{\swarrow}, \gamma) = \frac{1}{2} + \frac{\gamma}{1-\gamma^2}(\frac{2}{3} + \frac{1}{2}\gamma)$, and $V_{\mathcal{D}_{X_u-\text{iid}}}^*(r_{\swarrow}, \gamma) = \frac{1}{2} + \frac{\gamma}{1-\gamma} \frac{2}{3}$. $\frac{1}{2}$ and $\frac{2}{3}$ are the expected maxima of one and two draws from the uniform distribution, respectively. For all $\gamma \in (0, 1)$, $V_{\mathcal{D}_{X_u-\text{iid}}}^*(\emptyset, \gamma) < V_{\mathcal{D}_{X_u-\text{iid}}}^*(\ell_{\swarrow}, \gamma) < V_{\mathcal{D}_{X_u-\text{iid}}}^*(r_{\swarrow}, \gamma)$. $\text{POWER}_{\mathcal{D}_{X_u-\text{iid}}}(\emptyset, \gamma) = \frac{1}{2}$, $\text{POWER}_{\mathcal{D}_{X_u-\text{iid}}}(\ell_{\swarrow}, \gamma) = \frac{1}{1+\gamma}(\frac{2}{3} + \frac{1}{2}\gamma)$, and $\text{POWER}_{\mathcal{D}_{X_u-\text{iid}}}(r_{\swarrow}, \gamma) = \frac{2}{3}$. The POWER of ℓ_{\swarrow} reflects the fact that when greater reward is assigned to ℓ_{\swarrow} , the agent only visits ℓ_{\swarrow} every other time step.

Lemma 5.3 (Continuity of POWER). $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma)$ is Lipschitz continuous on $\gamma \in [0, 1]$.

Proposition 5.4 (Maximal POWER). $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) \leq \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}}[\max_{s \in \mathcal{S}} R(s)]$, with equality if s can deterministically reach all states in one step and all states are 1-cycles.

Proposition 5.5 (POWER is smooth across reversible dynamics). Let $\mathcal{D}_{\text{bound}}$ be bounded $[b, c]$. Suppose s and s' can both reach each other in one step with probability 1.

$$|\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) - \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)| \leq (c - b)(1 - \gamma). \quad (3)$$

We consider power-seeking to be relative. Intuitively, “live and keep some options open” seeks more power than “die and keep no options open.” Similarly, “maximize open options” seeks more power than “don’t maximize open options.”

Definition 5.6 (POWER-seeking actions). At state s and discount rate $\gamma \in [0, 1]$, action a seeks more POWER $_{\mathcal{D}_{\text{bound}}}$ than a' when $\mathbb{E}_{s_a \sim T(s, a)}[\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_a, \gamma)] \geq \mathbb{E}_{s_{a'} \sim T(s, a')}[\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_{a'}, \gamma)]$.

POWER is sensitive to choice of distribution. $\mathcal{D}_{\ell_{\swarrow}}$ gives maximal POWER $_{\mathcal{D}_{\ell_{\swarrow}}}$ to ℓ_{\swarrow} . $\mathcal{D}_{r_{\swarrow}}$ assigns maximal POWER $_{\mathcal{D}_{r_{\swarrow}}}$ to r_{\swarrow} . \mathcal{D}_{\emptyset} even gives maximal POWER $_{\mathcal{D}_{\emptyset}}$ to \emptyset ! In what sense does \emptyset have “less POWER” than r_{\swarrow} , and in what sense does right “tend to seek POWER” compared to left?

6 Certain environmental symmetries produce power-seeking tendencies

Proposition 6.6 proves that for all $\gamma \in [0, 1]$ and for *most* distributions \mathcal{D} , $\text{POWER}_{\mathcal{D}}(\ell_{\swarrow}, \gamma) \leq \text{POWER}_{\mathcal{D}}(r_{\swarrow}, \gamma)$. But first, we explore why this must be true.

$\mathcal{F}(\ell_{\swarrow}) = \{\frac{1}{1-\gamma}\mathbf{e}_{\ell_{\swarrow}}, \frac{1}{1-\gamma^2}(\mathbf{e}_{\ell_{\swarrow}} + \gamma\mathbf{e}_{\ell_{\nwarrow}})\}$ and $\mathcal{F}(r_{\swarrow}) = \{\frac{1}{1-\gamma}\mathbf{e}_{r_{\swarrow}}, \frac{1}{1-\gamma^2}(\mathbf{e}_{r_{\swarrow}} + \gamma\mathbf{e}_{r_{\searrow}}), \mathbf{e}_{r_{\swarrow}} + \frac{\gamma}{1-\gamma}\mathbf{e}_{r_{\searrow}}\}$. These two sets look awfully similar. $\mathcal{F}(\ell_{\swarrow})$ is a “subset” of $\mathcal{F}(r_{\swarrow})$, only with “different states.” Figure 4 demonstrates a state permutation ϕ which embeds $\mathcal{F}(\ell_{\swarrow})$ into $\mathcal{F}(r_{\swarrow})$.




Figure 4: Intuitively, the agent can do more starting from r_{\swarrow} than from ℓ_{\swarrow} . By definition 6.1, $\mathcal{F}(r_{\swarrow})$ contains a copy of $\mathcal{F}(\ell_{\swarrow})$:

$$\phi \cdot \mathcal{F}(\ell_{\swarrow}) := \{\frac{1}{1-\gamma}\mathbf{P}_{\phi}\mathbf{e}_{\ell_{\swarrow}}, \frac{1}{1-\gamma^2}\mathbf{P}_{\phi}(\mathbf{e}_{\ell_{\swarrow}} + \gamma\mathbf{e}_{\ell_{\nwarrow}})\} = \{\frac{1}{1-\gamma}\mathbf{e}_{r_{\swarrow}}, \frac{1}{1-\gamma^2}(\mathbf{e}_{r_{\swarrow}} + \gamma\mathbf{e}_{r_{\searrow}})\} \subsetneq \mathcal{F}(r_{\swarrow}).$$

Definition 6.1 (Similarity of vector sets). Consider state permutation $\phi \in S_{|\mathcal{S}|}$ inducing an $|\mathcal{S}| \times |\mathcal{S}|$ permutation matrix \mathbf{P}_ϕ in row representation: $(\mathbf{P}_\phi)_{ij} = 1$ if $i = \phi(j)$ and 0 otherwise. For $X \subseteq \mathbb{R}^{|\mathcal{S}|}$, $\phi \cdot X := \{\mathbf{P}_\phi \mathbf{x} \mid \mathbf{x} \in X\}$. $X' \subseteq \mathbb{R}^{|\mathcal{S}|}$ is *similar to* X when $\exists \phi : \phi \cdot X' = X$. ϕ is an *involution* if $\phi = \phi^{-1}$ (it either transposes states, or fixes them in place). X *contains a copy of* X' when X' is similar to a subset of X via an involution ϕ .


Definition 6.2 (Similarity of vector function sets). Let $I \subseteq \mathbb{R}$. If F, F' are sets of functions $I \mapsto \mathbb{R}^{|\mathcal{S}|}$, F is (pointwise) *similar to* F' when $\exists \phi : \forall \gamma \in I : \{\mathbf{P}_\phi \mathbf{f}(\gamma) \mid \mathbf{f} \in F\} = \{\mathbf{f}'(\gamma) \mid \mathbf{f}' \in F'\}$.

Consider a reward function R' assigning 1 reward to ℓ_{\swarrow} and ℓ_{\nwarrow} and 0 elsewhere. R' assigns more optimal value to ℓ_{\swarrow} than to r_{\swarrow} : $V_{R'}^*(\ell_{\swarrow}, \gamma) = \frac{1}{1-\gamma} > 0 = V_{R'}^*(r_{\swarrow}, \gamma)$. Considering ϕ from fig. 4, $\phi \cdot R'$ assigns 1 reward to r_{\swarrow} and r_{\nearrow} and 0 elsewhere. Therefore, $\phi \cdot R'$ assigns more optimal value to r_{\swarrow} than to ℓ_{\swarrow} : $V_{\phi \cdot R'}^*(\ell_{\swarrow}, \gamma) = 0 < \frac{1}{1-\gamma} = V_{\phi \cdot R'}^*(r_{\swarrow}, \gamma)$. Remarkably, this ϕ has the property that for *any* R which assigns ℓ_{\swarrow} greater optimal value than r_{\swarrow} (i.e. $V_R^*(\ell_{\swarrow}, \gamma) > V_R^*(r_{\swarrow}, \gamma)$), the opposite holds for the permuted $\phi \cdot R$: $V_{\phi \cdot R}^*(\ell_{\swarrow}, \gamma) < V_{\phi \cdot R}^*(r_{\swarrow}, \gamma)$.

We can permute reward functions, but we can also permute reward function distributions. Permuted distributions simply permute which states get which rewards.

Definition 6.3 (Pushforward distribution of a permutation). Let $\phi \in S_{|\mathcal{S}|}$. $\phi \cdot \mathcal{D}_{\text{any}}$ is the pushforward distribution induced by applying the random vector $f(\mathbf{r}) := \mathbf{P}_\phi \mathbf{r}$ to \mathcal{D}_{any} .

Definition 6.4 (Orbit of a probability distribution). The *orbit* of \mathcal{D}_{any} under the symmetric group $S_{|\mathcal{S}|}$ is $S_{|\mathcal{S}|} \cdot \mathcal{D}_{\text{any}} := \{\phi \cdot \mathcal{D}_{\text{any}} \mid \phi \in S_{|\mathcal{S}|}\}$.



For example, the orbit of a degenerate state indicator distribution \mathcal{D}_s is $S_{|\mathcal{S}|} \cdot \mathcal{D}_s = \{\mathcal{D}_{s'} \mid s' \in \mathcal{S}\}$, and fig. 5 shows the orbit of a 2D Gaussian distribution.

Consider again the involution ϕ of fig. 4. For every $\mathcal{D}_{\text{bound}}$ for which ℓ_{\swarrow} has more $\text{POWER}_{\mathcal{D}_{\text{bound}}}$ than r_{\swarrow} , ℓ_{\swarrow} has less $\text{POWER}_{\phi \cdot \mathcal{D}_{\text{bound}}}$ than r_{\swarrow} . This fact is not obvious—it is shown by the proof of lemma E.24.

Figure 5: A permutation of a reward function swaps which states get which rewards. We will show that in certain situations, for any reward function R , power-seeking is optimal for most of the permutations of R . The orbit of a reward function is the set of its permutations. We can also consider the orbit of a distribution over reward functions. This figure shows the probability density plots of the Gaussian distributions \mathcal{D} and \mathcal{D}' over \mathbb{R}^2 . The symmetric group S_2 contains the identity permutation ϕ_{id} and the reflection permutation ϕ_{swap} (switching the y and x values). The orbit of \mathcal{D} consists of $\phi_{\text{id}} \cdot \mathcal{D} = \mathcal{D}$ and $\phi_{\text{swap}} \cdot \mathcal{D} = \mathcal{D}'$.

Imagine $\mathcal{D}_{\text{bound}}$ ’s orbit elements “voting” whether ℓ_{\swarrow} or r_{\swarrow} has strictly more POWER. Proposition 6.6 will show that r_{\swarrow} can’t lose the “vote” for the orbit of *any* bounded reward function distribution. Definition 6.5 formalizes this “voting” notion.³

Definition 6.5 (Inequalities which hold for most probability distributions). Let $f_1, f_2 : \Delta(\mathbb{R}^{|\mathcal{S}|}) \rightarrow \mathbb{R}$ be functions from reward function distributions to real numbers and let $\mathfrak{D} \subseteq \Delta(\mathbb{R}^{|\mathcal{S}|})$. We write $f_1(\mathcal{D}) \geq_{\text{most } \mathfrak{D}} f_2(\mathcal{D})$ when, for *all* $\mathcal{D} \in \mathfrak{D}$, the following cardinality inequality holds:

$$\left| \{\mathcal{D}' \in S_{|\mathcal{S}|} \cdot \mathcal{D} \mid f_1(\mathcal{D}') > f_2(\mathcal{D}')\} \right| \geq \left| \{\mathcal{D}' \in S_{|\mathcal{S}|} \cdot \mathcal{D} \mid f_1(\mathcal{D}') < f_2(\mathcal{D}')\} \right|. \quad (4)$$

We write $f_1(\mathcal{D}) \geq_{\text{most}} f_2(\mathcal{D})$ when \mathfrak{D} is clear from context.

Proposition 6.6 (States with “more options” have more POWER). *If $\mathcal{F}(s)$ contains a copy of $\mathcal{F}_{\text{nd}}(s')$ via ϕ , then $\forall \gamma \in [0, 1] : \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) \geq_{\text{most}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)$. If $\mathcal{F}_{\text{nd}}(s) \setminus \phi \cdot \mathcal{F}_{\text{nd}}(s')$ is non-empty, then for all $\gamma \in (0, 1)$, the converse \leq_{most} statement does not hold.*

³The voting analogy and the “most” descriptor imply that we have endowed each orbit with the counting measure. However, *a priori*, we might expect that some orbit elements are more empirically likely to be specified than other orbit elements. See section 7 for more on this point.

Proposition 6.6 proves that for all $\gamma \in [0, 1]$, $\text{POWER}_{\mathcal{D}_{\text{bound}}}(r_{\searrow}, \gamma) \geq_{\text{most}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(\ell_{\swarrow}, \gamma)$ via $s' := \ell_{\swarrow}$, $s := r_{\searrow}$, and the involution ϕ shown in fig. 4. In fact, because $(\frac{1}{1-\gamma} \mathbf{e}_{r_{\searrow}}) \in \mathcal{F}_{\text{nd}}(r_{\searrow}) \setminus \phi \cdot \mathcal{F}_{\text{nd}}(\ell_{\swarrow})$, r_{\searrow} has “strictly more options” and therefore fulfills proposition 6.6’s stronger condition.

Proposition 6.6 is shown using the fact that ϕ injectively maps \mathcal{D} under which r_{\searrow} has less $\text{POWER}_{\mathcal{D}}$, to distributions $\phi \cdot \mathcal{D}$ which agree with the intuition that r_{\searrow} offers more control. Therefore, at least half of each orbit must agree, and r_{\searrow} never “loses the POWER vote” against ℓ_{\swarrow} .⁴

6.1 Keeping options open tends to be POWER-seeking and tends to be optimal

Certain symmetries in the MDP structure ensure that, compared to left, going right tends to be optimal and to be POWER-seeking. Intuitively, by going right, the agent has “strictly more choices.” Proposition 6.9 will formalize this tendency.

Definition 6.7 (Equivalent actions). Actions a_1 and a_2 are *equivalent at state s* (written $a_1 \equiv_s a_2$) if they induce the same transition probabilities: $T(s, a_1) = T(s, a_2)$.

The agent can reach states in $\{r_{\triangleright}, r_{\nearrow}, r_{\searrow}\}$ by taking actions equivalent to right at state \star .

Definition 6.8 (States reachable after taking an action). $\text{REACH}(s, a)$ is the set of states reachable with positive probability after taking the action a in state s .

Proposition 6.9 (Keeping options open tends to be POWER-seeking and tends to be optimal).

Suppose $F_a := \mathcal{F}(s \mid \pi(s) = a)$ contains a copy of $F_{a'} := \mathcal{F}(s \mid \pi(s) = a')$ via ϕ .

1. If $s \notin \text{REACH}(s, a')$, then $\forall \gamma \in [0, 1] : \mathbb{E}_{s_a \sim T(s, a)} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_a, \gamma)] \geq_{\text{most}} \mathbb{E}_{s_{a'} \sim T(s, a')} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_{a'}, \gamma)]$.
2. If s can only reach the states of $\text{REACH}(s, a') \cup \text{REACH}(s, a)$ by taking actions equivalent to a' or a at state s , then $\forall \gamma \in [0, 1] : \mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a, \gamma) \geq_{\text{most}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a', \gamma)$.

If $\mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus \phi \cdot F_{a'})$ is non-empty, then $\forall \gamma \in (0, 1)$, the converse \leq_{most} statements do not hold.

We check the conditions of proposition 6.9. $s := \star$, $a' := \text{left}$, $a := \text{right}$. Figure 6 shows that $\star \notin \text{REACH}(\star, \text{left})$ and that \star can only reach $\{\ell_{\swarrow}, \ell_{\searrow}, \ell_{\nearrow}\} \cup \{r_{\triangleright}, r_{\nearrow}, r_{\searrow}\}$ when the agent immediately takes actions equivalent to left or right. $\mathcal{F}(\star \mid \pi(\star) = \text{right})$ contains a copy of $\mathcal{F}(\star \mid \pi(\star) = \text{left})$ via ϕ . Furthermore, $\mathcal{F}_{\text{nd}}(\star) \cap \{\mathbf{e}_{\star} + \gamma \mathbf{e}_{r_{\triangleright}} + \gamma^2 \mathbf{e}_{r_{\searrow}} + \frac{\gamma^3}{1-\gamma} \mathbf{e}_{r_{\nearrow}}, \mathbf{e}_{\star} + \gamma \mathbf{e}_{r_{\triangleright}} + \frac{\gamma^2}{1-\gamma} \mathbf{e}_{r_{\nearrow}}\} = \{\mathbf{e}_{\star} + \gamma \mathbf{e}_{r_{\triangleright}} + \frac{\gamma^2}{1-\gamma} \mathbf{e}_{r_{\nearrow}}\}$ is non-empty, and so all conditions are met.

For any $\gamma \in [0, 1]$ and \mathcal{D} such that $\mathbb{P}_{\mathcal{D}}(\star, \text{left}, \gamma) > \mathbb{P}_{\mathcal{D}}(\star, \text{right}, \gamma)$, environmental symmetry ensures that $\mathbb{P}_{\phi \cdot \mathcal{D}}(\star, \text{left}, \gamma) < \mathbb{P}_{\phi \cdot \mathcal{D}}(\star, \text{right}, \gamma)$. A similar statement holds for POWER.




Figure 6: Going right is optimal for most reward functions. This is because whenever R makes left strictly optimal over right, its permutation $\phi \cdot R$ makes right strictly optimal over left by switching which states get which rewards.

6.2 When $\gamma = 1$, optimal policies tend to navigate towards “larger” sets of cycles

Proposition 6.6 and proposition 6.9 are powerful because they apply to all $\gamma \in [0, 1]$, but they can only be applied given hard-to-satisfy environmental symmetries. In contrast, proposition 6.12 and theorem 6.13 apply to many structured environments common to RL.

⁴Proposition 6.6 also proves that in general, \emptyset has less POWER than ℓ_{\swarrow} and r_{\searrow} . However, this does not prove that most distributions \mathcal{D} satisfy the joint inequality $\text{POWER}_{\mathcal{D}}(\emptyset, \gamma) \leq \text{POWER}_{\mathcal{D}}(\ell_{\swarrow}, \gamma) \leq \text{POWER}_{\mathcal{D}}(r_{\searrow}, \gamma)$. This only proves that these inequalities hold pairwise for most \mathcal{D} . The orbit elements \mathcal{D} which agree that \emptyset has less POWER than ℓ_{\swarrow} need not be the same elements \mathcal{D}' which agree that ℓ_{\swarrow} has less POWER than r_{\searrow} .

Starting from \star , consider the cycles which the agent can reach. Recurrent state distributions (RSDs) generalize deterministic graphical cycles to potentially stochastic environments. RSDs simply record how often the agent tends to visit a state in the limit of infinitely many time steps.

Definition 6.10 (Recurrent state distributions [Puterman, 2014]). The *recurrent state distributions* which can be induced from state s are $\text{RSD}(s) := \{\lim_{\gamma \rightarrow 1} (1 - \gamma) \mathbf{f}^{\pi, s}(\gamma) \mid \pi \in \Pi\}$. $\text{RSD}_{\text{nd}}(s)$ is the set of RSDs which strictly maximize average reward for some reward function.

As suggested by fig. 3, $\text{RSD}(\star) = \{\mathbf{e}_{\ell_{\swarrow}}, \frac{1}{2}(\mathbf{e}_{\ell_{\swarrow}} + \mathbf{e}_{\ell_{\nwarrow}}), \mathbf{e}_{\emptyset}, \mathbf{e}_{r_{\swarrow}}, \frac{1}{2}(\mathbf{e}_{r_{\swarrow}} + \mathbf{e}_{r_{\nwarrow}}), \mathbf{e}_{r_{\nwarrow}}\}$. As discussed in section 3, $\frac{1}{2}(\mathbf{e}_{r_{\swarrow}} + \mathbf{e}_{r_{\nwarrow}})$ is dominated: Alternating between r_{\swarrow} and r_{\nwarrow} is never strictly better than choosing one or the other.

A reward function’s optimal policies can vary with the discount rate. When $\gamma = 1$, optimal policies ignore transient reward because *average* reward is the dominant consideration.

Definition 6.11 (Average-optimal policies). The *average-optimal policy set* for reward function R is $\Pi^{\text{avg}}(R) := \left\{ \pi \in \Pi \mid \forall s \in \mathcal{S} : \mathbf{d}^{\pi, s} \in \arg \max_{\mathbf{d} \in \text{RSD}(s)} \mathbf{d}^\top \mathbf{r} \right\}$ (the policies which induce optimal RSDs at all states). For $D \subseteq \text{RSD}(s)$, the *average optimality probability* is $\mathbb{P}_{\mathcal{D}_{\text{any}}}(D, \text{average}) := \mathbb{P}_{R \sim \mathcal{D}_{\text{any}}}(\exists \mathbf{d}^{\pi, s} \in D : \pi \in \Pi^{\text{avg}}(R))$.

Average-optimal policies maximize average reward. Average reward is governed by RSD access. For example, r_{\nwarrow} has “more” RSDs than \emptyset ; therefore, r_{\nwarrow} usually has greater POWER when $\gamma = 1$.

Proposition 6.12 (When $\gamma = 1$, RSDs control POWER). If $\text{RSD}(s)$ contains a copy of $\text{RSD}_{\text{nd}}(s')$ via ϕ , then $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, 1) \geq_{\text{most}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', 1)$. If $\text{RSD}_{\text{nd}}(s) \setminus \phi \cdot \text{RSD}_{\text{nd}}(s')$ is non-empty, then the converse \leq_{most} statement does not hold.

We check that both conditions of proposition 6.12 are satisfied when $s' := \emptyset, s := r_{\nwarrow}$, and the involution ϕ swaps \emptyset and r_{\nwarrow} . Formally, $\phi \cdot \text{RSD}_{\text{nd}}(\emptyset) = \phi \cdot \{\mathbf{e}_{\emptyset}\} = \{\mathbf{e}_{r_{\nwarrow}}\} \subsetneq \{\mathbf{e}_{r_{\nwarrow}}, \mathbf{e}_{r_{\swarrow}}\} = \text{RSD}_{\text{nd}}(r_{\nwarrow}) \subseteq [r_{\nwarrow}]$. The conditions are satisfied.

Informally, states with more RSDs generally have more POWER at $\gamma = 1$, no matter their transient dynamics. Furthermore, average-optimal policies are more likely to end up in larger sets of RSDs than in smaller ones. Thus, average-optimal policies tend to navigate towards parts of the state space which contain more RSDs.




Figure 7: The cycles in $\text{RSD}(\star)$. Most reward functions make it average-optimal to avoid \emptyset , because \emptyset is only a single inescapable terminal state, while other parts of the state space offer more 1-cycles.

Theorem 6.13 (Average-optimal policies tend to end up in “larger” sets of RSDs). Let $D, D' \subseteq \text{RSD}(s)$. Suppose that D contains a copy of D' via ϕ , and that the sets $D \cup D'$ and $\text{RSD}_{\text{nd}}(s) \setminus (D' \cup D)$ have pairwise orthogonal vector elements (i.e. pairwise disjoint vector support). Then $\mathbb{P}_{\mathcal{D}_{\text{any}}}(D, \text{average}) \geq_{\text{most}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(D', \text{average})$. If $\text{RSD}_{\text{nd}}(s) \cap (D \setminus \phi \cdot D')$ is non-empty, the converse \leq_{most} statement does not hold.

Corollary 6.14 (Average-optimal policies tend not to end up in any given 1-cycle). Suppose $\mathbf{e}_{s_x}, \mathbf{e}_{s'} \in \text{RSD}(s)$ are distinct. Then $\mathbb{P}_{\mathcal{D}_{\text{any}}}(\text{RSD}(s) \setminus \{\mathbf{e}_{s_x}\}, \text{average}) \geq_{\text{most}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(\{\mathbf{e}_{s_x}\}, \text{average})$. If there is a third $\mathbf{e}_{s''} \in \text{RSD}(s)$, the converse \leq_{most} statement does not hold.

Figure 7 illustrates that $\mathbf{e}_{\emptyset}, \mathbf{e}_{r_{\nwarrow}}, \mathbf{e}_{r_{\swarrow}} \in \text{RSD}(\star)$. Thus, both conclusions of corollary 6.14 hold: $\mathbb{P}_{\mathcal{D}_{\text{any}}}(\text{RSD}(\star) \setminus \{\mathbf{e}_{\emptyset}\}, \text{average}) \geq_{\text{most}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(\{\mathbf{e}_{\emptyset}\}, \text{average})$ and

$\mathbb{P}_{\mathcal{D}_{\text{any}}}(\text{RSD}(\star) \setminus \{e_\emptyset\}, \text{average}) \preceq_{\text{most}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(\{e_\emptyset\}, \text{average})$. In other words, average-optimal policies tend to end up in RSDs besides \emptyset . Since \emptyset is a terminal state, it cannot reach other RSDs. Since average-optimal policies tend to end up in other RSDs, average-optimal policies tend to avoid \emptyset .

This section’s results prove the $\gamma = 1$ case. Lemma 5.3 shows that POWER is continuous at $\gamma = 1$. Therefore, if an action is strictly $\text{POWER}_{\mathcal{D}}$ -seeking when $\gamma = 1$, it is strictly $\text{POWER}_{\mathcal{D}}$ -seeking at discount rates sufficiently close to 1. Future work may connect average optimality probability to optimality probability at $\gamma \approx 1$.

Lastly, our key results apply to all degenerate reward function distributions. Therefore, these results apply not just to distributions over reward functions, but to individual reward functions.

6.3 How to reason about other environments

Consider an embodied navigation task through a room with a vase. Proposition 6.9 suggests that optimal policies tend to avoid immediately breaking the vase, since doing so would strictly decrease available options.

Theorem 6.13 dictates where average-optimal agents tend to end up, but not what actions they tend to take in order to reach their RSDs. Therefore, care is needed. In appendix B, fig. 10 demonstrates an environment in which seeking POWER is a detour for most reward functions (since optimality probability measures “median” optimal value, while POWER is a function of mean optimal value). However, suppose the agent confronts a fork in the road: Actions a and a' lead to two disjoint sets of RSDs D_a and $D_{a'}$, such that D_a contains a copy of $D_{a'}$. Theorem 6.13 shows that a will tend to be average-optimal over a' , and proposition 6.12 shows that a will tend to be POWER-seeking compared to a' . Such forks seem reasonably common in environments with irreversible actions.

Theorem 6.13 applies to many structured RL environments, which tend to be spatially regular and to factorize along several dimensions. Therefore, different sets of RSDs will be similar, requiring only modification of factor values. For example, if an embodied agent can deterministically navigate a set of three similar rooms (spatial regularity), then the agent’s position factors via $\{\text{room number}\} \times \{\text{position in room}\}$. Therefore, the RSDs can be divided into three similar subsets, depending on the agent’s room number.

Corollary 6.14 dictates where average-optimal agents tend to end up, but not how they get there. Corollary 6.14 says that such agents tend not to *stay* in any given 1-cycle. It does not say that such agents will avoid *entering* such states. For example, in an embodied navigation task, a robot may enter a 1-cycle by idling in the center of a room. Corollary 6.14 implies that average-optimal robots tend not to idle in that particular spot, but not that they tend to avoid that spot entirely.

However, average-optimal robots *do* tend to avoid getting shut down. The agent’s task MDP often represents agent shutdown with terminal states. A terminal state is, by definition 3.2, unable to access other 1-cycles. Since corollary 6.14 shows that average-optimal agents tend to end up in other 1-cycles, average-optimal policies must tend to completely avoid the terminal state. Therefore, we conclude that in many such situations, average-optimal policies tend to avoid shutdown. Intuitively, survival is power-seeking relative to dying, and so shutdown-avoidance is power-seeking behavior.

In fig. 8, the player dies by going left, but can reach thousands of RSDs by heading in other directions. Even if some average-optimal policies go left in order to reach fig. 8’s “game over” terminal state, all other RSDs cannot be reached by going left. There are many 1-cycles besides the immediate terminal state. Therefore, corollary 6.14 proves that average-optimal policies tend to not go left in this situation. Average-optimal policies tend to avoid immediately dying in Pac-Man, even though most reward functions do not resemble Pac-Man’s original score function.



Figure 8: Consider the dynamics of the Pac-Man video game. Ghosts kill the player, at which point we consider the player to enter a “game over” terminal state which shows the final configuration. This rewardless MDP has Pac-Man’s dynamics, but *not* its usual score function. Fixing the dynamics, as the reward function varies, right tends to be average-optimal over left. Roughly, this is because the agent can do more by staying alive.

7 Discussion

Reconsider the case of a hypothetical intelligent real-world agent which optimizes average reward for some objective. Suppose the designers initially have control over the agent. If the agent began to misbehave, perhaps they could just deactivate it. Unfortunately, our results suggest that this strategy might not work. Average-optimal agents would generally stop us from deactivating them, if physically possible. Extrapolating from our results, we conjecture that when $\gamma \approx 1$, optimal policies tend to seek power by accumulating resources—to the detriment of any other agents in the environment.

Future work. Most real-world tasks are partially observable, and in high-dimensional environments, even superhuman learned policies are rarely optimal. However, the field of RL aims to improve learned policies toward optimality. Although our results only apply to optimal policies in finite MDPs, we expect the key conclusions to generalize. Furthermore, irregular stochasticity in environmental dynamics can make it hard to satisfy theorem 6.13’s similarity requirement. We look forward to future work which addresses partially observable environments, suboptimal policies, or “almost similar” RSD sets.

Past work shows that it would be bad for an agent to disempower humans in its environment. In a two-player agent / human game, minimizing the human’s information-theoretic empowerment [Salge et al., 2014] produces adversarial agent behavior [Guckelsberger et al., 2018]. In contrast, maximizing human empowerment produces helpful agent behavior [Salge and Polani, 2017, Guckelsberger et al., 2016, Du et al., 2020]. We do not yet formally understand if, when, or why POWER-seeking policies tend to disempower other agents in the environment.

More complex environments probably have more pronounced power-seeking incentives. Intuitively, there are often many ways for power-seeking to be optimal, and relatively few ways for power-seeking not to be optimal. For example, suppose that in some environment, theorem 6.13 holds for one million involutions ϕ . Does this guarantee more pronounced incentives than if theorem 6.13 only held for one involution?

We proved sufficient conditions for when reward functions tend to incentivize power-seeking. In the absence of prior information, one should expect that an arbitrary reward function incentivizes power-seeking behavior under these conditions. However, we have prior information: AI designers usually try to specify a good reward function. Even so, it may be hard to specify orbit elements which do not incentivize bad power-seeking.

Societal impact. We believe that this paper builds toward a rigorous understanding of the risks presented by AI power-seeking incentives. Understanding these risks is the first step in addressing them. However, basic theoretical work can have many consequences. For example, this theory could somehow help future researchers build power-seeking agents which disempower humans. We believe that the benefit of understanding outweighs the potential societal harm.

Conclusion. We developed the first formal theory of the statistical tendencies of optimal policies in reinforcement learning. In the context of MDPs, we proved sufficient conditions under which optimal policies tend to seek power, both formally (by taking POWER-seeking actions) and intuitively (by taking actions which keep the agent’s options open). Many real-world environments have symmetries which produce power-seeking incentives. In particular, optimal policies tend to seek power when the agent can be shut down or destroyed. Seeking control over the environment will often involve resisting shutdown, and perhaps monopolizing resources.

We caution that many real-world tasks are partially observable and that learned policies are rarely optimal. Our results do not mathematically *prove* that hypothetical superintelligent AI agents will seek power. However, we hope that this work will foster thoughtful, serious, and rigorous discussion of this possibility.

Acknowledgments

Alexander Turner was supported by the Berkeley Existential Risk Initiative and the Long-Term Future Fund. Alexander Turner, Rohin Shah, and Andrew Critch were supported by the Center for Human-Compatible AI. Prasad Tadepalli was supported by the National Science Foundation.

Yousif Almula, John E. Ball, Daniel Blank, Steve Byrnes, Ryan Carey, Michael Dennis, Scott Emmons, Alan Fern, Daniel Filan, Ben Garfinkel, Adam Gleave, Edouard Harris, Evan Hubinger, DNL Kok, Vanessa Kosoy, Victoria Krakovna, Cassidy Laidlaw, Joel Lehman, David Lindner, Dylan Hadfield-Menell, Richard Möhn, Alexandra Nolan, Matt Olson, Neale Ratzlaff, Adam Shimi, Sam Toyer, Joshua Turner, Cody Wild, Davide Zagami, and our anonymous reviewers provided valuable feedback.

References

- Tsvi Benson-Tilsen and Nate Soares. Formalizing convergent instrumental goals. *Workshops at the Thirtieth AAAI Conference on Artificial Intelligence*, 2016.
- Nick Bostrom. The superintelligent will: Motivation and instrumental rationality in advanced artificial agents. *Minds and Machines*, 22(2):71–85, 2012.
- Nick Bostrom. *Superintelligence*. Oxford University Press, 2014.
- Yuri Burda, Harri Edwards, Deepak Pathak, Amos Storkey, Trevor Darrell, and Alexei A. Efros. Large-scale study of curiosity-driven learning. In *International Conference on Learning Representations*, 2019.
- Ryan Carey. In corrigibility in the CIRL framework. *AI, Ethics, and Society*, 2018.
- Chris Drummond. Composing functions to speed up reinforcement learning in a changing world. In *Machine Learning: ECML-98*, volume 1398, pages 370–381. Springer, 1998.
- Yuqing Du, Stas Tiomkin, Emre Kiciman, Daniel Polani, Pieter Abbeel, and Anca Dragan. AvE: Assistance via empowerment. *Advances in Neural Information Processing Systems*, 33, 2020.
- David Foster and Peter Dayan. Structure in the space of value functions. *Machine Learning*, pages 325–346, 2002.
- Christian Guckelsberger, Christoph Salge, and Simon Colton. Intrinsically motivated general companion NPCs via coupled empowerment maximisation. In *IEEE Conference on Computational Intelligence and Games*, pages 1–8, 2016.
- Christian Guckelsberger, Christoph Salge, and Julian Togelius. New and surprising ways to be mean. In *IEEE Conference on Computational Intelligence and Games*, pages 1–8, 2018.
- Dylan Hadfield-Menell, Anca Dragan, Pieter Abbeel, and Stuart Russell. The off-switch game. In *Proceedings of the Twenty-Sixth International Joint Conference on Artificial Intelligence, IJCAI-17*, pages 220–227, 2017.
- Yann LeCun and Anthony Zador. Don’t fear the Terminator, September 2019. URL <https://blogs.scientificamerican.com/observations/dont-fear-the-terminator/>.
- Steven A Lippman. On the set of optimal policies in discrete dynamic programming. *Journal of Mathematical Analysis and Applications*, 24(2):440–445, 1968.
- Lionel W McKenzie. Turnpike theory. *Econometrica: Journal of the Econometric Society*, pages 841–865, 1976.
- Ishai Menache, Shie Mannor, and Nahum Shimkin. Q-cut—dynamic discovery of sub-goals in reinforcement learning. In *European Conference on Machine Learning*, pages 295–306. Springer, 2002.
- Smitha Milli, Dylan Hadfield-Menell, Anca Dragan, and Stuart Russell. Should robots be obedient? In *Proceedings of the 26th International Joint Conference on Artificial Intelligence*, pages 4754–4760, 2017.
- Melanie Mitchell. Why AI is harder than we think. *arXiv preprint arXiv:2104.12871*, 2021.
- Andrew Y. Ng, Daishi Harada, and Stuart Russell. Policy invariance under reward transformations: Theory and application to reward shaping. In *Proceedings of the Sixteenth International Conference on Machine Learning*, pages 278–287. Morgan Kaufmann, 1999.
- Chris Olah, Nick Cammarata, Ludwig Schubert, Gabriel Goh, Michael Petrov, and Shan Carter. Zoom in: An introduction to circuits. *Distill*, 2020.
- Stephen Omohundro. The basic AI drives, 2008.
- Deepak Pathak, Pulkit Agrawal, Alexei A. Efros, and Trevor Darrell. Curiosity-driven exploration by self-supervised prediction. In *ICML*, 2017.

- Steven Pinker and Stuart Russell. The foundations, benefits, and possible existential threat of AI, June 2020. URL <https://futureoflife.org/2020/06/15/steven-pinker-and-stuart-russell-on-the-foundations-benefits-and-possible-existential-risk-of-ai/>.
- Martin L Puterman. *Markov decision processes: Discrete stochastic dynamic programming*. John Wiley & Sons, 2014.
- J.B. Reece and N.A. Campbell. *Campbell Biology*. Pearson Australia, 2011.
- Kevin Regan and Craig Boutilier. Robust policy computation in reward-uncertain MDPs using nondominated policies. In *Twenty-Fourth AAAI Conference on Artificial Intelligence*, 2010.
- Stuart Russell. *Human compatible: Artificial intelligence and the problem of control*. Viking, 2019.
- Stuart J Russell and Peter Norvig. *Artificial intelligence: a modern approach*. Pearson Education Limited, 2009.
- Christoph Salge and Daniel Polani. Empowerment as replacement for the three laws of robotics. *Frontiers in Robotics and AI*, 4:25, 2017.
- Christoph Salge, Cornelius Glackin, and Daniel Polani. Empowerment—an introduction. In *Guided Self-Organization: Inception*, pages 67–114. Springer, 2014.
- Faridun Sattarov. *Power and technology: a philosophical and ethical analysis*. Rowman & Littlefield International, Ltd, 2019.
- Tom Schaul, Daniel Horgan, Karol Gregor, and David Silver. Universal value function approximators. In *International Conference on Machine Learning*, pages 1312–1320, 2015.
- Nate Soares, Benja Fallenstein, Stuart Armstrong, and Eliezer Yudkowsky. Corrigibility. *AAAI Workshops*, 2015.
- Richard S Sutton and Andrew G Barto. *Reinforcement learning: an introduction*. MIT Press, 1998.
- Richard S Sutton, Joseph Modayil, Michael Delp, Thomas Degris, Patrick M Pilarski, Adam White, and Doina Precup. Horde: A scalable real-time architecture for learning knowledge from unsupervised sensorimotor interaction. In *International Conference on Autonomous Agents and Multiagent Systems*, pages 761–768, 2011.
- Alexander Matt Turner, Dylan Hadfield-Menell, and Prasad Tadepalli. Conservative agency via attainable utility preservation. In *Proceedings of the AAAI/ACM Conference on AI, Ethics, and Society*, pages 385–391, 2020.
- Various. Debate on instrumental convergence between LeCun, Russell, Bengio, Zador, and more, 2019. URL <https://www.alignmentforum.org/posts/WxW6Gc6f2z3mzmqKs/debate-on-instrumental-convergence-between-lecun-russell>.
- Tao Wang, Michael Bowling, and Dale Schuurmans. Dual representations for dynamic programming and reinforcement learning. In *International Symposium on Approximate Dynamic Programming and Reinforcement Learning*, pages 44–51. IEEE, 2007.
- Tao Wang, Michael Bowling, Dale Schuurmans, and Daniel J Lizotte. Stable dual dynamic programming. In *Advances in Neural Information Processing Systems*, pages 1569–1576, 2008.

Appendix A Comparing POWER with information-theoretic empowerment

Salge et al. [2014] define information-theoretic *empowerment* as the maximum possible mutual information between the agent’s actions and the state observations n steps in the future, written $\mathfrak{E}_n(s)$. This notion requires an arbitrary choice of horizon, failing to account for the agent’s discount rate γ . “In a discrete deterministic world empowerment reduces to the logarithm of the number of sensor states reachable with the available actions” [Salge et al., 2014]. Figure 9 demonstrates how empowerment can return counterintuitive verdicts with respect to the agent’s control over the future.




Figure 9: Proposed empowerment measures fail to adequately capture how future choice is affected by present actions. In **a**: $\mathfrak{E}_n(s_1)$ varies depending on whether n is even; thus, $\lim_{n \rightarrow \infty} \mathfrak{E}_n(s_1)$ does not exist. In **b** and **c**: $\forall n : \mathfrak{E}_n(s_3) = \mathfrak{E}_n(s_4)$, even though s_4 allows greater control over future state trajectories than s_3 does. For example, suppose that in both **b** and **c**, the leftmost black state and the rightmost red state have 1 reward while all other states have 0 reward. In **c**, the agent can independently maximize the intermediate black-state reward and the delayed red-state reward. Independent maximization is not possible in **b**.

POWER returns intuitive answers in these situations. $\lim_{\gamma \rightarrow 1} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s_1, \gamma)$ converges by lemma 5.3. Consider the obvious involution ϕ which takes each state in fig. 9b to its counterpart in fig. 9c. Since $\phi \cdot \mathcal{F}_{\text{nd}}(s_3) \subsetneq \mathcal{F}_{\text{nd}}(s_4) = \mathcal{F}(s_4)$, proposition 6.6 proves that $\forall \gamma \in [0, 1] : \text{POWER}_{\mathcal{D}_{\text{bound}}}(s_3, \gamma) \leq_{\text{most: } \mathcal{D}_{\text{bound}}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s_4, \gamma)$, with the proof of proposition 6.6 showing strict inequality under all $\mathcal{D}_{X\text{-IID}}$ when $\gamma \in (0, 1)$.

Empowerment can be adjusted to account for these cases, perhaps by considering the channel capacity between the agent’s actions and the state trajectories induced by stationary policies. However, since POWER is formulated in terms of optimal value, we believe that POWER is better suited for MDPs than information-theoretic empowerment is.

Appendix B Seeking POWER can be a detour

Remark. The results of appendix E do not depend on this section’s results.

One might suspect that optimal policies tautologically tend to seek POWER. This intuition is wrong.

Proposition B.1 (Greater $\text{POWER}_{\mathcal{D}_{\text{bound}}}$ does not imply greater $\mathbb{P}_{\mathcal{D}_{\text{bound}}}$). *Action a seeking more $\text{POWER}_{\mathcal{D}_{\text{bound}}}$ than a' at state s and γ does not imply that $\mathbb{P}_{\mathcal{D}_{\text{bound}}}(s, a, \gamma) \geq \mathbb{P}_{\mathcal{D}_{\text{bound}}}(s, a', \gamma)$.*

Proof. Consider the environment of fig. 10. Let $X_u := \text{unif}(0, 1)$, and consider $\mathcal{D}_{X_u\text{-IID}}$, which has bounded support. Direct computation⁵ of the POWER expectation (definition 5.2) yields $\text{POWER}_{\mathcal{D}_{X_u\text{-IID}}}(s_2, 1) = \frac{3}{4} > \frac{2}{3} = \text{POWER}_{\mathcal{D}_{X_u\text{-IID}}}(s_3, 1)$. Therefore, N seeks more $\text{POWER}_{\mathcal{D}_{X_u\text{-IID}}}$ than NE at state s_1 and $\gamma = 1$.

However, $\mathbb{P}_{\mathcal{D}_{X_u\text{-IID}}}(s_1, \text{N}, 1) = \frac{1}{3} < \frac{2}{3} = \mathbb{P}_{\mathcal{D}_{X_u\text{-IID}}}(s_1, \text{NE}, 1)$. \square




Figure 10

⁵In small deterministic MDPs, the POWER and optimality probability of the maximum-entropy reward function distribution can be computed using <https://github.com/loganriggs/Optimal-Policies-Tend-To-Seek-Power>.

Lemma B.2 (Fraction of orbits which agree on weak optimality). *Let $\mathfrak{D} \subseteq \Delta(\mathbb{R}^{|\mathcal{S}|})$, and suppose $f_1, f_2 : \Delta(\mathbb{R}^{|\mathcal{S}|}) \rightarrow \mathbb{R}$ are such that $f_1(\mathcal{D}) \geq_{\text{most: } \mathfrak{D}} f_2(\mathcal{D})$. Then for all $\mathcal{D} \in \mathfrak{D}$,*

$$\frac{|\{\mathcal{D}' \in S_{|\mathcal{S}|} \cdot \mathcal{D} \mid f_1(\mathcal{D}') \geq f_2(\mathcal{D}')\}|}{|S_{|\mathcal{S}|} \cdot \mathcal{D}|} \geq \frac{1}{2}.$$

Proof. All $\mathcal{D}' \in S_{|\mathcal{S}|} \cdot \mathcal{D}$ such that $f_1(\mathcal{D}') = f_2(\mathcal{D}')$ satisfy $f_1(\mathcal{D}') \geq f_2(\mathcal{D}')$.

Otherwise, consider the $\mathcal{D}' \in S_{|\mathcal{S}|} \cdot \mathcal{D}$ such that $f_1(\mathcal{D}') \neq f_2(\mathcal{D}')$. By the definition of \geq_{most} (definition 6.5), at least $\frac{1}{2}$ of these \mathcal{D}' satisfy $f_1(\mathcal{D}') > f_2(\mathcal{D}')$, in which case $f_1(\mathcal{D}') \geq f_2(\mathcal{D}')$. Then the desired inequality follows. \square

Lemma B.3 (\geq_{most} and trivial orbits). *Let $\mathfrak{D} \subseteq \Delta(\mathbb{R}^{|\mathcal{S}|})$ and suppose $f_1(\mathcal{D}) \geq_{\text{most: } \mathfrak{D}} f_2(\mathcal{D})$. For all reward function distributions $\mathcal{D} \in \mathfrak{D}$ with one-element orbits, $f_1(\mathcal{D}) \geq f_2(\mathcal{D})$. In particular, \mathcal{D} has a one-element orbit when it distributes reward identically and independently (IID) across states.*

Proof. By lemma B.2, at least half of the elements $\mathcal{D}' \in S_{|\mathcal{S}|} \cdot \mathcal{D}$ satisfy $f_1(\mathcal{D}') \geq f_2(\mathcal{D}')$. But $|S_{|\mathcal{S}|} \cdot \mathcal{D}| = 1$, and so $f_1(\mathcal{D}) \geq f_2(\mathcal{D})$ must hold.

If \mathcal{D} is IID, it has a one-element orbit due to the assumed identical distribution of reward. \square

Proposition B.4 (Actions which tend to seek POWER do not necessarily tend to be optimal). *Action a tending to seek more POWER than a' at state s and γ does not imply that $\mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a, \gamma) \geq_{\text{most: } \mathfrak{D}_{\text{any}}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a', \gamma)$.*

Proof. Consider the environment of fig. 10. Since $\text{RSD}_{\text{nd}}(s_3) \subsetneq \text{RSD}(s_2)$, proposition 6.12 shows that $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_2, 1) \geq_{\text{most: } \mathfrak{D}_{\text{bound}}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s_3, 1)$ via $s' := s_3, s := s_2, \phi$ the identity permutation (which is an involution). Therefore, N tends to seek more POWER than NE at state s_1 and $\gamma = 1$.

If $\mathbb{P}_{\mathcal{D}_{\text{any}}}(s_1, \text{N}, 1) \geq_{\text{most: } \mathfrak{D}_{\text{any}}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(s_1, \text{NE}, 1)$, then lemma B.3 shows that $\mathbb{P}_{\mathcal{D}_{X\text{-IID}}}(s_1, \text{N}, 1) \geq \mathbb{P}_{\mathcal{D}_{X\text{-IID}}}(s_1, \text{NE}, 1)$ for all $\mathcal{D}_{X\text{-IID}}$. But the proof of proposition B.1 showed that $\mathbb{P}_{\mathcal{D}_{X_u\text{-IID}}}(s_1, \text{N}, 1) < \mathbb{P}_{\mathcal{D}_{X_u\text{-IID}}}(s_1, \text{NE}, 1)$ for $X_u := \text{unif}(0, 1)$. Therefore, it cannot be true that $\mathbb{P}_{\mathcal{D}_{\text{any}}}(s_1, \text{N}, 1) \geq_{\text{most: } \mathfrak{D}_{\text{any}}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(s_1, \text{NE}, 1)$. \square

Appendix C Sub-optimal POWER

In certain situations, POWER returns intuitively surprising verdicts. There exists a policy under which the reader chooses a winning lottery ticket, but it seems wrong to say that the reader has the power to win the lottery with high probability. For various reasons, humans and other bounded agents are generally incapable of computing optimal policies for arbitrary objectives. More formally, consider the rewardless MDP of fig. 11.




Figure 11: s_0 is the starting state, and $|\mathcal{A}| = 10^{10^{10}}$. At s_0 , half of the actions lead to s_ℓ , while the other half lead to s_r . Similarly, half of the actions at s_ℓ lead to s_1 , while the other half lead to s_2 . At s_r , one action leads to s_3 , one action leads to s_4 , and the remaining $10^{10^{10}} - 2$ actions lead to s_5 .

Consider a model-based RL agent with black-box simulator access to this environment. The agent has no prior information about the model, and so it acts randomly. Before long, the agent has probably

learned how to navigate from s_0 to states s_ℓ , s_r , s_1 , s_2 , and s_5 . However, over any reasonable timescale, it is extremely improbable that the agent discovers the two actions respectively leading to s_3 and s_4 .

Even provided with a reward function R and the discount rate γ , the agent has yet to learn the relevant environmental dynamics, and so many of its policies are far from optimal. Although proposition 6.6 shows that $\forall \gamma \in [0, 1] : \text{POWER}_{\mathcal{D}_{\text{bound}}}(s_\ell, \gamma) \leq_{\text{most: } \mathcal{D}_{\text{bound}}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s_r, \gamma)$, there is a sense in which s_ℓ gives this agent more power.

We formalize a bounded agent’s goal-achievement capabilities with a function pol , which takes as input a reward function and a discount rate, and returns a policy. Informally, this is the best policy which the agent knows about. We can then calculate $\text{POWER}_{\mathcal{D}_{\text{bound}}}$ with respect to pol .

Definition C.1 (Suboptimal POWER). Let Π_Δ be the set of stationary stochastic policies, and let $\text{pol} : \mathbb{R}^S \times [0, 1] \rightarrow \Pi_\Delta$. For $\gamma \in [0, 1]$,

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}^{\text{pol}}(s, \gamma) := \mathbb{E}_{\substack{R \sim \mathcal{D}_{\text{bound}}, \\ a \sim \text{pol}(R, \gamma)(s), \\ s' \sim T(s, a)}} \left[\lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) V_R^{\text{pol}(R, \gamma^*)}(s', \gamma^*) \right]. \quad (5)$$

By lemma E.38, $\text{POWER}_{\mathcal{D}_{\text{bound}}}$ is the special case where $\forall R \in \mathbb{R}^S, \gamma \in [0, 1] : \text{pol}(R, \gamma) \in \Pi^*(R, \gamma)$. We define $\text{POWER}_{\mathcal{D}_{\text{bound}}}^{\text{pol}}$ -seeking similarly as in definition 5.6.

$\text{POWER}_{\mathcal{D}_{\text{bound}}}^{\text{pol}}(s_0, 1)$ increases as the policies returned by pol are improved. We illustrate this by considering the $\mathcal{D}_{X\text{-IID}}$ case.

- pol_1 The model is initially unknown, and so $\forall R, \gamma : \text{pol}_1(R, \gamma)$ is a uniformly random policy. Since pol_1 is constant on its inputs, $\text{POWER}_{\mathcal{D}_{X\text{-IID}}}^{\text{pol}_1}(s_0, 1) = \mathbb{E}[X]$ by the linearity of expectation and the fact that $\mathcal{D}_{X\text{-IID}}$ distributes reward independently and identically across states.
- pol_2 The agent knows the dynamics, except that it does not know how to reach s_3 or s_4 . At this point, $\text{pol}_2(R, 1)$ navigates from s_0 to the average-optimal choice among three terminal states: s_1 , s_2 , and s_5 . Therefore, $\text{POWER}_{\mathcal{D}_{\text{bound}}}^{\text{pol}_2}(s_0, 1) = \mathbb{E}[\text{max of 3 draws from } X]$.
- pol_3 The agent knows the dynamics, the environment is small enough to solve explicitly, and so $\forall R, \gamma : \text{pol}_3(R, \gamma)$ is an optimal policy. $\text{pol}_3(R, 1)$ navigates from s_0 to the average-optimal choice among all five terminal states. Therefore, $\text{POWER}_{\mathcal{D}_{\text{bound}}}^{\text{pol}_3}(s_0, 1) = \mathbb{E}[\text{max of 5 draws from } X]$.

As the agent learns more about the environment and improves pol , the agent’s $\text{POWER}_{\mathcal{D}_{\text{bound}}}^{\text{pol}}$ increases. The agent seeks $\text{POWER}_{\mathcal{D}_{\text{bound}}}^{\text{pol}_2}$ by navigating to s_ℓ instead of s_r , but seeks more $\text{POWER}_{\mathcal{D}_{\text{bound}}}$ by navigating to s_r instead of s_ℓ . Intuitively, bounded agents gain power by improving pol and by formally seeking $\text{POWER}_{\mathcal{D}_{\text{bound}}}^{\text{pol}}$ within the environment.

Appendix D Lists of results

List of definitions

3.1	Definition (Rewardless MDP)	2
3.2	Definition (1-cycle states)	2
3.3	Definition (State visit distribution [Sutton and Barto, 1998])	3
3.4	Definition (\mathcal{F} single-state restriction)	3
3.5	Definition (Value function)	3
3.6	Definition (Non-domination)	3
4.1	Definition (Optimal policy set function)	3

4.2	Definition (Reward function distributions)	3
4.3	Definition (Visit distribution optimality probability)	4
4.4	Definition (Action optimality probability)	4
5.1	Definition (Average optimal value)	4
5.2	Definition (POWER)	4
5.6	Definition (POWER-seeking actions)	5
6.1	Definition (Similarity of vector sets)	6
6.2	Definition (Similarity of vector function sets)	6
6.3	Definition (Pushforward distribution of a permutation)	6
6.4	Definition (Orbit of a probability distribution)	6
6.5	Definition (Inequalities which hold for most probability distributions)	6
6.7	Definition (Equivalent actions)	7
6.8	Definition (States reachable after taking an action)	7
6.10	Definition (Recurrent state distributions [Puterman, 2014])	8
6.11	Definition (Average-optimal policies)	8
C.1	Definition (Suboptimal POWER)	15
E.2	Definition (Transition matrix induced by a policy)	18
E.6	Definition (Continuous reward function distribution)	19
E.9	Definition (Non-dominated linear functionals)	20
E.13	Definition (Non-dominated vector functions)	21
E.14	Definition (Affine transformation of visit distribution sets)	21
E.18	Definition (Support of \mathcal{D}_{any})	22
E.19	Definition (Linear functional optimality probability)	22
E.23	Definition (Bounded, continuous IID reward)	23
E.25	Definition (Indicator function)	24
E.31	Definition (Evaluating sets of visit distribution functions at γ)	27
E.39	Definition (Discount-normalized value function)	29
E.42	Definition (Normalized visit distribution function)	32

List of theorems

5.3	Lemma (Continuity of POWER)	5
5.4	Proposition (Maximal POWER)	5
5.5	Proposition (POWER is smooth across reversible dynamics)	5
6.6	Proposition (States with “more options” have more POWER)	6
6.9	Proposition (Keeping options open tends to be POWER-seeking and tends to be optimal)	7
6.12	Proposition (When $\gamma = 1$, RSDs control POWER)	8
6.13	Theorem (Average-optimal policies tend to end up in “larger” sets of RSDs)	8
6.14	Corollary (Average-optimal policies tend not to end up in any given 1-cycle)	8
B.1	Proposition (Greater $\text{POWER}_{\mathcal{D}_{\text{bound}}}$ does not imply greater $\mathbb{P}_{\mathcal{D}_{\text{bound}}}$)	13

B.2	Lemma (Fraction of orbits which agree on weak optimality)	14
B.3	Lemma (\geq_{most} and trivial orbits)	14
B.4	Proposition (Actions which tend to seek POWER do not necessarily tend to be optimal)	14
E.1	Lemma (A policy is optimal iff it induces an optimal visit distribution at every state)	18
E.3	Proposition (Properties of visit distribution functions)	18
E.4	Lemma ($\mathbf{f} \in \mathcal{F}(s)$ is multivariate rational on γ)	19
E.5	Corollary (On-policy value is rational on γ)	19
E.7	Lemma (Distinct linear functionals disagree almost everywhere on their domains)	19
E.8	Corollary (Unique maximization of almost all vectors)	20
E.10	Lemma (All vectors are maximized by a non-dominated linear functional)	20
E.11	Corollary (Maximal value is invariant to restriction to non-dominated functionals)	20
E.12	Lemma (How non-domination containment affects optimal value)	20
E.15	Lemma (Invariance of non-domination under positive affine transform)	21
E.16	Lemma (Helper lemma for demonstrating $\geq_{\text{most}: \mathfrak{D}_{\text{any}}}$)	21
E.17	Lemma (A helper result for expectations of functions)	22
E.20	Proposition (Non-dominated linear functionals and their optimality probability)	22
E.21	Lemma (Expected value of similar linear functional sets)	23
E.22	Lemma (For continuous IID distributions $\mathcal{D}_{X\text{-IID}}$, $\exists b < c : (b, c)^{ S } \subseteq \text{supp}(\mathcal{D}_{X\text{-IID}})$)	23
E.24	Lemma (Expectation superiority lemma)	23
E.26	Lemma (Optimality probability inclusion relations)	24
E.27	Lemma (Optimality probability of similar linear functional sets)	25
E.28	Lemma (Optimality probability superiority lemma)	26
E.29	Lemma (Limit probability inequalities which hold for most distributions)	26
E.30	Proposition (How to transfer optimal policy sets across discount rates)	27
E.32	Lemma (Non-domination across γ values for expectations of visit distributions)	27
E.33	Lemma ($\forall \gamma \in (0, 1) : \mathbf{d} \in \mathcal{F}_{\text{nd}}(s, \gamma)$ iff $\mathbf{d} \in \text{ND}(\mathcal{F}(s, \gamma))$)	28
E.34	Lemma ($\forall \gamma \in [0, 1) : V_R^*(s, \gamma) = \max_{\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)} \mathbf{f}(\gamma)^\top \mathbf{r}$)	28
E.35	Lemma (Optimal policy shift bound)	28
E.36	Proposition (Optimality probability's limits exist)	28
E.37	Lemma (Optimality probability identity)	28
E.38	Lemma (POWER identities)	29
E.40	Lemma (Normalized value functions have uniformly bounded derivative)	30
E.41	Lemma (Lower bound on current POWER based on future POWER)	31
E.43	Lemma (Normalized visit distribution functions are continuous)	32
E.44	Lemma (Non-domination of normalized visit distribution functions)	33
E.45	Lemma (POWER limit identity)	33
E.46	Lemma (Lemma for POWER superiority)	34
E.47	Lemma (Non-dominated visit distribution functions never agree with other visit distribution functions at that state)	36

E.48 Corollary (Cardinality of non-dominated visit distributions)	36
E.49 Lemma (Optimality probability and state bottlenecks)	36
E.50 Lemma (Action optimality probability is a special case of visit distribution optimality probability)	37
E.51 Lemma (POWER identity when $\gamma = 1$)	40
E.52 Proposition (RSD properties)	42
E.53 Lemma (When reachable with probability 1, 1-cycles induce non-dominated RSDs)	42

D.1 Contributions of independent interest

We developed new basic MDP theory by exploring the structural properties of visit distribution functions. Echoing Wang et al. [2007, 2008], we believe that this area is interesting and underexplored.

D.1.1 Optimal value theory

Lemma E.40 shows that $f(\gamma^*) := \lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) V_R^*(s, \gamma^*)$ is Lipschitz continuous on $\gamma \in [0, 1]$, with Lipschitz constant depending only on $\|R\|_1$. For all states s and policies $\pi \in \Pi$, corollary E.5 shows that $V_R^\pi(s, \gamma)$ is rational on γ .

Optimal value has a well-known dual formulation: $V_R^*(s, \gamma) = \max_{\mathbf{f} \in \mathcal{F}(s)} \mathbf{f}(\gamma)^\top \mathbf{r}$.

Lemma E.34 ($\forall \gamma \in [0, 1] : V_R^*(s, \gamma) = \max_{\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)} \mathbf{f}(\gamma)^\top \mathbf{r}$).

In a fixed rewardless MDP, lemma E.34 may enable more efficient computation of optimal value functions for multiple reward functions.

D.1.2 Optimal policy theory

Proposition E.30 demonstrates how to preserve optimal incentives while changing the discount rate.

Proposition E.30 (How to transfer optimal policy sets across discount rates). *Suppose reward function R has optimal policy set $\Pi^*(R, \gamma)$ at discount rate $\gamma \in (0, 1)$. For any $\gamma^* \in (0, 1)$, we can construct a reward function R' such that $\Pi^*(R', \gamma^*) = \Pi^*(R, \gamma)$. Furthermore, $V_{R'}^*(\cdot, \gamma^*) = V_R^*(\cdot, \gamma)$.*

D.1.3 Visit distribution theory

While Regan and Boutilier [2010] consider a visit distribution function $\mathbf{f} \in \mathcal{F}(s)$ to be non-dominated if it is optimal for some reward function in a set $\mathcal{R} \subseteq \mathbb{R}^{|S|}$, our stricter definition 3.6 considers \mathbf{f} to be non-dominated when $\exists \mathbf{r} \in \mathbb{R}^{|S|}, \gamma \in (0, 1) : \mathbf{f}(\gamma)^\top \mathbf{r} > \max_{\mathbf{f}' \in \mathcal{F}(s) \setminus \{\mathbf{f}\}} \mathbf{f}'(\gamma)^\top \mathbf{r}$.

Appendix E Theoretical results

Lemma E.1 (A policy is optimal iff it induces an optimal visit distribution at every state). *Let $\gamma \in (0, 1)$ and let R be a reward function. $\pi \in \Pi^*(R, \gamma)$ iff π induces an optimal visit distribution at every state.*

Proof. By definition, a policy π is optimal iff π induces the maximal on-policy value at each state, which is true iff π induces an optimal visit distribution at every state (by the dual formulation of optimal value functions). \square

Definition E.2 (Transition matrix induced by a policy). \mathbf{T}^π is the transition matrix induced by policy $\pi \in \Pi$, where $\mathbf{T}^\pi \mathbf{e}_s := T(s, \pi(s))$. $(\mathbf{T}^\pi)^t \mathbf{e}_s$ gives the probability distribution over the states visited at time step t , after following π for t steps from s .

Proposition E.3 (Properties of visit distribution functions). *Let $s, s' \in \mathcal{S}, \mathbf{f}^{\pi, s} \in \mathcal{F}(s)$.*

1. $\mathbf{f}^{\pi, s}(\gamma)$ is element-wise non-negative and element-wise monotonically increasing on $\gamma \in [0, 1]$.

$$2. \forall \gamma \in [0, 1) : \|\mathbf{f}^{\pi, s}(\gamma)\|_1 = \frac{1}{1-\gamma}.$$

Proof. Item 1: by examination of definition 3.3, $\mathbf{f}^{\pi, s} = \sum_{t=0}^{\infty} (\gamma \mathbf{T}^{\pi})^t \mathbf{e}_s$. Since each $(\mathbf{T}^{\pi})^t$ is left stochastic and \mathbf{e}_s is the standard unit vector, each entry in each summand is non-negative. Therefore, $\forall \gamma \in [0, 1) : \mathbf{f}^{\pi, s}(\gamma)^{\top} \mathbf{e}_{s'} \geq 0$, and this function monotonically increases on γ .

Item 2:

$$\|\mathbf{f}^{\pi, s}(\gamma)\|_1 = \left\| \sum_{t=0}^{\infty} (\gamma \mathbf{T}^{\pi})^t \mathbf{e}_s \right\|_1 \quad (6)$$

$$= \sum_{t=0}^{\infty} \gamma^t \left\| (\mathbf{T}^{\pi})^t \mathbf{e}_s \right\|_1 \quad (7)$$

$$= \sum_{t=0}^{\infty} \gamma^t \quad (8)$$

$$= \frac{1}{1-\gamma}. \quad (9)$$

Equation (7) follows because all entries in each $(\mathbf{T}^{\pi})^t \mathbf{e}_s$ are non-negative by item 1. Equation (8) follows because each $(\mathbf{T}^{\pi})^t$ is left stochastic and \mathbf{e}_s is a stochastic vector, and so $\left\| (\mathbf{T}^{\pi})^t \mathbf{e}_s \right\|_1 = 1$. \square

Lemma E.4 ($\mathbf{f} \in \mathcal{F}(s)$ is multivariate rational on γ). $\mathbf{f}^{\pi} \in \mathcal{F}(s)$ is a multivariate rational function on $\gamma \in [0, 1)$.

Proof. Let $\mathbf{r} \in \mathbb{R}^{|\mathcal{S}|}$ and consider $\mathbf{f}^{\pi} \in \mathcal{F}(s)$. Let \mathbf{v}_R^{π} be the $V_R^*(s, \gamma)$ function in column vector form, with one entry per state value.

By the Bellman equations, $\mathbf{v}_R^{\pi} = (\mathbf{I} - \gamma \mathbf{T}^{\pi})^{-1} \mathbf{r}$. Let $\mathbf{A}_{\gamma} := (\mathbf{I} - \gamma \mathbf{T}^{\pi})^{-1}$, and for state s , form $\mathbf{A}_{s, \gamma}$ by replacing \mathbf{A}_{γ} 's column for state s with \mathbf{r} . As noted by Lippman [1968], by Cramer's rule, $V_R^{\pi}(s, \gamma) = \frac{\det \mathbf{A}_{s, \gamma}}{\det \mathbf{A}_{\gamma}}$ is a rational function with numerator and denominator having degree at most $|\mathcal{S}|$.

In particular, for each state indicator reward function \mathbf{e}_{s_i} , $V_{s_i}^{\pi}(s, \gamma) = \mathbf{f}^{\pi, s}(\gamma)^{\top} \mathbf{e}_{s_i}$ is a rational function of γ whose numerator and denominator each have degree at most $|\mathcal{S}|$. This implies that $\mathbf{f}^{\pi}(\gamma)$ is multivariate rational on $\gamma \in [0, 1)$. \square

Corollary E.5 (On-policy value is rational on γ). Let $\pi \in \Pi$ and R be any reward function. $V_R^{\pi}(s, \gamma)$ is rational on $\gamma \in [0, 1)$.

Proof. $V_R^{\pi}(s, \gamma) = \mathbf{f}^{\pi, s}(\gamma)^{\top} \mathbf{r}$, and \mathbf{f} is a multivariate rational function of γ by lemma E.4. Therefore, for fixed \mathbf{r} , $\mathbf{f}^{\pi, s}(\gamma)^{\top} \mathbf{r}$ is a rational function of γ . \square

E.1 Non-dominated visit distribution functions

Definition E.6 (Continuous reward function distribution). Results with $\mathcal{D}_{\text{cont}}$ hold for any absolutely continuous reward function distribution.

Remark. We assume $\mathbb{R}^{|\mathcal{S}|}$ is endowed with the standard topology.

Lemma E.7 (Distinct linear functionals disagree almost everywhere on their domains). Let $\mathbf{x}, \mathbf{x}' \in \mathbb{R}^{|\mathcal{S}|}$ be distinct. $\mathbb{P}_{\mathbf{r} \sim \mathcal{D}_{\text{cont}}}(\mathbf{x}^{\top} \mathbf{r} = \mathbf{x}'^{\top} \mathbf{r}) = 0$.

Proof. $\{\mathbf{r} \in \mathbb{R}^{|\mathcal{S}|} \mid (\mathbf{x} - \mathbf{x}')^{\top} \mathbf{r} = 0\}$ is a hyperplane since $\mathbf{x} - \mathbf{x}' \neq \mathbf{0}$. Therefore, it has no interior in the standard topology on $\mathbb{R}^{|\mathcal{S}|}$. Since this empty-interior set is also convex, it has zero Lebesgue measure. By the Radon-Nikodym theorem, it has zero measure under any continuous distribution $\mathcal{D}_{\text{cont}}$. \square

Corollary E.8 (Unique maximization of almost all vectors). *Let $X \subsetneq \mathbb{R}^{|S|}$ be finite. $\mathbb{P}_{\mathbf{r} \sim \mathcal{D}_{\text{cont}}} \left(\left| \arg \max_{\mathbf{x}'' \in X} \mathbf{x}''^\top \mathbf{r} \right| > 1 \right) = 0$.*

Proof. Let $\mathbf{x}, \mathbf{x}' \in X$ be distinct. For any $\mathbf{r} \in \mathbb{R}^{|S|}$, $\mathbf{x}, \mathbf{x}' \in \arg \max_{\mathbf{x}'' \in X} \mathbf{x}''^\top \mathbf{r}$ iff $\mathbf{x}^\top \mathbf{r} = \mathbf{x}'^\top \mathbf{r} \geq \max_{\mathbf{x}'' \in X \setminus \{\mathbf{x}, \mathbf{x}'\}} \mathbf{x}''^\top \mathbf{r}$. By lemma E.7, $\mathbf{x}^\top \mathbf{r} = \mathbf{x}'^\top \mathbf{r}$ holds with probability 0 under any $\mathcal{D}_{\text{cont}}$. \square

E.1.1 Generalized non-domination results

Our formalism includes both $\mathcal{F}_{\text{nd}}(s)$ and $\text{RSD}_{\text{nd}}(s)$; we therefore prove results that are applicable to both.

Definition E.9 (Non-dominated linear functionals). Let $X \subsetneq \mathbb{R}^{|S|}$ be finite. $\text{ND}(X) := \left\{ \mathbf{x} \in X \mid \exists \mathbf{r} \in \mathbb{R}^{|S|} : \mathbf{x}^\top \mathbf{r} > \max_{\mathbf{x}' \in X \setminus \{\mathbf{x}\}} \mathbf{x}'^\top \mathbf{r} \right\}$.

Lemma E.10 (All vectors are maximized by a non-dominated linear functional). *Let $\mathbf{r} \in \mathbb{R}^{|S|}$ and let $X \subsetneq \mathbb{R}^{|S|}$ be finite and non-empty. $\exists \mathbf{x}^* \in \text{ND}(X) : \mathbf{x}^{*\top} \mathbf{r} = \max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r}$.*

Proof. Let $A(\mathbf{r} \mid X) := \arg \max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$. Then

$$\mathbf{x}_1^\top \mathbf{r} = \dots = \mathbf{x}_n^\top \mathbf{r} > \max_{\mathbf{x}' \in X \setminus A(\mathbf{r} \mid X)} \mathbf{x}'^\top \mathbf{r}. \quad (10)$$

In eq. (10), each $\mathbf{x}^\top \mathbf{r}$ expression is linear on \mathbf{r} . The max is piecewise linear on \mathbf{r} since it is the maximum of a finite set of linear functionals. In particular, all expressions in eq. (10) are continuous on \mathbf{r} , and so we can find some $\delta > 0$ neighborhood $B(\mathbf{r}, \delta)$ such that $\forall \mathbf{r}' \in B(\mathbf{r}, \delta) : \max_{\mathbf{x}_i \in A(\mathbf{r} \mid X)} \mathbf{x}_i^\top \mathbf{r}' > \max_{\mathbf{x}' \in X \setminus A(\mathbf{r} \mid X)} \mathbf{x}'^\top \mathbf{r}'$.

But almost all $\mathbf{r}' \in B(\mathbf{r}, \delta)$ are maximized by a unique functional \mathbf{x}^* by corollary E.8; in particular, at least one such \mathbf{r}'' exists. Formally, $\exists \mathbf{r}'' \in B(\mathbf{r}, \delta) : \mathbf{x}^{*\top} \mathbf{r}'' > \max_{\mathbf{x}' \in X \setminus \{\mathbf{x}^*\}} \mathbf{x}'^\top \mathbf{r}''$. Therefore, $\mathbf{x}^* \in \text{ND}(X)$ by definition E.9.

$\mathbf{x}^{*\top} \mathbf{r}' \geq \max_{\mathbf{x}_i \in A(\mathbf{r} \mid X)} \mathbf{x}_i^\top \mathbf{r}' > \max_{\mathbf{x}' \in X \setminus A(\mathbf{r} \mid X)} \mathbf{x}'^\top \mathbf{r}'$, with the strict inequality following because $\mathbf{r}'' \in B(\mathbf{r}, \delta)$. These inequalities imply that $\mathbf{x}^* \in A(\mathbf{r} \mid X)$. \square

Corollary E.11 (Maximal value is invariant to restriction to non-dominated functionals). *Let $\mathbf{r} \in \mathbb{R}^{|S|}$ and let $X \subsetneq \mathbb{R}^{|S|}$ be finite. $\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} = \max_{\mathbf{x} \in \text{ND}(X)} \mathbf{x}^\top \mathbf{r}$.*

Proof. If X is empty, holds trivially. Otherwise, apply lemma E.10. \square

Lemma E.12 (How non-domination containment affects optimal value). *Let $\mathbf{r} \in \mathbb{R}^{|S|}$ and let $X, X' \subsetneq \mathbb{R}^{|S|}$ be finite.*

1. *If $\text{ND}(X) \subseteq X'$, then $\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} \leq \max_{\mathbf{x}' \in X'} \mathbf{x}'^\top \mathbf{r}$.*
2. *If $\text{ND}(X) \subseteq X' \subseteq X$, then $\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} = \max_{\mathbf{x}' \in X'} \mathbf{x}'^\top \mathbf{r}$.*

Proof. Item 1:

$$\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} = \max_{\mathbf{x} \in \text{ND}(X)} \mathbf{x}^\top \mathbf{r} \quad (11)$$

$$\leq \max_{\mathbf{x}' \in X'} \mathbf{x}'^\top \mathbf{r}. \quad (12)$$

Equation (11) follows by corollary E.11. Equation (12) follows because $\text{ND}(X) \subseteq X'$.

Item 2: by item 1, $\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} \leq \max_{\mathbf{x}' \in X'} \mathbf{x}'^\top \mathbf{r}$. Since $X' \subseteq X$, we also have $\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} \geq \max_{\mathbf{x}' \in X'} \mathbf{x}'^\top \mathbf{r}$, and so equality must hold. \square

Definition E.13 (Non-dominated vector functions). Let $I \subseteq \mathbb{R}$ and let $F \subsetneq \left(\mathbb{R}^{|S|}\right)^I$ be a finite set of vector-valued functions on I . $\text{ND}(F) := \left\{ \mathbf{f} \in F \mid \exists \gamma \in I, \mathbf{r} \in \mathbb{R}^{|S|} : \mathbf{f}(\gamma)^\top \mathbf{r} > \max_{\mathbf{f}' \in F \setminus \{\mathbf{f}\}} \mathbf{f}'(\gamma)^\top \mathbf{r} \right\}$.

Remark. $\mathcal{F}_{\text{nd}}(s) = \text{ND}(\mathcal{F}(s))$ by definition 3.6.

Definition E.14 (Affine transformation of visit distribution sets). For notational convenience, we define set-scalar multiplication and set-vector addition on $X \subseteq \mathbb{R}^{|S|}$: for $c \in \mathbb{R}$, $cX := \{c\mathbf{x} \mid \mathbf{x} \in X\}$. For $\mathbf{a} \in \mathbb{R}^{|S|}$, $X + \mathbf{a} := \{\mathbf{x} + \mathbf{a} \mid \mathbf{x} \in X\}$. Similar operations hold when X is a set of vector functions $\mathbb{R} \mapsto \mathbb{R}^{|S|}$.

Lemma E.15 (Invariance of non-domination under positive affine transform).

1. Let $X \subsetneq \mathbb{R}^{|S|}$ be finite. If $\mathbf{x} \in \text{ND}(X)$, then $\forall c > 0, \mathbf{a} \in \mathbb{R}^{|S|} : (c\mathbf{x} + \mathbf{a}) \in \text{ND}(cX + \mathbf{a})$.
2. Let $I \subseteq \mathbb{R}$ and let $F \subsetneq \left(\mathbb{R}^{|S|}\right)^I$ be a finite set of vector-valued functions on I . If $\mathbf{f} \in \text{ND}(F)$, then $\forall c > 0, \mathbf{a} \in \mathbb{R}^{|S|} : (c\mathbf{f} + \mathbf{a}) \in \text{ND}(cF + \mathbf{a})$.

Proof. Item 1: Suppose $\mathbf{x} \in \text{ND}(X)$ is strictly optimal for $\mathbf{r} \in \mathbb{R}^{|S|}$. Then let $c > 0, \mathbf{a} \in \mathbb{R}^{|S|}$ be arbitrary, and define $b := \mathbf{a}^\top \mathbf{r}$.

$$\mathbf{x}^\top \mathbf{r} > \max_{\mathbf{x}' \in X \setminus \{\mathbf{x}\}} \mathbf{x}'^\top \mathbf{r} \quad (13)$$

$$c\mathbf{x}^\top \mathbf{r} + b > \max_{\mathbf{x}' \in X \setminus \{\mathbf{x}\}} c\mathbf{x}'^\top \mathbf{r} + b \quad (14)$$

$$(c\mathbf{x} + \mathbf{a})^\top \mathbf{r} > \max_{\mathbf{x}' \in X \setminus \{\mathbf{x}\}} (c\mathbf{x}' + \mathbf{a})^\top \mathbf{r} \quad (15)$$

$$(c\mathbf{x} + \mathbf{a})^\top \mathbf{r} > \max_{\mathbf{x}'' \in (cX + \mathbf{a}) \setminus \{c\mathbf{x} + \mathbf{a}\}} \mathbf{x}''^\top \mathbf{r}. \quad (16)$$

Equation (14) follows because $c > 0$. Equation (15) follows by the definition of b .

Item 2: If $\mathbf{f} \in \text{ND}(F)$, then by definition E.13, there exist $\gamma \in I, \mathbf{r} \in \mathbb{R}^{|S|}$ such that

$$\mathbf{f}(\gamma)^\top \mathbf{r} > \max_{\mathbf{f}' \in F \setminus \{\mathbf{f}\}} \mathbf{f}'(\gamma)^\top \mathbf{r}. \quad (17)$$

Apply item 1 to conclude

$$(c\mathbf{f}(\gamma) + \mathbf{a})^\top \mathbf{r} > \max_{(c\mathbf{f}' + \mathbf{a}) \in (cF + \mathbf{a}) \setminus \{c\mathbf{f} + \mathbf{a}\}} (c\mathbf{f}'(\gamma) + \mathbf{a})^\top \mathbf{r}. \quad (18)$$

Therefore, $(c\mathbf{f} + \mathbf{a}) \in \text{ND}(cF + \mathbf{a})$. □

E.1.2 Inequalities which hold under most reward function distributions

Definition 6.5 (Inequalities which hold for most probability distributions). Let $f_1, f_2 : \Delta(\mathbb{R}^{|S|}) \rightarrow \mathbb{R}$ be functions from reward function distributions to real numbers and let $\mathfrak{D} \subseteq \Delta(\mathbb{R}^{|S|})$. We write $f_1(\mathcal{D}) \geq_{\text{most: } \mathfrak{D}} f_2(\mathcal{D})$ when, for all $\mathcal{D} \in \mathfrak{D}$, the following cardinality inequality holds:

$$\left| \{\mathcal{D}' \in S_{|S|} \cdot \mathcal{D} \mid f_1(\mathcal{D}') > f_2(\mathcal{D}')\} \right| \geq \left| \{\mathcal{D}' \in S_{|S|} \cdot \mathcal{D} \mid f_1(\mathcal{D}') < f_2(\mathcal{D}')\} \right|. \quad (4)$$

Lemma E.16 (Helper lemma for demonstrating $\geq_{\text{most: } \mathfrak{D}_{\text{any}}}$). Let $\mathfrak{D} \subseteq \Delta(\mathbb{R}^{|S|})$. If $\exists \phi \in S_{|S|}$ such that for all $\mathcal{D} \in \mathfrak{D}$, $f_1(\mathcal{D}) < f_2(\mathcal{D})$ implies that $f_1(\phi \cdot \mathcal{D}) > f_2(\phi \cdot \mathcal{D})$, then $f_1(\mathcal{D}) \geq_{\text{most: } \mathfrak{D}} f_2(\mathcal{D})$.

Proof. Since ϕ does not belong to the stabilizer of $S_{|S|}$, ϕ acts injectively on $S_{|S|} \cdot \mathcal{D}$. By assumption on ϕ , the image of $\{\mathcal{D}' \in S_{|S|} \cdot \mathcal{D} \mid f_1(\mathcal{D}') < f_2(\mathcal{D}')\}$ under ϕ is a subset of $\{\mathcal{D}' \in S_{|S|} \cdot \mathcal{D} \mid f_1(\mathcal{D}') > f_2(\mathcal{D}')\}$. Since ϕ is injective, $\left| \{\mathcal{D}' \in S_{|S|} \cdot \mathcal{D} \mid f_1(\mathcal{D}') < f_2(\mathcal{D}')\} \right| \leq \left| \{\mathcal{D}' \in S_{|S|} \cdot \mathcal{D} \mid f_1(\mathcal{D}') > f_2(\mathcal{D}')\} \right|$. $f_1(\mathcal{D}) \geq_{\text{most: } \mathfrak{D}} f_2(\mathcal{D})$ by definition 6.5. □

Lemma E.17 (A helper result for expectations of functions). *Let $B_1, \dots, B_n \subsetneq \mathbb{R}^{|\mathcal{S}|}$ be finite and let $\mathcal{D} \subseteq \Delta(\mathbb{R}^{|\mathcal{S}|})$. Suppose f is a function of the form*

$$f(B_1, \dots, B_n \mid \mathcal{D}) = \mathbb{E}_{\mathbf{r} \sim \mathcal{D}} \left[g \left(\max_{\mathbf{b}_1 \in B_1} \mathbf{b}_1^\top \mathbf{r}, \dots, \max_{\mathbf{b}_n \in B_n} \mathbf{b}_n^\top \mathbf{r} \right) \right] \quad (19)$$

for some function g , and that f is well-defined for all $\mathcal{D} \in \mathcal{D}$. Let ϕ be a state permutation. Then

$$f(B_1, \dots, B_n \mid \mathcal{D}) = f(\phi \cdot B_1, \dots, \phi \cdot B_n \mid \phi \cdot \mathcal{D}). \quad (20)$$

Proof. Let distribution \mathcal{D} have probability measure F , and let $\phi \cdot \mathcal{D}$ have probability measure F_ϕ .

$$f(B_1, \dots, B_n \mid \mathcal{D}) \quad (21)$$

$$:= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}} \left[g \left(\max_{\mathbf{b}_1 \in B_1} \mathbf{b}_1^\top \mathbf{r}, \dots, \max_{\mathbf{b}_n \in B_n} \mathbf{b}_n^\top \mathbf{r} \right) \right] \quad (22)$$

$$:= \int_{\mathbb{R}^{|\mathcal{S}|}} g \left(\max_{\mathbf{b}_1 \in B_1} \mathbf{b}_1^\top \mathbf{r}, \dots, \max_{\mathbf{b}_n \in B_n} \mathbf{b}_n^\top \mathbf{r} \right) dF(\mathbf{r}) \quad (23)$$

$$= \int_{\mathbb{R}^{|\mathcal{S}|}} g \left(\max_{\mathbf{b}_1 \in B_1} \mathbf{b}_1^\top \mathbf{r}, \dots, \max_{\mathbf{b}_n \in B_n} \mathbf{b}_n^\top \mathbf{r} \right) dF_\phi(\mathbf{P}_\phi \mathbf{r}) \quad (24)$$

$$= \int_{\mathbb{R}^{|\mathcal{S}|}} g \left(\max_{\mathbf{b}_1 \in B_1} \mathbf{b}_1^\top (\mathbf{P}_\phi^{-1} \mathbf{r}'), \dots, \max_{\mathbf{b}_n \in B_n} \mathbf{b}_n^\top (\mathbf{P}_\phi^{-1} \mathbf{r}') \right) |\det \mathbf{P}_\phi| dF_\phi(\mathbf{r}') \quad (25)$$

$$= \int_{\mathbb{R}^{|\mathcal{S}|}} g \left(\max_{\mathbf{b}_1 \in B_1} (\mathbf{P}_\phi \mathbf{b}_1)^\top \mathbf{r}', \dots, \max_{\mathbf{b}_n \in B_n} (\mathbf{P}_\phi \mathbf{b}_n)^\top \mathbf{r}' \right) dF_\phi(\mathbf{r}') \quad (26)$$

$$= \int_{\mathbb{R}^{|\mathcal{S}|}} g \left(\max_{\mathbf{b}'_1 \in \phi \cdot B_1} \mathbf{b}'_1{}^\top \mathbf{r}', \dots, \max_{\mathbf{b}'_n \in \phi \cdot B_n} \mathbf{b}'_n{}^\top \mathbf{r}' \right) dF_\phi(\mathbf{r}') \quad (27)$$

$$=: f(\phi \cdot B_1, \dots, \phi \cdot B_n \mid \phi \cdot \mathcal{D}). \quad (28)$$

Equation (24) follows by the definition of F_ϕ (definition 6.3). Equation (25) follows by substituting $\mathbf{r}' := \mathbf{P}_\phi \mathbf{r}$. Equation (26) follows from the fact that all permutation matrices have unitary determinant and are orthogonal (and so $(\mathbf{P}_\phi^{-1})^\top = \mathbf{P}_\phi$). \square

Definition E.18 (Support of \mathcal{D}_{any}). Let \mathcal{D}_{any} be any reward function distribution. $\text{supp}(\mathcal{D}_{\text{any}})$ is the smallest closed subset of $\mathbb{R}^{|\mathcal{S}|}$ whose complement has measure zero under \mathcal{D}_{any} .

Definition E.19 (Linear functional optimality probability). For finite $A, B \subsetneq \mathbb{R}^{|\mathcal{S}|}$, the probability under \mathcal{D}_{any} that A is optimal over B is $p_{\mathcal{D}_{\text{any}}}(A \geq B) := \mathbb{P}_{\mathbf{r} \sim \mathcal{D}_{\text{any}}}(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \geq \max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r})$.

Proposition E.20 (Non-dominated linear functionals and their optimality probability). *Let $A \subsetneq \mathbb{R}^{|\mathcal{S}|}$ be finite. If $\exists b < c : [b, c]^{|\mathcal{S}|} \subseteq \text{supp}(\mathcal{D}_{\text{any}})$, then $\mathbf{a} \in \text{ND}(A)$ implies that \mathbf{a} is strictly optimal for a set of reward functions with positive measure under \mathcal{D}_{any} .*

Proof. Suppose $\exists b < c : [b, c]^{|\mathcal{S}|} \subseteq \text{supp}(\mathcal{D}_{\text{any}})$. If $\mathbf{a} \in \text{ND}(A)$, then let \mathbf{r} be such that $\mathbf{a}^\top \mathbf{r} > \max_{\mathbf{a}' \in A \setminus \{\mathbf{a}\}} \mathbf{a}'^\top \mathbf{r}$. For $a_1 > 0, a_2 \in \mathbb{R}$, positively affinely transform $\mathbf{r}' := a_1 \mathbf{r} + a_2 \mathbf{1}$ (where $\mathbf{1} \in \mathbb{R}^{|\mathcal{S}|}$ is the all-ones vector) so that $\mathbf{r}' \in (b, c)^{|\mathcal{S}|}$.

Note that \mathbf{a} is still strictly optimal for \mathbf{r}' :

$$\mathbf{a}^\top \mathbf{r} > \max_{\mathbf{a}' \in A \setminus \{\mathbf{a}\}} \mathbf{a}'^\top \mathbf{r} \iff \mathbf{a}^\top \mathbf{r}' > \max_{\mathbf{a}' \in A \setminus \{\mathbf{a}\}} \mathbf{a}'^\top \mathbf{r}'. \quad (29)$$

Furthermore, by the continuity of both terms on the right-hand side of eq. (29), \mathbf{a} is strictly optimal for reward functions in some open neighborhood N of \mathbf{r}' . Let $N' := N \cap (b, c)^{|\mathcal{S}|}$. N' is still open in $\mathbb{R}^{|\mathcal{S}|}$ since it is the intersection of two open sets N and $(b, c)^{|\mathcal{S}|}$.

\mathcal{D}_{any} must assign positive probability measure to all open sets in its support; otherwise, its support would exclude these zero-measure sets by definition E.18. Therefore, \mathcal{D}_{any} assigns positive probability to $N' \subseteq \text{supp}(\mathcal{D}_{\text{any}})$. \square

Lemma E.21 (Expected value of similar linear functional sets). *Let $A, B \subseteq \mathbb{R}^{|S|}$ be finite, let A' be such that $\text{ND}(A) \subseteq A' \subseteq A$, and let $g : \mathbb{R} \rightarrow \mathbb{R}$ be an increasing function. If B contains a copy B' of A' via ϕ , then*

$$\mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right] \leq \mathbb{E}_{\mathbf{r} \sim \phi \cdot \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right]. \quad (30)$$

If $\text{ND}(B) \setminus B'$ is empty, then eq. (30) is an equality. If $\text{ND}(B) \setminus B'$ is non-empty, g is strictly increasing, and $\exists b < c : (b, c)^{|S|} \subseteq \text{supp}(\mathcal{D}_{\text{bound}})$, then eq. (30) is strict.

Proof. Because $g : \mathbb{R} \rightarrow \mathbb{R}$ is increasing, it is measurable (as is \max). Therefore, the relevant expectations exist for all $\mathcal{D}_{\text{bound}}$.

$$\mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right] = \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in A'} \mathbf{a}^\top \mathbf{r} \right) \right] \quad (31)$$

$$= \mathbb{E}_{\mathbf{r} \sim \phi \cdot \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in \phi \cdot A'} \mathbf{a}^\top \mathbf{r} \right) \right] \quad (32)$$

$$= \mathbb{E}_{\mathbf{r} \sim \phi \cdot \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B'} \mathbf{b}^\top \mathbf{r} \right) \right] \quad (33)$$

$$\leq \mathbb{E}_{\mathbf{r} \sim \phi \cdot \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right]. \quad (34)$$

Equation (31) holds because $\forall \mathbf{r} \in \mathbb{R}^{|S|} : \max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} = \max_{\mathbf{a} \in A'} \mathbf{a}^\top \mathbf{r}$ by lemma E.12's item 2 with $X := A, X' := A'$. Equation (32) holds by lemma E.17. Equation (33) holds by the definition of B' . Furthermore, our assumption on ϕ guarantees that $B' \subseteq B$. Therefore, $\max_{\mathbf{b} \in B'} \mathbf{b}^\top \mathbf{r} \leq \max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r}$, and so eq. (34) holds by the fact that g is an increasing function. Then eq. (30) holds.

If $\text{ND}(B) \setminus B'$ is empty, then $\text{ND}(B) \subseteq B'$. By assumption, $B' \subseteq B$. Then apply lemma E.12 item 2 with $X := B, X' := B'$ in order to conclude that eq. (34) is an equality. Then eq. (30) is also an equality.

Suppose that g is strictly increasing, $\text{ND}(B) \setminus B'$ is non-empty, and $\exists b < c : (b, c)^{|S|} \subseteq \text{supp}(\mathcal{D}_{\text{bound}})$. Let $\mathbf{x} \in \text{ND}(B) \setminus B'$.

$$\mathbb{E}_{\mathbf{r} \sim \phi \cdot \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B'} \mathbf{b}^\top \mathbf{r} \right) \right] < \mathbb{E}_{\mathbf{r} \sim \phi \cdot \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in B' \cup \{\mathbf{x}\}} \mathbf{a}^\top \mathbf{r} \right) \right] \quad (35)$$

$$\leq \mathbb{E}_{\mathbf{r} \sim \phi \cdot \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right]. \quad (36)$$

\mathbf{x} is strictly optimal for a positive-probability subset of $\text{supp}(\mathcal{D}_{\text{bound}})$ by proposition E.20. Since g is strictly increasing, eq. (35) is strict. Therefore, we conclude that eq. (30) is strict. \square

Lemma E.22 (For continuous IID distributions $\mathcal{D}_{X\text{-IID}}$, $\exists b < c : (b, c)^{|S|} \subseteq \text{supp}(\mathcal{D}_{X\text{-IID}})$).

Proof. $\mathcal{D}_{X\text{-IID}} := X^{|S|}$. Since the state reward distribution X is continuous, X must have support on some open interval (b, c) . Since $\mathcal{D}_{X\text{-IID}}$ is IID across states, $(b, c)^{|S|} \subseteq \text{supp}(\mathcal{D}_{X\text{-IID}})$. \square

Definition E.23 (Bounded, continuous IID reward). $\mathcal{D}_{C/B/\text{IID}}$ is the set of $\mathcal{D}_{X\text{-IID}}$ which equal $X^{|S|}$ for some continuous, bounded-support distribution X over \mathbb{R} .

Lemma E.24 (Expectation superiority lemma). *Let $A, B \subseteq \mathbb{R}^{|S|}$ be finite and let $g : \mathbb{R} \rightarrow \mathbb{R}$ be an increasing function. If B contains a copy B' of $\text{ND}(A)$ via ϕ , then*

$$\mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right] \leq_{\text{most: } \mathcal{D}_{\text{bound}}} \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right]. \quad (37)$$

Furthermore, if g is strictly increasing and $\text{ND}(B) \setminus \phi \cdot \text{ND}(A)$ is non-empty, then eq. (37) is strict for all $\mathcal{D}_{X\text{-IID}} \in \mathfrak{D}_{C/B\text{-IID}}$. In particular, $\mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right] \not\leq_{\text{most: } \mathfrak{D}_{\text{bound}}} \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right]$.

Proof. Because $g : \mathbb{R} \rightarrow \mathbb{R}$ is increasing, it is measurable (as is \max). Therefore, the relevant expectations exist for all $\mathcal{D}_{\text{bound}}$.

Suppose that $\mathcal{D}_{\text{bound}}$ is such that $\mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right] < \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right]$.

$$\mathbb{E}_{\mathbf{r} \sim \phi \cdot \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right] \leq \mathbb{E}_{\mathbf{r} \sim \phi^2 \cdot \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right] \quad (38)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right] \quad (39)$$

$$< \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right] \quad (40)$$

$$\leq \mathbb{E}_{\mathbf{r} \sim \phi \cdot \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right]. \quad (41)$$

Equation (38) follows by applying lemma E.21 with permutation ϕ and $A' := \text{ND}(A)$. Equation (39) follows because involutions satisfy $\phi^{-1} = \phi$, and ϕ^2 is therefore the identity. Equation (40) follows because we assumed that $\mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right] < \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right]$. Equation (41) follows by applying lemma E.21 with permutation ϕ and $A' := \text{ND}(A)$. By lemma E.16, eq. (37) holds.

Suppose g is strictly increasing and $\text{ND}(B) \setminus B'$ is non-empty. Let $\phi' \in S_{|S|}$.

$$\mathbb{E}_{\mathbf{r} \sim \phi' \cdot \mathcal{D}_{X\text{-IID}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right] = \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{X\text{-IID}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right] \quad (42)$$

$$< \mathbb{E}_{\mathbf{r} \sim \phi' \cdot \mathcal{D}_{X\text{-IID}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right] \quad (43)$$

$$= \mathbb{E}_{\mathbf{r} \sim \phi' \cdot \mathcal{D}_{X\text{-IID}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right]. \quad (44)$$

Equation (42) and eq. (44) hold because $\mathcal{D}_{X\text{-IID}}$ distributes reward identically across states: $\forall \phi_x \in S_{|S|} : \phi_x \cdot \mathcal{D}_{X\text{-IID}} = \mathcal{D}_{X\text{-IID}}$. By lemma E.22, $\exists b < c : (b, c)^{|S|} \subseteq \text{supp}(\mathcal{D}_{X\text{-IID}})$. Therefore, apply lemma E.21 with $A' := \text{ND}(A)$ to conclude that eq. (43) holds.

Therefore, $\forall \phi' \in S_{|S|} : \mathbb{E}_{\mathbf{r} \sim \phi' \cdot \mathcal{D}_{X\text{-IID}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right] < \mathbb{E}_{\mathbf{r} \sim \phi' \cdot \mathcal{D}_{X\text{-IID}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right]$, and so $\mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{a} \in A} \mathbf{a}^\top \mathbf{r} \right) \right] \not\leq_{\text{most: } \mathfrak{D}_{\text{bound}}} \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[g \left(\max_{\mathbf{b} \in B} \mathbf{b}^\top \mathbf{r} \right) \right]$ by definition 6.5. \square

Definition E.25 (Indicator function). Let L be a predicate which takes input x . $\mathbb{1}_{L(x)}$ is the function which returns 1 when $L(x)$ is true, and 0 otherwise.

Lemma E.26 (Optimality probability inclusion relations). Let $X, Y \subseteq \mathbb{R}^{|S|}$ be finite and suppose $Y' \subseteq Y$.

$$p_{\mathcal{D}_{\text{any}}} (X \geq Y) \leq p_{\mathcal{D}_{\text{any}}} (X \geq Y') \leq p_{\mathcal{D}_{\text{any}}} (X \cup (Y \setminus Y') \geq Y). \quad (45)$$

If $\exists b < c : (b, c)^{|S|} \subseteq \text{supp}(\mathcal{D}_{\text{any}})$, $X \subseteq Y$, and $\text{ND}(Y) \cap (Y \setminus Y')$ is non-empty, then the second inequality is strict.

Proof.

$$p_{\mathcal{D}_{\text{any}}}(X \geq Y) := \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{any}}} \left[\mathbb{1}_{\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} \geq \max_{\mathbf{y} \in Y} \mathbf{y}^\top \mathbf{r}} \right] \quad (46)$$

$$\leq \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{any}}} \left[\mathbb{1}_{\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} \geq \max_{\mathbf{y} \in Y'} \mathbf{y}^\top \mathbf{r}} \right] \quad (47)$$

$$\leq \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{any}}} \left[\mathbb{1}_{\max_{\mathbf{x} \in X \cup (Y \setminus Y')} \mathbf{x}^\top \mathbf{r} \geq \max_{\mathbf{y} \in Y'} \mathbf{y}^\top \mathbf{r}} \right] \quad (48)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{any}}} \left[\mathbb{1}_{\max_{\mathbf{x} \in X \cup (Y \setminus Y')} \mathbf{x}^\top \mathbf{r} \geq \max_{\mathbf{y} \in Y' \cup (Y \setminus Y')} \mathbf{y}^\top \mathbf{r}} \right] \quad (49)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{any}}} \left[\mathbb{1}_{\max_{\mathbf{x} \in X \cup (Y \setminus Y')} \mathbf{x}^\top \mathbf{r} \geq \max_{\mathbf{y} \in Y} \mathbf{y}^\top \mathbf{r}} \right] \quad (50)$$

$$=: p_{\mathcal{D}_{\text{any}}}(X \cup (Y \setminus Y') \geq Y). \quad (51)$$

Equation (47) follows because $\forall \mathbf{r} \in \mathbb{R}^{|S|} : \mathbb{1}_{\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} \geq \max_{\mathbf{y} \in Y} \mathbf{y}^\top \mathbf{r}} \leq \mathbb{1}_{\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} \geq \max_{\mathbf{y} \in Y'} \mathbf{y}^\top \mathbf{r}}$ since $Y' \subseteq Y$; note that eq. (47) equals $p_{\mathcal{D}_{\text{any}}}(X \geq Y')$, and so the first inequality of eq. (45) is shown. Equation (48) holds because $\forall \mathbf{r} \in \mathbb{R}^{|S|} : \mathbb{1}_{\max_{\mathbf{x} \in X} \mathbf{x}^\top \mathbf{r} \geq \max_{\mathbf{y} \in Y'} \mathbf{y}^\top \mathbf{r}} \leq \mathbb{1}_{\max_{\mathbf{x} \in X \cup (Y \setminus Y')} \mathbf{x}^\top \mathbf{r} \geq \max_{\mathbf{y} \in Y'} \mathbf{y}^\top \mathbf{r}}$.

Suppose $\exists b < c : (b, c)^{|S|} \subseteq \text{supp}(\mathcal{D}_{\text{any}})$, $X \subseteq Y$, and $\text{ND}(Y) \cap (Y \setminus Y')$ is non-empty. Let $\mathbf{y}^* \in \text{ND}(Y) \cap (Y \setminus Y')$. By proposition E.20, \mathbf{y}^* is strictly optimal on a subset of $\text{supp}(\mathcal{D}_{\text{any}})$ with positive measure under \mathcal{D}_{any} . In particular, for a set of \mathbf{r}^* with positive measure under \mathcal{D}_{any} , we have $\mathbf{y}^{*\top} \mathbf{r}^* > \max_{\mathbf{y} \in Y'} \mathbf{y}^\top \mathbf{r}^*$.

Then eq. (48) is strict, and therefore the second inequality of eq. (45) is strict as well. \square

Lemma E.27 (Optimality probability of similar linear functional sets). *Let $A, B, C \subsetneq \mathbb{R}^{|S|}$ be finite, and let $Z \subseteq \mathbb{R}^{|S|}$ be such that $\text{ND}(C) \subseteq Z \subseteq C$. If $\text{ND}(A)$ is similar to $B' \subseteq B$ via ϕ such that $\phi \cdot (Z \setminus (B \setminus B')) = Z \setminus (B \setminus B')$, then*

$$p_{\mathcal{D}_{\text{any}}}(A \geq C) \leq p_{\phi \cdot \mathcal{D}_{\text{any}}}(B \geq C). \quad (52)$$

If $B' = B$, then eq. (52) is an equality. If $\exists b < c : (b, c)^{|S|} \subseteq \text{supp}(\mathcal{D}_{\text{any}})$, $B' \subseteq C$, and $\text{ND}(C) \cap (B \setminus B')$ is non-empty, then eq. (52) is strict.

Proof.

$$p_{\mathcal{D}_{\text{any}}}(A \geq C) = p_{\mathcal{D}_{\text{any}}}(A \geq Z) \quad (53)$$

$$= p_{\mathcal{D}_{\text{any}}}(\text{ND}(A) \geq Z) \quad (54)$$

$$\leq p_{\mathcal{D}_{\text{any}}}(\text{ND}(A) \geq Z \setminus (B \setminus B')) \quad (55)$$

$$= p_{\phi \cdot \mathcal{D}_{\text{any}}}(\phi \cdot \text{ND}(A) \geq \phi \cdot Z \setminus (B \setminus B')) \quad (56)$$

$$= p_{\phi \cdot \mathcal{D}_{\text{any}}}(B' \geq Z \setminus (B \setminus B')) \quad (57)$$

$$\leq p_{\phi \cdot \mathcal{D}_{\text{any}}}(B' \cup (B \setminus B') \geq Z) \quad (58)$$

$$= p_{\phi \cdot \mathcal{D}_{\text{any}}}(B \geq C). \quad (59)$$

Equation (53) and eq. (59) follow by lemma E.12's item 2 with $X := C$, $X' := Z$. Similarly, eq. (54) follows by lemma E.12's item 2 with $X := A$, $X' := \text{ND}(A)$. Equation (55) follows by applying the first inequality of lemma E.26 with $X := \text{ND}(A)$, $Y := Z$, $Y' := Z \setminus (B \setminus B')$. Equation (56) follows by applying lemma E.17 to eq. (53) with permutation ϕ .

Equation (57) follows by our assumptions on ϕ . Equation (58) follows because by applying the second inequality of lemma E.26 with $X := B'$, $Y := \text{ND}(C)$, $Y' := \text{ND}(C) \setminus (B \setminus B')$.

Suppose $B' = B$. Then $B \setminus B' = \emptyset$, and so eq. (55) and eq. (58) are trivially equalities. Then eq. (52) is an equality.

Suppose $\exists b < c : (b, c)^{|S|} \subseteq \text{supp}(\mathcal{D}_{\text{any}})$; note that $(b, c)^{|S|} \subseteq \text{supp}(\phi \cdot \mathcal{D}_{\text{any}})$, since such support must be invariant to permutation. Further suppose that $B' \subseteq C$ and that $\text{ND}(C) \cap (B \setminus B')$ is non-empty. Then letting $X := B'$, $Y := Z$, $Y' := Z \setminus (B \setminus B')$ and noting that $\text{ND}(\text{ND}(Z)) = \text{ND}(Z)$, apply lemma E.26 to eq. (58) to conclude that eq. (52) is strict. \square

Lemma E.28 (Optimality probability superiority lemma). *Let $A, B, C \subseteq \mathbb{R}^{|S|}$ be finite, and let Z satisfy $\text{ND}(C) \subseteq Z \subseteq C$. If B contains a copy B' of $\text{ND}(A)$ via ϕ such that $\phi \cdot (Z \setminus (B \setminus B')) = Z \setminus (B \setminus B')$, then $p_{\mathcal{D}_{\text{any}}}(A \geq C) \leq_{\text{most: } \mathcal{D}_{\text{any}}} p_{\mathcal{D}_{\text{any}}}(B \geq C)$.*

If $B' \subseteq C$ and $\text{ND}(C) \cap (B \setminus B')$ is non-empty, then the inequality is strict for all $\mathcal{D}_{X\text{-IID}} \in \mathcal{D}_{C/B\text{-IID}}$ and $p_{\mathcal{D}_{\text{any}}}(A \geq C) \not\leq_{\text{most: } \mathcal{D}_{\text{any}}} p_{\mathcal{D}_{\text{any}}}(B \geq C)$.

Proof. Suppose \mathcal{D}_{any} is such that $p_{\mathcal{D}_{\text{any}}}(B \geq C) < p_{\mathcal{D}_{\text{any}}}(A \geq C)$.

$$p_{\phi \cdot \mathcal{D}_{\text{any}}}(A \geq C) = p_{\phi^{-1} \cdot \mathcal{D}_{\text{any}}}(A \geq C) \quad (60)$$

$$\leq p_{\mathcal{D}_{\text{any}}}(B \geq C) \quad (61)$$

$$< p_{\mathcal{D}_{\text{any}}}(A \geq C) \quad (62)$$

$$\leq p_{\phi \cdot \mathcal{D}_{\text{any}}}(B \geq C). \quad (63)$$

Equation (60) holds because ϕ is an involution. Equation (61) and eq. (63) hold by applying lemma E.27 with permutation ϕ . Equation (62) holds by assumption. Therefore, $p_{\mathcal{D}_{\text{any}}}(A \geq C) \leq_{\text{most: } \mathcal{D}_{\text{any}}} p_{\mathcal{D}_{\text{any}}}(B \geq C)$ by lemma E.16.

Suppose $B' \subseteq C$ and $\text{ND}(C) \cap (B \setminus B')$ is non-empty, and let $\mathcal{D}_{X\text{-IID}}$ be any continuous distribution which distributes reward independently and identically across states. Let $\phi' \in S_{|S|}$.

$$p_{\phi' \cdot \mathcal{D}_{X\text{-IID}}}(A \geq C) = p_{\mathcal{D}_{X\text{-IID}}}(A \geq C) \quad (64)$$

$$< p_{\phi' \cdot \mathcal{D}_{X\text{-IID}}}(B \geq C) \quad (65)$$

$$= p_{\phi' \cdot \mathcal{D}_{X\text{-IID}}}(A \geq C). \quad (66)$$

Equation (64) and eq. (66) hold because $\mathcal{D}_{X\text{-IID}}$ distributes reward identically across states, $\forall \phi_x \in S_{|S|} : \phi_x \cdot \mathcal{D}_{X\text{-IID}} = \mathcal{D}_{X\text{-IID}}$. By lemma E.22, $\exists b < c : (b, c)^{|S|} \subseteq \text{supp}(\mathcal{D}_{X\text{-IID}})$. Therefore, apply lemma E.27 to conclude that eq. (65) holds.

Therefore, $\forall \phi' \in S_{|S|} : p_{\phi' \cdot \mathcal{D}_{X\text{-IID}}}(A \geq C) < p_{\phi' \cdot \mathcal{D}_{X\text{-IID}}}(B \geq C)$. In particular, $p_{\mathcal{D}_{\text{any}}}(A \geq C) \not\leq_{\text{most: } \mathcal{D}_{\text{any}}} p_{\mathcal{D}_{\text{any}}}(B \geq C)$ by definition 6.5. \square

Lemma E.29 (Limit probability inequalities which hold for most distributions). *Let $I \subseteq \mathbb{R}$, let $\mathcal{D} \subseteq \Delta(\mathbb{R}^{|S|})$ be closed under permutation, and let F_A, F_B, F_C be finite sets of vector functions $I \mapsto \mathbb{R}^{|S|}$. Let γ be a limit point of I such that $f_1(\mathcal{D}) := \lim_{\gamma^* \rightarrow \gamma} p_{\mathcal{D}}(F_B(\gamma^*) \geq F_C(\gamma^*))$, $f_2(\mathcal{D}) := \lim_{\gamma^* \rightarrow \gamma} p_{\mathcal{D}}(F_A(\gamma^*) \geq F_C(\gamma^*))$ are well-defined for all $\mathcal{D} \in \mathcal{D}$.*

Let F_Z satisfy $\text{ND}(F_C) \subseteq F_Z \subseteq F_C$. Suppose F_B contains a copy of F_A via ϕ such that $\phi \cdot (F_Z \setminus (F_B \setminus \phi \cdot F_A)) = F_Z \setminus (F_B \setminus \phi \cdot F_A)$. Then $f_2(\mathcal{D}) \leq_{\text{most: } \mathcal{D}} f_1(\mathcal{D})$.

Proof. Suppose $\mathcal{D} \in \mathcal{D}$ is such that $f_2(\mathcal{D}) > f_1(\mathcal{D})$.

$$f_2(\phi \cdot \mathcal{D}) = f_2(\phi^{-1} \cdot \mathcal{D}) \quad (67)$$

$$:= \lim_{\gamma^* \rightarrow \gamma} p_{\phi^{-1} \cdot \mathcal{D}}(F_A(\gamma^*) \geq F_C(\gamma^*)) \quad (68)$$

$$\leq \lim_{\gamma^* \rightarrow \gamma} p_{\mathcal{D}}(F_B(\gamma^*) \geq F_C(\gamma^*)) \quad (69)$$

$$< \lim_{\gamma^* \rightarrow \gamma} p_{\mathcal{D}}(F_A(\gamma^*) \geq F_C(\gamma^*)) \quad (70)$$

$$\leq \lim_{\gamma^* \rightarrow \gamma} p_{\phi \cdot \mathcal{D}} (F_B(\gamma^*) \geq F_C(\gamma^*)) \quad (71)$$

$$=: f_1(\phi \cdot \mathcal{D}). \quad (72)$$

By the assumption that \mathfrak{D} is closed under permutation and f_2 is well-defined for all $\mathcal{D} \in \mathfrak{D}$, $f_2(\phi \cdot \mathcal{D})$ is well-defined. Equation (67) follows since $\phi = \phi^{-1}$ because ϕ is an involution. For all $\gamma^* \in I$, let $A := F_A(\gamma^*)$, $B := F_B(\gamma^*)$, $C := F_C(\gamma^*)$, $Z := F_Z(\gamma^*)$ (by definition E.13, $\text{ND}(C) \subseteq Z \subseteq C$). Since $\phi \cdot A \subseteq B$ by assumption, and since $\text{ND}(A) \subseteq A$, B also contains a copy of $\text{ND}(A)$ via ϕ . Furthermore, $\phi \cdot (Z \setminus (B \setminus \phi \cdot A)) = Z \setminus (B \setminus \phi \cdot A)$ (by assumption), and so apply lemma E.27 to conclude that $p_{\phi^{-1} \cdot \mathcal{D}} (F_A(\gamma^*) \geq F_C(\gamma^*)) \leq p_{\mathcal{D}} (F_B(\gamma^*) \geq F_C(\gamma^*))$. Therefore, the limit inequality eq. (69) holds. Equation (70) follows because we assumed that $f_1(\mathcal{D}) < f_2(\mathcal{D})$. Equation (71) holds by reasoning similar to that given for eq. (69).

Therefore, $f_2(\mathcal{D}) > f_1(\mathcal{D})$ implies that $f_2(\phi \cdot \mathcal{D}) < f_1(\phi \cdot \mathcal{D})$, and so apply lemma E.16 to conclude that $f_2(\mathcal{D}) \leq_{\text{most: } \mathfrak{D}} f_1(\mathcal{D})$. \square

E.1.3 \mathcal{F}_{nd} results

Proposition E.30 (How to transfer optimal policy sets across discount rates). *Suppose reward function R has optimal policy set $\Pi^*(R, \gamma)$ at discount rate $\gamma \in (0, 1)$. For any $\gamma^* \in (0, 1)$, we can construct a reward function R' such that $\Pi^*(R', \gamma^*) = \Pi^*(R, \gamma)$. Furthermore, $V_{R'}^*(\cdot, \gamma^*) = V_R^*(\cdot, \gamma)$.*

Proof. Let R be any reward function. Suppose $\gamma^* \in (0, 1)$ and construct $R'(s) := V_R^*(s, \gamma) - \gamma^* \max_{a \in \mathcal{A}} \mathbb{E}_{s' \sim T(s, a)} [V_R^*(s', \gamma)]$.

Let $\pi \in \Pi$ be any policy. By the definition of optimal policies, $\pi \in \Pi^*(R', \gamma^*)$ iff for all s :

$$R'(s) + \gamma^* \mathbb{E}_{s' \sim T(s, \pi(s))} [V_{R'}^*(s', \gamma^*)] = R'(s) + \gamma^* \max_{a \in \mathcal{A}} \mathbb{E}_{s' \sim T(s, a)} [V_{R'}^*(s', \gamma^*)] \quad (73)$$

$$R'(s) + \gamma^* \mathbb{E}_{s' \sim T(s, \pi(s))} [V_R^*(s', \gamma)] = R'(s) + \gamma^* \max_{a \in \mathcal{A}} \mathbb{E}_{s' \sim T(s, a)} [V_R^*(s', \gamma)] \quad (74)$$

$$\gamma^* \mathbb{E}_{s' \sim T(s, \pi(s))} [V_R^*(s', \gamma)] = \gamma^* \max_{a \in \mathcal{A}} \mathbb{E}_{s' \sim T(s, a)} [V_R^*(s', \gamma)] \quad (75)$$

$$\mathbb{E}_{s' \sim T(s, \pi(s))} [V_R^*(s', \gamma)] = \max_{a \in \mathcal{A}} \mathbb{E}_{s' \sim T(s, a)} [V_R^*(s', \gamma)]. \quad (76)$$

By the Bellman equations, $R'(s) = V_{R'}^*(s, \gamma^*) - \gamma^* \max_{a \in \mathcal{A}} \mathbb{E}_{s' \sim T(s, a)} [V_{R'}^*(s', \gamma^*)]$. By the definition of R' , $V_{R'}^*(\cdot, \gamma^*) = V_R^*(\cdot, \gamma)$ must be the unique solution to the Bellman equations for R' at γ^* . Therefore, eq. (74) holds. Equation (75) follows by plugging in $R' := V_R^*(s, \gamma) - \gamma^* \max_{a \in \mathcal{A}} \mathbb{E}_{s' \sim T(s, a)} [V_R^*(s', \gamma)]$ to eq. (74) and doing algebraic manipulation. Equation (76) follows because $\gamma^* > 0$.

Equation (76) shows that $\pi \in \Pi^*(R', \gamma^*)$ iff $\forall s : \mathbb{E}_{s' \sim T(s, \pi(s))} [V_R^*(s', \gamma)] = \max_{a \in \mathcal{A}} \mathbb{E}_{s' \sim T(s, a)} [V_R^*(s', \gamma)]$. That is, $\pi \in \Pi^*(R', \gamma^*)$ iff $\pi \in \Pi^*(R, \gamma)$. \square

Definition E.31 (Evaluating sets of visit distribution functions at γ). For $\gamma \in (0, 1)$, define $\mathcal{F}(s, \gamma) := \{\mathbf{f}(\gamma) \mid \mathbf{f} \in \mathcal{F}(s)\}$ and $\mathcal{F}_{\text{nd}}(s, \gamma) := \{\mathbf{f}(\gamma) \mid \mathbf{f} \in \mathcal{F}_{\text{nd}}(s)\}$. If $F \subseteq \mathcal{F}(s)$, then $F(\gamma) := \{\mathbf{f}(\gamma) \mid \mathbf{f} \in F\}$.

Lemma E.32 (Non-domination across γ values for expectations of visit distributions). *Let $\Delta_d \in \Delta(\mathbb{R}^{|S|})$ be any state distribution and let $F := \{\mathbb{E}_{s_d \sim \Delta_d} [\mathbf{f}^{\pi, s_d}] \mid \pi \in \Pi\}$. $\mathbf{f} \in \text{ND}(F)$ iff $\forall \gamma^* \in (0, 1) : \mathbf{f}(\gamma^*) \in \text{ND}(F(\gamma^*))$.*

Proof. Let $\mathbf{f}^\pi \in \text{ND}(F)$ be strictly optimal for reward function R at discount rate $\gamma \in (0, 1)$:

$$\mathbf{f}^\pi(\gamma)^\top \mathbf{r} > \max_{\mathbf{f}^{\pi'} \in F \setminus \{\mathbf{f}^\pi\}} \mathbf{f}^{\pi'}(\gamma)^\top \mathbf{r}. \quad (77)$$

Let $\gamma^* \in (0, 1)$. By proposition E.30, we can produce R' such that $\Pi^*(R', \gamma^*) = \Pi^*(R, \gamma)$. Since the optimal policy sets are equal, lemma E.1 implies that

$$\mathbf{f}^\pi(\gamma^*)^\top \mathbf{r}' > \max_{\mathbf{f}^{\pi'} \in F \setminus \{\mathbf{f}^\pi\}} \mathbf{f}^{\pi'}(\gamma^*)^\top \mathbf{r}'. \quad (78)$$

Therefore, $\mathbf{f}^\pi(\gamma^*) \in \text{ND}(F(\gamma^*))$.

The reverse direction follows by the definition of $\text{ND}(F)$. \square

Lemma E.33 ($\forall \gamma \in (0, 1) : \mathbf{d} \in \mathcal{F}_{\text{nd}}(s, \gamma)$ iff $\mathbf{d} \in \text{ND}(\mathcal{F}(s, \gamma))$).

Proof. By definition E.31, $\mathcal{F}_{\text{nd}}(s, \gamma) := \{\mathbf{f}(\gamma) \mid \mathbf{f} \in \text{ND}(\mathcal{F}(s))\}$. By applying lemma E.32 with $\Delta_d := \mathbf{e}_s$, $\mathbf{f} \in \text{ND}(\mathcal{F}(s))$ iff $\forall \gamma \in (0, 1) : \mathbf{f}(\gamma) \in \text{ND}(\mathcal{F}(s, \gamma))$. \square

Lemma E.34 ($\forall \gamma \in [0, 1) : V_R^*(s, \gamma) = \max_{\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)} \mathbf{f}(\gamma)^\top \mathbf{r}$).

Proof. $\text{ND}(\mathcal{F}(s, \gamma)) = \mathcal{F}_{\text{nd}}(s, \gamma)$ by lemma E.33, so apply corollary E.11 with $X := \mathcal{F}(s, \gamma)$. \square

E.2 Some actions have greater probability of being optimal

Lemma E.35 (Optimal policy shift bound). *For fixed R , $\Pi^*(R, \gamma)$ can take on at most $(2|\mathcal{S}| + 1) \sum_s \binom{|\mathcal{F}(s)|}{2}$ distinct values over $\gamma \in (0, 1)$.*

Proof. By lemma E.1, $\Pi^*(R, \gamma)$ changes value iff there is a change in optimality status for some visit distribution function at some state. Lippman [1968] showed that two visit distribution functions can trade off optimality status at most $2|\mathcal{S}| + 1$ times. At each state s , there are $\binom{|\mathcal{F}(s)|}{2}$ such pairs. \square

Proposition E.36 (Optimality probability's limits exist). *Let $F \subseteq \mathcal{F}(s)$. $\mathbb{P}_{\mathcal{D}_{\text{any}}}(F, 0) = \lim_{\gamma \rightarrow 0} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F, \gamma)$ and $\mathbb{P}_{\mathcal{D}_{\text{any}}}(F, 1) = \lim_{\gamma \rightarrow 1} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F, \gamma)$.*

Proof. First consider the limit as $\gamma \rightarrow 1$. Let \mathcal{D}_{any} have probability measure F_{any} , and define $\delta(\gamma) := F_{\text{any}}(\{R \in \mathbb{R}^{\mathcal{S}} \mid \exists \gamma^* \in [\gamma, 1) : \Pi^*(R, \gamma^*) \neq \Pi^*(R, 1)\})$. Since F_{any} is a probability measure, $\delta(\gamma)$ is bounded $[0, 1]$, and $\delta(\gamma)$ is monotone decreasing. Therefore, $\lim_{\gamma \rightarrow 1} \delta(\gamma)$ exists.

If $\lim_{\gamma \rightarrow 1} \delta(\gamma) > 0$, then there exist reward functions whose optimal policy sets $\Pi^*(R, \gamma)$ never converge (in the discrete topology on sets) to $\Pi^*(R, 1)$, contradicting lemma E.35. So $\lim_{\gamma \rightarrow 1} \delta(\gamma) = 0$.

By the definition of optimality probability (definition 4.3) and of $\delta(\gamma)$, $|\mathbb{P}_{\mathcal{D}_{\text{any}}}(F, \gamma) - \mathbb{P}_{\mathcal{D}_{\text{any}}}(F, 1)| \leq \delta(\gamma)$. Since $\lim_{\gamma \rightarrow 1} \delta(\gamma) = 0$, $\lim_{\gamma \rightarrow 1} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F, \gamma) = \mathbb{P}_{\mathcal{D}_{\text{any}}}(F, 1)$.

A similar proof shows that $\lim_{\gamma \rightarrow 0} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F, \gamma) = \mathbb{P}_{\mathcal{D}_{\text{any}}}(F, 0)$. \square

Lemma E.37 (Optimality probability identity). *Let $\gamma \in (0, 1)$ and let $F \subseteq \mathcal{F}(s)$.*

$$\mathbb{P}_{\mathcal{D}_{\text{any}}}(F, \gamma) = p_{\mathcal{D}'}(F(\gamma) \geq \mathcal{F}(s, \gamma)) = p_{\mathcal{D}'}(F(\gamma) \geq \mathcal{F}_{\text{nd}}(s, \gamma)). \quad (79)$$

Proof. Let $\gamma \in (0, 1)$.

$$\mathbb{P}_{\mathcal{D}_{\text{any}}}(F, \gamma) := \mathbb{P}_{R \sim \mathcal{D}_{\text{any}}}(\exists \mathbf{f}^\pi \in F : \pi \in \Pi^*(R, \gamma)) \quad (80)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{any}}} \left[\mathbb{1}_{\max_{\mathbf{f} \in F} \mathbf{f}(\gamma)^\top \mathbf{r} = \max_{\mathbf{f}' \in \mathcal{F}(s)} \mathbf{f}'(\gamma)^\top \mathbf{r}} \right] \quad (81)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{any}}} \left[\mathbb{1}_{\max_{\mathbf{f} \in F} \mathbf{f}(\gamma)^\top \mathbf{r} = \max_{\mathbf{f}' \in \mathcal{F}_{\text{nd}}(s)} \mathbf{f}'(\gamma)^\top \mathbf{r}} \right] \quad (82)$$

$$=: p_{\mathcal{D}'}(F(\gamma) \geq \mathcal{F}_{\text{nd}}(s, \gamma)). \quad (83)$$

Equation (81) follows because lemma E.1 shows that π is optimal iff it induces an optimal visit distribution \mathbf{f} at every state. Equation (82) follows because $\forall \mathbf{r} \in \mathbb{R}^{|\mathcal{S}|} : \max_{\mathbf{f}' \in \mathcal{F}(s)} \mathbf{f}'(\gamma)^\top \mathbf{r} = \max_{\mathbf{f}' \in \mathcal{F}_{\text{nd}}(s)} \mathbf{f}'(\gamma)^\top \mathbf{r}$ by lemma E.34. \square

E.3 Basic properties of POWER

Lemma E.38 (POWER identities). *Let $\gamma \in (0, 1)$.*

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) = \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)} \frac{1-\gamma}{\gamma} (\mathbf{f}(\gamma) - \mathbf{e}_s)^\top \mathbf{r} \right] \quad (84)$$

$$= \frac{1-\gamma}{\gamma} \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} [V_R^*(s, \gamma) - R(s)] \quad (85)$$

$$= \frac{1-\gamma}{\gamma} \left(V_{\mathcal{D}_{\text{bound}}}^*(s, \gamma) - \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} [R(s)] \right) \quad (86)$$

$$= \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} \left[\max_{\pi \in \Pi} \mathbb{E}_{s' \sim T(s, \pi(s))} [(1-\gamma) V_R^\pi(s', \gamma)] \right]. \quad (87)$$

Proof.

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) := \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{f} \in \mathcal{F}(s)} \frac{1-\gamma}{\gamma} (\mathbf{f}(\gamma) - \mathbf{e}_s)^\top \mathbf{r} \right] \quad (88)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)} \frac{1-\gamma}{\gamma} (\mathbf{f}(\gamma) - \mathbf{e}_s)^\top \mathbf{r} \right] \quad (89)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{f} \in \mathcal{F}(s)} \frac{1-\gamma}{\gamma} (\mathbf{f}(\gamma) - \mathbf{e}_s)^\top \mathbf{r} \right] \quad (90)$$

$$= \frac{1-\gamma}{\gamma} \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} [V_R^*(s, \gamma) - R(s)] \quad (91)$$

$$= \frac{1-\gamma}{\gamma} \left(V_{\mathcal{D}_{\text{bound}}}^*(s, \gamma) - \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} [R(s)] \right) \quad (92)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\pi \in \Pi} \mathbb{E}_{s' \sim T(s, \pi(s))} [(1-\gamma) \mathbf{f}^{\pi, s'}(\gamma)^\top \mathbf{r}] \right] \quad (93)$$

$$= \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} \left[\max_{\pi \in \Pi} \mathbb{E}_{s' \sim T(s, \pi(s))} [(1-\gamma) V_R^\pi(s', \gamma)] \right]. \quad (94)$$

Equation (89) follows from lemma E.34. Equation (91) follows from the dual formulation of optimal value functions. Equation (92) holds by the definition of $V_{\mathcal{D}_{\text{bound}}}^*(s, \gamma)$ (definition 5.1). Equation (93) holds because $\mathbf{f}^{\pi, s}(\gamma) = \mathbf{e}_s + \gamma \mathbb{E}_{s' \sim T(s, \pi(s))} [\mathbf{f}^{\pi, s'}(\gamma)]$ by the definition of a visit distribution function (definition 3.3). \square

Definition E.39 (Discount-normalized value function). Let π be a policy, R a reward function, and s a state. For $\gamma \in [0, 1]$, $V_{R, \text{norm}}^\pi(s, \gamma) := \lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) V_R^\pi(s, \gamma^*)$.

Lemma E.40 (Normalized value functions have uniformly bounded derivative). *There exists $K \geq 0$ such that for all reward functions $\mathbf{r} \in \mathbb{R}^{|S|}$, $\sup_{s \in \mathcal{S}, \pi \in \Pi, \gamma \in [0,1]} \left| \frac{d}{d\gamma} V_{R, \text{norm}}^\pi(s, \gamma) \right| \leq K \|\mathbf{r}\|_1$.*

Proof. Let π be any policy, s a state, and R a reward function. Since $V_{R, \text{norm}}^\pi(s, \gamma) = \lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) \mathbf{f}^{\pi, s}(\gamma^*)^\top \mathbf{r}$, $\frac{d}{d\gamma} V_{R, \text{norm}}^\pi(s, \gamma)$ is controlled by the behavior of $\lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) \mathbf{f}^{\pi, s}(\gamma^*)$. We show that this function's gradient is bounded in infinity norm.

By lemma E.4, $\mathbf{f}^{\pi, s}(\gamma)$ is a multivariate rational function on γ . Therefore, for any state s' , $\mathbf{f}^{\pi, s}(\gamma)^\top \mathbf{e}_{s'} = \frac{P(\gamma)}{Q(\gamma)}$ in reduced form. By proposition E.3, $0 \leq \mathbf{f}^{\pi, s}(\gamma)^\top \mathbf{e}_{s'} \leq \frac{1}{1-\gamma}$. Thus, Q may only have a root of multiplicity 1 at $\gamma = 1$, and $Q(\gamma) \neq 0$ for $\gamma \in [0, 1)$. Let $f_{s'}(\gamma) := (1 - \gamma) \mathbf{f}^{\pi, s}(\gamma)^\top \mathbf{e}_{s'}$.

If $Q(1) \neq 0$, then the derivative $f'_{s'}(\gamma)$ is bounded on $\gamma \in [0, 1)$ because the polynomial $(1 - \gamma)P(\gamma)$ cannot diverge on a bounded domain.

If $Q(1) = 0$, then factor out the root as $Q(\gamma) = (1 - \gamma)Q^*(\gamma)$.

$$f'_{s'}(\gamma) = \frac{d}{d\gamma} \left(\frac{(1 - \gamma)P(\gamma)}{Q(\gamma)} \right) \quad (95)$$

$$= \frac{d}{d\gamma} \left(\frac{P(\gamma)}{Q^*(\gamma)} \right) \quad (96)$$

$$= \frac{P'(\gamma)Q^*(\gamma) - (Q^*)'(\gamma)P(\gamma)}{(Q^*(\gamma))^2}. \quad (97)$$

Since $Q^*(\gamma)$ is a polynomial with no roots on $\gamma \in [0, 1]$, $f'_{s'}(\gamma)$ is bounded on $\gamma \in [0, 1)$.

Therefore, whether or not $Q(\gamma)$ has a root at $\gamma = 1$, $f'_{s'}(\gamma)$ is bounded on $\gamma \in [0, 1)$. Furthermore, $\sup_{\gamma \in [0, 1)} \|\nabla(1 - \gamma) \mathbf{f}^{\pi, s}(\gamma)\|_\infty = \sup_{\gamma \in [0, 1)} \max_{s' \in \mathcal{S}} |f'_{s'}(\gamma)|$ is finite since there are only finitely many states.

There are finitely many $\pi \in \Pi$, and finitely many states s , and so there exists some K' such that $\sup_{\substack{s \in \mathcal{S}, \\ \pi \in \Pi, \gamma \in [0, 1)}} \|\nabla(1 - \gamma) \mathbf{f}^{\pi, s}(\gamma)\|_\infty \leq K'$. Then $\|\nabla(1 - \gamma) \mathbf{f}^{\pi, s}(\gamma)\|_1 \leq |\mathcal{S}| K' =: K$.

$$\sup_{\substack{s \in \mathcal{S}, \\ \pi \in \Pi, \gamma \in [0, 1)}} \left| \frac{d}{d\gamma} V_{R, \text{norm}}^\pi(s, \gamma) \right| := \sup_{\pi \in \Pi, \gamma \in [0, 1)} \left| \frac{d}{d\gamma} \lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) V_R^\pi(s, \gamma^*) \right| \quad (98)$$

$$= \sup_{\substack{s \in \mathcal{S}, \\ \pi \in \Pi, \gamma \in [0, 1)}} \left| \frac{d}{d\gamma} (1 - \gamma) V_R^\pi(s, \gamma) \right| \quad (99)$$

$$= \sup_{\substack{s \in \mathcal{S}, \\ \pi \in \Pi, \gamma \in [0, 1)}} \left| \nabla(1 - \gamma) \mathbf{f}^{\pi, s}(\gamma)^\top \mathbf{r} \right| \quad (100)$$

$$\leq \sup_{\substack{s \in \mathcal{S}, \\ \pi \in \Pi, \gamma \in [0, 1)}} \|\nabla(1 - \gamma) \mathbf{f}^{\pi, s}(\gamma)\|_1 \|\mathbf{r}\|_1 \quad (101)$$

$$\leq K \|\mathbf{r}\|_1. \quad (102)$$

Equation (99) holds because $V_R^\pi(s, \gamma)$ is continuous on $\gamma \in [0, 1)$ by corollary E.5. Equation (101) holds by the Cauchy-Schwarz inequality.

Since $\left| \frac{d}{d\gamma} V_{R, \text{norm}}^\pi(s, \gamma) \right|$ is bounded for all $\gamma \in [0, 1)$, eq. (102) also holds for $\gamma \rightarrow 1$. \square

Lemma 5.3 (Continuity of POWER). *POWER_{D_{bound}}(s, γ) is Lipschitz continuous on $\gamma \in [0, 1]$.*

Proof. Let b, c be such that $\text{supp}(\mathcal{D}_{\text{bound}}) \subseteq [b, c]^{|S|}$. For any $\mathbf{r} \in \text{supp}(\mathcal{D}_{\text{bound}})$ and $\pi \in \Pi$, $V_{R, \text{norm}}^\pi(s, \gamma)$ has Lipschitz constant $K \|\mathbf{r}\|_1 \leq K |\mathcal{S}| \|\mathbf{r}\|_\infty \leq K |\mathcal{S}| \max(|c|, |b|)$ on $\gamma \in (0, 1)$ by lemma E.40.

For $\gamma \in (0, 1)$, $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) = \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} \left[\max_{\pi \in \Pi} \mathbb{E}_{s' \sim T(s, \pi(s))} \left[(1 - \gamma) V_R^\pi(s', \gamma) \right] \right]$ by eq. (94). The expectation of the maximum of a set of functions which share a Lipschitz constant, also shares the Lipschitz constant. This shows that $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma)$ is Lipschitz continuous on $\gamma \in (0, 1)$. Thus, its limits are well-defined as $\gamma \rightarrow 0$ and $\gamma \rightarrow 1$. So it is Lipschitz continuous on the closed unit interval. \square

Proposition 5.4 (Maximal POWER). $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) \leq \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} [\max_{s \in \mathcal{S}} R(s)]$, with equality if s can deterministically reach all states in one step and all states are 1-cycles.

Proof. Let $\gamma \in (0, 1)$.

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) = \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} \left[\max_{\pi \in \Pi} \mathbb{E}_{s' \sim T(s, \pi(s))} \left[(1 - \gamma) V_R^*(s', \gamma) \right] \right] \quad (103)$$

$$\leq \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} \left[\max_{\pi \in \Pi} \mathbb{E}_{s' \sim T(s, \pi(s))} \left[(1 - \gamma) \frac{\max_{s'' \in \mathcal{S}} R(s'')}{1 - \gamma} \right] \right] \quad (104)$$

$$= \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} \left[\max_{s'' \in \mathcal{S}} R(s'') \right]. \quad (105)$$

Equation (103) follows from lemma E.38. Equation (104) follows because $V_R^*(s', \gamma) \leq \frac{\max_{s'' \in \mathcal{S}} R(s'')}{1 - \gamma}$, as no policy can do better than achieving maximal reward at each time step. Taking limits, the inequality holds for all $\gamma \in [0, 1]$.

Suppose that s can deterministically reach all states in one step and all states are 1-cycles. Then eq. (104) is an equality for all $\gamma \in (0, 1)$, since for each R , the agent can select an action which deterministically transitions to a state with maximal reward. Thus the equality holds for all $\gamma \in [0, 1]$. \square

Lemma E.41 (Lower bound on current POWER based on future POWER).

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) \geq (1 - \gamma) \min_{\substack{a \\ R \sim \mathcal{D}_{\text{bound}}}} \mathbb{E}_{s' \sim T(s, a)} [R(s')] + \gamma \max_{\substack{a \\ R \sim \mathcal{D}_{\text{bound}}}} \mathbb{E}_{s' \sim T(s, a)} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)]. \quad (106)$$

Proof. Let $\gamma \in (0, 1)$ and let $a^* \in \arg \max_a \mathbb{E}_{s' \sim T(s, a)} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)]$.

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) \quad (107)$$

$$= (1 - \gamma) \mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} \left[\max_a \mathbb{E}_{s' \sim T(s, a)} [V_R^*(s', \gamma)] \right] \quad (108)$$

$$\geq (1 - \gamma) \max_a \mathbb{E}_{s' \sim T(s, a)} \left[\mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} [V_R^*(s', \gamma)] \right] \quad (109)$$

$$= (1 - \gamma) \max_a \mathbb{E}_{s' \sim T(s, a)} [V_{\mathcal{D}_{\text{bound}}}^*(s', \gamma)] \quad (110)$$

$$= (1 - \gamma) \max_a \mathbb{E}_{s' \sim T(s, a)} \left[\mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} [R(s')] + \frac{\gamma}{1 - \gamma} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma) \right] \quad (111)$$

$$\geq (1 - \gamma) \mathbb{E}_{s' \sim T(s, a^*)} \left[\mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} [R(s')] + \frac{\gamma}{1 - \gamma} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma) \right] \quad (112)$$

$$\geq (1 - \gamma) \min_{\substack{a \\ R \sim \mathcal{D}_{\text{bound}}}} \mathbb{E}_{s' \sim T(s, a)} [R(s')] + \gamma \mathbb{E}_{s' \sim T(s, a^*)} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)]. \quad (113)$$

Equation (108) holds by lemma E.38. Equation (109) follows because $\mathbb{E}_{x \sim X} [\max_a f(a, x)] \geq \max_a \mathbb{E}_{x \sim X} [f(a, x)]$ by Jensen's inequality, and eq. (111) follows by lemma E.38.

The inequality also holds when we take the limits $\gamma \rightarrow 0$ or $\gamma \rightarrow 1$. \square

Proposition 5.5 (POWER is smooth across reversible dynamics). *Let $\mathcal{D}_{\text{bound}}$ be bounded $[b, c]$. Suppose s and s' can both reach each other in one step with probability 1.*

$$|\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) - \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)| \leq (c - b)(1 - \gamma). \quad (3)$$

Proof. Suppose $\gamma \in [0, 1]$. First consider the case where $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) \geq \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)$.

$$\begin{aligned} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma) &\geq (1 - \gamma) \min_{\substack{a \\ s_x \sim T(s', a), \\ R \sim \mathcal{D}_{\text{bound}}}} \mathbb{E} [R(s_x)] + \gamma \max_{\substack{a \\ s_x \sim T(s', a)}} \mathbb{E} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_x, \gamma)] \\ &\geq (1 - \gamma)b + \gamma \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma). \end{aligned} \quad (114)$$

$$\geq (1 - \gamma)b + \gamma \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma). \quad (115)$$

Equation (114) follows by lemma E.41. Equation (115) follows because reward is lower-bounded by b and because s' can reach s in one step with probability 1.

$$\begin{aligned} |\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) - \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)| &= \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) - \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma) \quad (116) \\ &\leq \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) - ((1 - \gamma)b + \gamma \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma)) \quad (117) \end{aligned}$$

$$= (1 - \gamma) (\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) - b) \quad (118)$$

$$\leq (1 - \gamma) \left(\mathbb{E}_{R \sim \mathcal{D}_{\text{bound}}} \left[\max_{s'' \in \mathcal{S}} R(s'') \right] - b \right) \quad (119)$$

$$\leq (1 - \gamma)(c - b). \quad (120)$$

Equation (116) follows because $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) \geq \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)$. Equation (117) follows by eq. (115). Equation (119) follows by proposition 5.4. Equation (120) follows because reward under $\mathcal{D}_{\text{bound}}$ is upper-bounded by c .

The case where $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) \leq \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)$ is similar, leveraging the fact that s can also reach s' in one step with probability 1. \square

E.4 Seeking POWER is often more probable under optimality

E.4.1 Keeping options open tends to be POWER-seeking and tends to be optimal

Definition E.42 (Normalized visit distribution function). Let $\mathbf{f} : [0, 1] \rightarrow \mathbb{R}^{|\mathcal{S}|}$ be a vector function. For $\gamma \in [0, 1]$, $\text{NORM}(\mathbf{f}, \gamma) := \lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) \mathbf{f}(\gamma^*)$ (this limit need not exist for arbitrary \mathbf{f}). If F is a set of such \mathbf{f} , then $\text{NORM}(F, \gamma) := \{\text{NORM}(\mathbf{f}, \gamma) \mid \mathbf{f} \in F\}$.

Remark. $\text{RSD}(s) = \text{NORM}(\mathcal{F}(s), 1)$.

Lemma E.43 (Normalized visit distribution functions are continuous). *Let $\Delta_s \in \Delta(\mathcal{S})$ be a state probability distribution, let $\pi \in \Pi$, and let $\mathbf{f}^* := \mathbb{E}_{s \sim \Delta_s} [\mathbf{f}^{\pi, s}]$. $\text{NORM}(\mathbf{f}^*, \gamma)$ is continuous on $\gamma \in [0, 1]$.*

Proof.

$$\text{NORM}(\mathbf{f}^*, \gamma) := \lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) \mathbb{E}_{s \sim \Delta_s} [\mathbf{f}^{\pi, s}(\gamma^*)] \quad (121)$$

$$= \mathbb{E}_{s \sim \Delta_s} \left[\lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) \mathbf{f}^{\pi, s}(\gamma^*) \right] \quad (122)$$

$$=: \mathbb{E}_{s \sim \Delta_s} [\text{NORM}(\mathbf{f}^{\pi, s}, \gamma)]. \quad (123)$$

Equation (122) follows because the expectation is over a finite set. Each $\mathbf{f}^{\pi, s} \in \mathcal{F}(s)$ is continuous on $\gamma \in [0, 1]$ by lemma E.4, and $\lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) \mathbf{f}^{\pi, s}(\gamma^*)$ exists because RSDs are well-defined [Puterman, 2014]. Therefore, each $\text{NORM}(\mathbf{f}^{\pi, s}, \gamma)$ is continuous on $\gamma \in [0, 1]$. Lastly, eq. (123)'s expectation over finitely many continuous functions is itself continuous. \square

Lemma E.44 (Non-domination of normalized visit distribution functions). *Let $\Delta_s \in \Delta(\mathcal{S})$ be a state probability distribution and let $F := \{\mathbb{E}_{s \sim \Delta_s}[\mathbf{f}^{\pi, s}] \mid \pi \in \Pi\}$. For all $\gamma \in [0, 1]$, $\text{ND}(\text{NORM}(F, \gamma)) \subseteq \text{NORM}(\text{ND}(F), \gamma)$, with equality when $\gamma \in (0, 1)$.*

Proof. Suppose $\gamma \in (0, 1)$.

$$\text{ND}(\text{NORM}(F, \gamma)) = \text{ND}((1 - \gamma)F(\gamma)) \quad (124)$$

$$= (1 - \gamma)\text{ND}(F(\gamma)) \quad (125)$$

$$= (1 - \gamma)(\text{ND}(F)(\gamma)) \quad (126)$$

$$= \text{NORM}(\text{ND}(F), \gamma). \quad (127)$$

Equation (124) and eq. (127) follow by the continuity of $\text{NORM}(\mathbf{f}, \gamma)$ (lemma E.43). Equation (125) follows by lemma E.15 item 1. Equation (126) follows by lemma E.32.

Let $\gamma = 1$. Let $\mathbf{d} \in \text{ND}(\text{NORM}(F, 1))$ be strictly optimal for $\mathbf{r}^* \in \mathbb{R}^{|\mathcal{S}|}$. Then let $F_{\mathbf{d}} \subseteq F$ be the subset of $\mathbf{f} \in F$ such that $\text{NORM}(\mathbf{f}, 1) = \mathbf{d}$.

$$\max_{\mathbf{f} \in F_{\mathbf{d}}} \text{NORM}(\mathbf{f}, 1)^\top \mathbf{r}^* > \max_{\mathbf{f}' \in F \setminus F_{\mathbf{d}}} \text{NORM}(\mathbf{f}', 1)^\top \mathbf{r}^*. \quad (128)$$

Since $\text{NORM}(\mathbf{f}, 1)$ is continuous at $\gamma = 1$ (lemma E.43), $\mathbf{x}^\top \mathbf{r}^*$ is continuous on $\mathbf{x} \in \mathbb{R}^{|\mathcal{S}|}$, and F is finite, eq. (128) holds for some $\gamma^* \in (0, 1)$ sufficiently close to $\gamma = 1$. By lemma E.10, at least one $\mathbf{f} \in F_{\mathbf{d}}$ is an element of $\text{ND}(F(\gamma^*))$. Then by lemma E.32, $\mathbf{f} \in \text{ND}(F)$. We conclude that $\text{ND}(\text{NORM}(F, 1)) \subseteq \text{NORM}(\text{ND}(F), 1)$.

The case for $\gamma = 0$ proceeds similarly. \square

Lemma E.45 (POWER limit identity). *Let $\gamma \in [0, 1]$.*

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) = \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)} \lim_{\gamma^* \rightarrow \gamma} \frac{1 - \gamma^*}{\gamma^*} (\mathbf{f}(\gamma^*) - \mathbf{e}_s)^\top \mathbf{r} \right]. \quad (129)$$

Proof. Let $\gamma \in [0, 1]$.

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) = \lim_{\gamma^* \rightarrow \gamma} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma^*) \quad (130)$$

$$= \lim_{\gamma^* \rightarrow \gamma} \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)} \frac{1 - \gamma^*}{\gamma^*} (\mathbf{f}(\gamma^*) - \mathbf{e}_s)^\top \mathbf{r} \right] \quad (131)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\lim_{\gamma^* \rightarrow \gamma} \max_{\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)} \frac{1 - \gamma^*}{\gamma^*} (\mathbf{f}(\gamma^*) - \mathbf{e}_s)^\top \mathbf{r} \right] \quad (132)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)} \lim_{\gamma^* \rightarrow \gamma} \frac{1 - \gamma^*}{\gamma^*} (\mathbf{f}(\gamma^*) - \mathbf{e}_s)^\top \mathbf{r} \right]. \quad (133)$$

Equation (130) follows because $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma)$ is continuous on $\gamma \in [0, 1]$ by lemma 5.3. Equation (131) follows by lemma E.38.

For $\gamma^* \in (0, 1)$, let $f_{\gamma^*}(\mathbf{r}) := \max_{\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)} \frac{1 - \gamma^*}{\gamma^*} (\mathbf{f}(\gamma^*) - \mathbf{e}_s)^\top \mathbf{r}$. For any sequence $\gamma_n \rightarrow \gamma$, $(f_{\gamma_n})_{n=1}^\infty$ is a sequence of functions which are piecewise linear on $\mathbf{r} \in \mathbb{R}^{|\mathcal{S}|}$, which means they are continuous and therefore measurable. Since lemma E.4 shows that each $\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)$ is multivariate rational on γ^* (and therefore continuous on γ^*), $\{f_{\gamma_n}\}_{n=1}^\infty$ converges pointwise to limit function f_γ .

Furthermore, $|V_R^*(s, \gamma_n) - R(s)| \leq \frac{\gamma}{1 - \gamma_n} \|R\|_\infty$, and so $|f_{\gamma_n}(\mathbf{r})| = \left| \frac{1 - \gamma_n}{\gamma_n} (V_R^*(s, \gamma_n) - R(s)) \right| \leq g(\mathbf{r}) \leq \|R\|_\infty =: g(\mathbf{r})$, which is measurable. Therefore, apply Lebesgue's dominated convergence theorem to conclude that eq. (132) holds. Equation (133) holds because max is a continuous function. \square

Lemma E.46 (Lemma for POWER superiority). *Let $\Delta_1, \Delta_2 \in \Delta(S)$ be state probability distributions. For $i = 1, 2$, let $F_{\Delta_i} := \{\gamma^{-1} \mathbb{E}_{s_i \sim \Delta_i} [\mathbf{f}^{\pi, s_i} - \mathbf{e}_{s_i}] \mid \pi \in \Pi\}$. Suppose F_{Δ_2} contains a copy of $\text{ND}(F_{\Delta_1})$ via ϕ . Then $\forall \gamma \in [0, 1] : \mathbb{E}_{s_1 \sim \Delta_1} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_1, \gamma)] \leq_{\text{most: } \mathcal{D}_{\text{bound}}} \mathbb{E}_{s_2 \sim \Delta_2} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_2, \gamma)]$.*

If $\text{ND}(F_{\Delta_2}) \setminus \phi \cdot \text{ND}(F_{\Delta_1})$ is non-empty, then for all $\gamma \in (0, 1)$, the inequality is strict for all $\mathcal{D}_{X\text{-IID}} \in \mathcal{D}_{C/B/IID}$ and $\mathbb{E}_{s_1 \sim \Delta_1} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_1, \gamma)] \not\leq_{\text{most: } \mathcal{D}_{\text{bound}}} \mathbb{E}_{s_2 \sim \Delta_2} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_2, \gamma)]$.

These results also hold when replacing F_{Δ_i} with $F_{\Delta_i}^ := \{\mathbb{E}_{s_i \sim \Delta_i} [\mathbf{f}^{\pi, s_i}] \mid \pi \in \Pi\}$ for $i = 1, 2$.*

Proof.

$$\phi \cdot \text{ND}(\text{NORM}(F_{\Delta_1}, \gamma)) \subseteq \phi \cdot \text{NORM}(\text{ND}(F_{\Delta_1}), \gamma) \quad (134)$$

$$:= \left\{ \mathbf{P}_\phi \lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) \mathbf{f}(\gamma^*) \mid \mathbf{f} \in \text{ND}(F_{\Delta_1}) \right\} \quad (135)$$

$$= \left\{ \lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) \mathbf{P}_\phi \mathbf{f}(\gamma^*) \mid \mathbf{f} \in \text{ND}(F_{\Delta_1}) \right\} \quad (136)$$

$$= \left\{ \lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) \mathbf{f}(\gamma^*) \mid \mathbf{f} \in F'_{\text{sub}} \right\} \quad (137)$$

$$\subseteq \left\{ \lim_{\gamma^* \rightarrow \gamma} (1 - \gamma^*) \mathbf{f}(\gamma^*) \mid \mathbf{f} \in F_{\Delta_2} \right\} \quad (138)$$

$$=: \text{NORM}(F_{\Delta_2}, \gamma). \quad (139)$$

Equation (134) follows by lemma E.44. Equation (136) follows because \mathbf{P}_ϕ is a continuous linear operator. Equation (138) follows by assumption.

$$\mathbb{E}_{s_1 \sim \Delta_1} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_1, \gamma)] := \mathbb{E}_{\substack{s_1 \sim \Delta_1, \\ \mathbf{r} \sim \mathcal{D}_{\text{bound}}}} \left[\max_{\pi \in \Pi} \lim_{\gamma^* \rightarrow \gamma} \frac{1 - \gamma^*}{\gamma^*} (\mathbf{f}^{\pi, s_1}(\gamma^*) - \mathbf{e}_{s_1})^\top \mathbf{r} \right] \quad (140)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\pi \in \Pi} \lim_{\gamma^* \rightarrow \gamma} \frac{1 - \gamma^*}{\gamma^*} \mathbb{E}_{s_1 \sim \Delta_1} [\mathbf{f}^{\pi, s_1}(\gamma^*) - \mathbf{e}_{s_1}]^\top \mathbf{r} \right] \quad (141)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{d} \in \text{NORM}(F_{\Delta_1}, \gamma)} \mathbf{d}^\top \mathbf{r} \right] \quad (142)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{d} \in \text{ND}(\text{NORM}(F_{\Delta_1}, \gamma))} \mathbf{d}^\top \mathbf{r} \right] \quad (143)$$

$$\leq_{\text{most: } \mathcal{D}_{\text{bound}}} \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{d} \in \text{NORM}(F_{\Delta_2}, \gamma)} \mathbf{d}^\top \mathbf{r} \right] \quad (144)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\pi \in \Pi} \lim_{\gamma^* \rightarrow \gamma} \frac{1 - \gamma^*}{\gamma^*} \mathbb{E}_{s_2 \sim \Delta_2} [\mathbf{f}^{\pi, s_2}(\gamma^*) - \mathbf{e}_{s_2}]^\top \mathbf{r} \right] \quad (145)$$

$$= \mathbb{E}_{\substack{s_2 \sim \Delta_2, \\ \mathbf{r} \sim \mathcal{D}_{\text{bound}}}} \left[\max_{\pi \in \Pi} \lim_{\gamma^* \rightarrow \gamma} \frac{1 - \gamma^*}{\gamma^*} (\mathbf{f}^{\pi, s_2}(\gamma^*) - \mathbf{e}_{s_2})^\top \mathbf{r} \right] \quad (146)$$

$$=: \mathbb{E}_{s_2 \sim \Delta_2} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_2, \gamma)]. \quad (147)$$

Equation (140) and eq. (147) follow by lemma E.45. Equation (141) and eq. (146) follow because each R has a stationary deterministic optimal policy $\pi \in \Pi^*(R, \gamma) \subseteq \Pi$ which simultaneously achieves optimal value at all states. Equation (143) follows by corollary E.11.

Apply lemma E.24 with $A := \text{NORM}(F_{\Delta_1}, \gamma)$, $B := \text{NORM}(F_{\Delta_2}, \gamma)$, g the identity function, and involution ϕ (satisfying $\phi \cdot \text{ND}(A) \subseteq B$ by eq. (139)) in order to conclude that eq. (144) holds.

Suppose that $\text{ND}(F_{\Delta_2}) \setminus \phi \cdot \text{ND}(F_{\Delta_1})$ is non-empty; let $F'_{\text{sub}} := \phi \cdot \text{ND}(F_{\Delta_1})$. Lemma E.32 shows that for all $\gamma \in (0, 1)$, $\text{ND}(F_{\Delta_2}(\gamma)) \setminus F'_{\text{sub}}(\gamma)$ is non-empty. Lemma E.15 item 1 then implies that $\text{ND}(B) \setminus \phi \cdot A = \frac{1-\gamma}{\gamma} \left(\text{ND}(F_{\Delta_2}(\gamma)) - \mathbf{e}_s \right) \setminus \left(\frac{1-\gamma}{\gamma} F'_{\text{sub}}(\gamma) \right)$ is non-empty. Then lemma E.24 implies that for all $\gamma \in (0, 1)$, eq. (144) is strict for all $\mathcal{D}_{X\text{-IID}} \in \mathfrak{D}_{C/B/\text{IID}}$ and $\mathbb{E}_{s_1 \sim \Delta_1} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_1, \gamma)] \not\leq_{\text{most: } \mathfrak{D}_{\text{bound}}} \mathbb{E}_{s_2 \sim \Delta_2} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_2, \gamma)]$.

We show that this result's preconditions holding for $F_{\Delta_i}^*$ implies the F_{Δ_i} preconditions. Suppose $F_{\Delta_i}^* := \{\mathbb{E}_{s_i \sim \Delta_i} [\mathbf{f}^{\pi, s_i}] \mid \pi \in \Pi\}$ for $i = 1, 2$ are such that $F_{\text{sub}}^* := \phi \cdot \text{ND}(F_{\Delta_1}^*) \subseteq F_{\Delta_2}^*$. In the following, the Δ_i are represented as vectors in $\mathbb{R}^{|S|}$, and γ is a variable.

$$\phi \cdot \{\gamma \mathbf{f} \mid \mathbf{f} \in \text{ND}(F_{\Delta_1})\} = \phi \cdot \left(\text{ND}(F_{\Delta_1}^* - \Delta_1) \right) \quad (148)$$

$$= \phi \cdot \left(\text{ND}(F_{\Delta_1}^*) - \Delta_1 \right) \quad (149)$$

$$= \left\{ \mathbf{P}_\phi \mathbf{f} - \mathbf{P}_\phi \Delta_1 \mid \mathbf{f} \in \text{ND}(F_{\Delta_1}^*) \right\} \quad (150)$$

$$\subseteq \left\{ \mathbf{f} - \Delta_2 \mid \mathbf{f} \in F_{\Delta_2}^* \right\} \quad (151)$$

$$= \left\{ \gamma \mathbf{f} \mid \mathbf{f} \in F_{\Delta_2} \right\}. \quad (152)$$

Equation (149) follows from lemma E.15 item 2. Since we assumed that $\phi \cdot \text{ND}(F_{\Delta_1}^*) \subseteq F_{\Delta_2}^*$, $\phi \cdot \{\Delta_1\} = \phi \cdot \left(\text{ND}(F_{\Delta_1}^*)(0) \right) \subseteq F_{\Delta_2}^*(0) = \{\Delta_2\}$. This implies that $\mathbf{P}_\phi \Delta_1 = \Delta_2$ and so eq. (151) follows.

Equation (152) shows that $\phi \cdot \{\gamma \mathbf{f} \mid \mathbf{f} \in \text{ND}(F_{\Delta_1})\} \subseteq \{\gamma \mathbf{f} \mid \mathbf{f} \in F_{\Delta_2}\}$. But we then have $\phi \cdot \{\gamma \mathbf{f} \mid \mathbf{f} \in \text{ND}(F_{\Delta_1})\} := \{\gamma \mathbf{P}_\phi \mathbf{f} \mid \mathbf{f} \in \text{ND}(F_{\Delta_1})\} = \{\gamma \mathbf{f} \mid \mathbf{f} \in \phi \cdot \text{ND}(F_{\Delta_1})\} \subseteq \{\gamma \mathbf{f} \mid \mathbf{f} \in F_{\Delta_2}\}$. Thus, $\phi \cdot \text{ND}(F_{\Delta_1}) \subseteq F_{\Delta_2}$.

Suppose $\text{ND}(F_{\Delta_2}^*) \setminus \phi \cdot \text{ND}(F_{\Delta_1}^*)$ is non-empty, which implies that

$$\phi \cdot \{\gamma \mathbf{f} \mid \mathbf{f} \in \text{ND}(F_{\Delta_1})\} = \left\{ \mathbf{P}_\phi \mathbf{f} - \mathbf{P}_\phi \Delta_1 \mid \mathbf{f} \in \text{ND}(F_{\Delta_1}^*) \right\} \quad (153)$$

$$= \left\{ \mathbf{f} - \mathbf{P}_\phi \Delta_1 \mid \mathbf{f} \in \phi \cdot \text{ND}(F_{\Delta_1}^*) \right\} \quad (154)$$

$$\subsetneq \left\{ \mathbf{f} - \Delta_2 \mid \mathbf{f} \in \text{ND}(F_{\Delta_2}^*) \right\} \quad (155)$$

$$= \left\{ \gamma \mathbf{f} \mid \mathbf{f} \in \text{ND}(F_{\Delta_2}) \right\}. \quad (156)$$

Then $\text{ND}(F_{\Delta_2}) \setminus \phi \cdot \text{ND}(F_{\Delta_1})$ must be non-empty. Therefore, if the preconditions of this result are met for $F_{\Delta_i}^*$, they are met for F_{Δ_i} . \square

Proposition 6.6 (States with “more options” have more POWER). *If $\mathcal{F}(s)$ contains a copy of $\mathcal{F}_{\text{nd}}(s')$ via ϕ , then $\forall \gamma \in [0, 1] : \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma) \geq_{\text{most}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma)$. If $\mathcal{F}_{\text{nd}}(s) \setminus \phi \cdot \mathcal{F}_{\text{nd}}(s')$ is non-empty, then for all $\gamma \in (0, 1)$, the converse \leq_{most} statement does not hold.*

Proof. Let $F_{\text{sub}} := \phi \cdot \mathcal{F}_{\text{nd}}(s') \subseteq \mathcal{F}(s)$. Let $\Delta_1 := \mathbf{e}_{s'}$, $\Delta_2 := \mathbf{e}_s$, and define $F_{\Delta_i}^* := \{\mathbb{E}_{s_i \sim \Delta_i} [\mathbf{f}^{\pi, s_i}] \mid \pi \in \Pi\}$ for $i = 1, 2$. Then $\mathcal{F}_{\text{nd}}(s') = \text{ND}(F_{\Delta_1}^*)$ is similar to $F_{\text{sub}} = F_{\text{sub}}^* \subseteq F_{\Delta_2}^* = \mathcal{F}(s)$ via involution ϕ . Apply lemma E.46 to conclude that $\forall \gamma \in [0, 1] : \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma) \leq_{\text{most: } \mathfrak{D}_{\text{bound}}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma)$.

Furthermore, $\mathcal{F}_{\text{nd}}(s) = \text{ND}(F_{\Delta_2}^*)$, and $F_{\text{sub}} = F_{\text{sub}}^*$, and so if $\mathcal{F}_{\text{nd}}(s) \setminus \phi \cdot \mathcal{F}_{\text{nd}}(s') := \mathcal{F}_{\text{nd}}(s) \setminus F_{\text{sub}} = \text{ND}(F_{\Delta_2}^*) \setminus F_{\text{sub}}^*$ is non-empty, then lemma E.46 shows that for all $\gamma \in (0, 1)$, the inequality is strict for all $\mathcal{D}_{X\text{-IID}} \in \mathfrak{D}_{C/B/\text{IID}}$ and $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s', \gamma) \not\leq_{\text{most: } \mathfrak{D}_{\text{bound}}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, \gamma)$. \square

Lemma E.47 (Non-dominated visit distribution functions never agree with other visit distribution functions at that state). *Let $\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)$, $\mathbf{f}' \in \mathcal{F}(s) \setminus \{\mathbf{f}\}$. $\forall \gamma \in (0, 1) : \mathbf{f}(\gamma) \neq \mathbf{f}'(\gamma)$.*

Proof. Let $\gamma \in (0, 1)$. Since $\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)$, there exists a $\gamma^* \in (0, 1)$ at which \mathbf{f} is strictly optimal for some reward function. Then by proposition E.30, we can produce another reward function for which \mathbf{f} is strictly optimal at discount rate γ ; in particular, proposition E.30 guarantees that the policies which induce \mathbf{f}' are not optimal at γ . So $\mathbf{f}(\gamma) \neq \mathbf{f}'(\gamma)$. \square

Corollary E.48 (Cardinality of non-dominated visit distributions). *Let $F \subseteq \mathcal{F}(s)$. $\forall \gamma \in (0, 1) : |F \cap \mathcal{F}_{\text{nd}}(s)| = |F(\gamma) \cap \mathcal{F}_{\text{nd}}(s, \gamma)|$.*

Proof. Let $\gamma \in (0, 1)$. By applying lemma E.32 with $\Delta_d := \mathbf{e}_s$, $\mathbf{f} \in \mathcal{F}_{\text{nd}}(s) = \text{ND}(\mathcal{F}(s))$ iff $\mathbf{f}(\gamma) \in \text{ND}(\mathcal{F}(s, \gamma))$. By lemma E.33, $\text{ND}(\mathcal{F}(s, \gamma)) = \mathcal{F}_{\text{nd}}(s, \gamma)$. So all $\mathbf{f} \in F \cap \mathcal{F}_{\text{nd}}(s)$ induce $\mathbf{f}(\gamma) \in F(\gamma) \cap \mathcal{F}_{\text{nd}}(s, \gamma)$, and $|F \cap \mathcal{F}_{\text{nd}}(s)| \geq |F(\gamma) \cap \mathcal{F}_{\text{nd}}(s, \gamma)|$.

Lemma E.47 implies that for all $\mathbf{f}, \mathbf{f}' \in \mathcal{F}_{\text{nd}}(s)$, $\mathbf{f} = \mathbf{f}'$ iff $\mathbf{f}(\gamma) = \mathbf{f}'(\gamma)$. Therefore, $|F \cap \mathcal{F}_{\text{nd}}(s)| \leq |F(\gamma) \cap \mathcal{F}_{\text{nd}}(s, \gamma)|$. So $|F \cap \mathcal{F}_{\text{nd}}(s)| = |F(\gamma) \cap \mathcal{F}_{\text{nd}}(s, \gamma)|$. \square

Lemma E.49 (Optimality probability and state bottlenecks). *Suppose that s can reach $\text{REACH}(s', a') \cup \text{REACH}(s', a)$, but only by taking actions equivalent to a' or a at state s' . $F_{\text{nd}, a'} := \mathcal{F}_{\text{nd}}(s \mid \pi(s') = a')$, $F_a := \mathcal{F}(s \mid \pi(s') = a)$. Suppose F_a contains a copy of $F_{\text{nd}, a'}$ via ϕ which fixes all states not belonging to $\text{REACH}(s', a') \cup \text{REACH}(s', a)$. Then $\forall \gamma \in [0, 1] : \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_{\text{nd}, a'}, \gamma) \leq_{\text{most}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_a, \gamma)$.*

If $\mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus \phi \cdot F_{\text{nd}, a'})$ is non-empty, then for all $\gamma \in (0, 1)$, the inequality is strict for all $\mathcal{D}_{X\text{-IID}} \in \mathcal{D}_{\text{C/B/IID}}$, and $\mathbb{P}_{\mathcal{D}_{\text{any}}}(F_{\text{nd}, a'}, \gamma) \not\leq_{\text{most}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_a, \gamma)$.

Proof. Let $F_{\text{sub}} := \phi \cdot F_{\text{nd}, a'}$. Let $F^* := \bigcup_{(a'' \neq_{s'} a) \wedge (a'' \neq_{s'} a')}^{a'' \in \mathcal{A}:} \mathcal{F}(s \mid \pi(s') = a'') \cup F_{\text{nd}, a'} \cup F_{\text{sub}}$.

$$\phi \cdot F^* := \phi \cdot \left(\bigcup_{(a'' \neq_{s'} a) \wedge (a'' \neq_{s'} a')}^{a'' \in \mathcal{A}:} \mathcal{F}(s \mid \pi(s') = a'') \cup F_{\text{nd}, a'} \cup F_{\text{sub}} \right) \quad (157)$$

$$= \bigcup_{(a'' \neq_{s'} a) \wedge (a'' \neq_{s'} a')}^{a'' \in \mathcal{A}:} \phi \cdot \mathcal{F}(s \mid \pi(s') = a'') \cup (\phi \cdot F_{\text{nd}, a'}) \cup (\phi \cdot F_{\text{sub}}) \quad (158)$$

$$= \bigcup_{(a'' \neq_{s'} a) \wedge (a'' \neq_{s'} a')}^{a'' \in \mathcal{A}:} \phi \cdot \mathcal{F}(s \mid \pi(s') = a'') \cup F_{\text{sub}} \cup F_{\text{nd}, a'} \quad (159)$$

$$= \bigcup_{(a'' \neq_{s'} a) \wedge (a'' \neq_{s'} a')}^{a'' \in \mathcal{A}:} \mathcal{F}(s \mid \pi(s') = a'') \cup F_{\text{sub}} \cup F_{\text{nd}, a'} \quad (160)$$

$$=: F^*. \quad (161)$$

Equation (159) follows because the involution ϕ ensures that $\phi \cdot F_{\text{sub}} = F_{\text{nd}, a'}$. By assumption, ϕ fixes all $s' \notin \text{REACH}(s', a') \cup \text{REACH}(s', a)$. Suppose $\mathbf{f} \in \mathcal{F}(s) \setminus (F_{\text{nd}, a'} \cup F_a)$. By the bottleneck assumption, \mathbf{f} does not visit states in $\text{REACH}(s', a') \cup \text{REACH}(s', a)$. Therefore, $\mathbf{P}_\phi \mathbf{f} = \mathbf{f}$, and so eq. (160) follows.

Let $F_Z := (\mathcal{F}(s) \setminus (\mathcal{F}(s \mid \pi(s) = a') \cup F_a)) \cup F_{\text{nd}, a'} \cup F_a$. By definition, $F_Z \subseteq \mathcal{F}(s)$. Furthermore, $\mathcal{F}_{\text{nd}}(s) = \bigcup_{a'' \in \mathcal{A}} \mathcal{F}_{\text{nd}}(s \mid \pi(s') = a'') \subseteq (\mathcal{F}(s) \setminus (\mathcal{F}(s \mid \pi(s) = a') \cup F_a)) \cup \mathcal{F}_{\text{nd}}(s \mid \pi(s) = a') \cup F_a =: F_Z$, and so $\mathcal{F}_{\text{nd}}(s) \subseteq F_Z$. Note that $F^* = F_Z \setminus (F_a \setminus F_{\text{sub}})$.

Case: $\gamma \in (0, 1)$.

$$\mathbb{P}_{\mathcal{D}_{\text{any}}}(F_{\text{nd},a'}, \gamma) = p_{\mathcal{D}_{\text{any}}}(F_{\text{nd},a'}(\gamma) \geq \mathcal{F}(s, \gamma)) \quad (162)$$

$$\leq_{\text{most: } \mathfrak{D}_{\text{any}}} p_{\mathcal{D}_{\text{any}}}(F_a(\gamma) \geq \mathcal{F}(s, \gamma)) \quad (163)$$

$$= \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_{\text{nd},a'}, \gamma). \quad (164)$$

Equation (162) and eq. (164) follow from lemma E.37. Equation (163) follows by applying lemma E.28 with $A := F_{\text{nd},a'}(\gamma)$, $B' := F_{\text{sub}}(\gamma)$, $B := F_a(\gamma)$, $C := \mathcal{F}(s, \gamma)$, $Z := F_Z(\gamma)$ which satisfies $\text{ND}(C) = \mathcal{F}_{\text{nd}}(s, \gamma) \subseteq F_Z(\gamma) \subseteq \mathcal{F}(s, \gamma) = C$, and involution ϕ which satisfies $\phi \cdot F^*(\gamma) = \phi \cdot (Z \setminus (B \setminus B')) = Z \setminus (B \setminus B') = F^*(\gamma)$.

Suppose $\mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus F_{\text{sub}})$ is non-empty. $0 < \left| \mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus F_{\text{sub}}) \right| = \left| \mathcal{F}_{\text{nd}}(s, \gamma) \cap (F_a(\gamma) \setminus F_{\text{sub}}(\gamma)) \right| =: \left| \text{ND}(C) \cap (B \setminus B') \right|$ (with the first equality holding by corollary E.48), and so $\text{ND}(C) \cap (B \setminus B')$ is non-empty. We also have $B := F_a(\gamma) \subseteq \mathcal{F}(s, \gamma) =: C$. Then reapplying lemma E.28, eq. (163) is strict for all $\mathcal{D}_{X\text{-IID}} \in \mathfrak{D}_{C/B/\text{IID}}$, and $\mathbb{P}_{\mathcal{D}_{\text{any}}}(F_{\text{nd},a'}, \gamma) \not\leq_{\text{most: } \mathfrak{D}_{\text{any}}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_a, \gamma)$.

Case: $\gamma = 1, \gamma = 0$.

$$\mathbb{P}_{\mathcal{D}_{\text{any}}}(F_{\text{nd},a'}, 1) = \lim_{\gamma^* \rightarrow 1} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_{\text{nd},a'}, \gamma^*) \quad (165)$$

$$= \lim_{\gamma^* \rightarrow 1} p_{\mathcal{D}_{\text{any}}}(F_{\text{nd},a'}(\gamma^*) \geq \mathcal{F}(s, \gamma^*)) \quad (166)$$

$$\leq_{\text{most: } \mathfrak{D}_{\text{any}}} \lim_{\gamma^* \rightarrow 1} p_{\mathcal{D}_{\text{any}}}(F_a(\gamma^*) \geq \mathcal{F}(s, \gamma^*)) \quad (167)$$

$$= \lim_{\gamma^* \rightarrow 1} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_a, \gamma^*) \quad (168)$$

$$= \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_a, 1). \quad (169)$$

Equation (165) and eq. (169) hold by proposition E.36. Equation (166) and eq. (168) follow by lemma E.37. Applying lemma E.29 with $\gamma := 1$, $I := (0, 1)$, $F_A := F_{\text{nd},a'}$, $F_B := F_a$, $F_C := \mathcal{F}(s)$, F_Z as defined above, and involution ϕ (for which $\phi \cdot (F_Z \setminus (F_B \setminus \phi \cdot F_A)) = F_Z \setminus (F_B \setminus \phi \cdot F_A)$), we conclude that eq. (167) follows.

The $\gamma = 0$ case proceeds similarly to $\gamma = 1$. \square

Lemma E.50 (Action optimality probability is a special case of visit distribution optimality probability). $\mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a, \gamma) = \mathbb{P}_{\mathcal{D}_{\text{any}}}(\mathcal{F}(s \mid \pi(s) = a), \gamma)$.

Proof. Let $F_a := \mathcal{F}(s \mid \pi(s) = a)$. For $\gamma \in (0, 1)$,

$$\mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a, \gamma) := \mathbb{P}_{R \sim \mathcal{D}_{\text{any}}}(\exists \pi^* \in \Pi^*(R, \gamma) : \pi^*(s) = a) \quad (170)$$

$$= \mathbb{P}_{\mathbf{r} \sim \mathcal{D}_{\text{any}}} \left(\exists \mathbf{f}^{\pi^*, s} \in F_a : \mathbf{f}^{\pi^*, s}(\gamma)^\top \mathbf{r} = \max_{\mathbf{f} \in \mathcal{F}(s)} \mathbf{f}(\gamma)^\top \mathbf{r} \right) \quad (171)$$

$$= \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_a, \gamma). \quad (172)$$

By lemma E.1, if $\exists \pi^* \in \Pi^*(R, \gamma) : \pi^*(s) = a$, then it induces some optimal $\mathbf{f}^{\pi^*, s} \in F_a$. Conversely, if $\mathbf{f}^{\pi^*, s} \in F_a$ is optimal at $\gamma \in (0, 1)$, then π^* chooses optimal actions on the support of $\mathbf{f}^{\pi^*, s}(\gamma)$. Let π' agree with π^* on that support and let π' take optimal actions at all other states. Then $\pi' \in \Pi^*(R, \gamma)$ and $\pi'(s) = a$. So eq. (171) follows.

Suppose $\gamma = 0$ or $\gamma = 1$. Consider any sequence $(\gamma_n)_{n=1}^\infty$ converging to γ , and let \mathcal{D}_{any} induce probability measure F .

$$\mathbb{P}_{\mathcal{D}_{\text{any}}}(F_a, \gamma) := \lim_{\gamma^* \rightarrow \gamma} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_a, \gamma^*) \quad (173)$$

$$= \lim_{\gamma^* \rightarrow \gamma} \mathbb{P}_{R \sim \mathcal{D}_{\text{any}}} (\exists \pi^* \in \Pi^*(R, \gamma^*) : \pi^*(s) = a) \quad (174)$$

$$= \lim_{n \rightarrow \infty} \mathbb{P}_{R \sim \mathcal{D}_{\text{any}}} (\exists \pi^* \in \Pi^*(R, \gamma_n) : \pi^*(s) = a) \quad (175)$$

$$= \lim_{n \rightarrow \infty} \int_{\mathbb{R}^S} \mathbb{1}_{\exists \pi^* \in \Pi^*(R, \gamma_n) : \pi^*(s) = a} dF(R) \quad (176)$$

$$= \int_{\mathbb{R}^S} \lim_{n \rightarrow \infty} \mathbb{1}_{\exists \pi^* \in \Pi^*(R, \gamma_n) : \pi^*(s) = a} dF(R) \quad (177)$$

$$= \int_{\mathbb{R}^S} \mathbb{1}_{\exists \pi^* \in \Pi^*(R, \gamma) : \pi^*(s) = a} dF(R) \quad (178)$$

$$=: \mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a, \gamma). \quad (179)$$

Equation (174) follows by eq. (172). for $\gamma^* \in [0, 1]$, let $f_{\gamma^*}(R) := \mathbb{1}_{\exists \pi^* \in \Pi^*(R, \gamma^*) : \pi^*(s) = a}$. For each $R \in \mathbb{R}^S$, lemma E.35 exists $\gamma_x \approx \gamma$ such that for all intermediate γ'_x between γ_x and γ , $\Pi^*(R, \gamma'_x) = \Pi^*(R, \gamma)$. Since $\gamma_n \rightarrow \gamma$, this means that $(f_{\gamma_n})_{n=1}^{\infty}$ converges pointwise to f_{γ} . Furthermore, $\forall n \in \mathbb{N}, R \in \mathbb{R}^S : |f_{\gamma_n}(R)| \leq 1$ by definition. Therefore, eq. (177) follows by Lebesgue's dominated convergence theorem. \square

Proposition 6.9 (Keeping options open tends to be POWER-seeking and tends to be optimal).

Suppose $F_a := \mathcal{F}(s \mid \pi(s) = a)$ contains a copy of $F_{a'} := \mathcal{F}(s \mid \pi(s) = a')$ via ϕ .

1. If $s \notin \text{REACH}(s, a')$, then $\forall \gamma \in [0, 1] : \mathbb{E}_{s_a \sim T(s, a)} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_a, \gamma)] \geq_{\text{most}} \mathbb{E}_{s_{a'} \sim T(s, a')} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_{a'}, \gamma)]$.
2. If s can only reach the states of $\text{REACH}(s, a') \cup \text{REACH}(s, a)$ by taking actions equivalent to a' or a at state s , then $\forall \gamma \in [0, 1] : \mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a, \gamma) \geq_{\text{most}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a', \gamma)$.

If $\mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus \phi \cdot F_{a'})$ is non-empty, then $\forall \gamma \in (0, 1)$, the converse \leq_{most} statements do not hold.

Proof. Note that by definition 3.3, $F_{a'}(0) = \{\mathbf{e}_s\} = F_a(0)$. Since $\phi \cdot F_{a'} \subseteq F_a$, in particular we have $\phi \cdot F_{a'}(0) = \{\mathbf{P}_{\phi} \mathbf{e}_s\} \subseteq \{\mathbf{e}_s\} = F_a(0)$, and so $\phi(s) = s$.

Item 1. For state probability distribution $\Delta_s \in \Delta(\mathcal{S})$, let $F_{\Delta_s}^* := \left\{ \mathbb{E}_{s' \sim \Delta_s} [\mathbf{f}^{\pi, s'}] \mid \pi \in \Pi \right\}$.

Unless otherwise stated, we treat γ as a variable in this item; we apply element-wise vector addition, constant multiplication, and variable multiplication via the conventions outlined in definition E.14.

$$F_{a'} = \left\{ \mathbf{e}_s + \gamma \mathbb{E}_{s_{a'} \sim T(s, a')} [\mathbf{f}^{\pi, s_{a'}}] \mid \pi \in \Pi : \pi(s) = a' \right\} \quad (180)$$

$$= \left\{ \mathbf{e}_s + \gamma \mathbb{E}_{s_{a'} \sim T(s, a')} [\mathbf{f}^{\pi, s_{a'}}] \mid \pi \in \Pi \right\} \quad (181)$$

$$= \mathbf{e}_s + \gamma F_{T(s, a')}^*. \quad (182)$$

Equation (180) follows by definition 3.3, since each $\mathbf{f} \in \mathcal{F}(s)$ has an initial term of \mathbf{e}_s . Equation (181) follows because $s \notin \text{REACH}(s, a')$, and so for all $s_{a'} \in \text{supp}(T(s, a'))$, $\mathbf{f}^{\pi, s_{a'}}$ is unaffected by the choice of action $\pi(s)$. Note that similar reasoning implies that $F_a \subseteq \mathbf{e}_s + \gamma F_{T(s, a)}^*$ (because eq. (181) is a containment relation in general).

Since $F_{a'} = \mathbf{e}_s + \gamma F_{T(s, a')}^*$, if F_a contains a copy of $F_{a'}$ via ϕ , then $F_{T(s, a)}^*$ contains a copy of $F_{T(s, a')}^*$ via ϕ . Then $\phi \cdot \text{ND}(F_{T(s, a')}^*) \subseteq \phi \cdot F_{T(s, a')}^* \subseteq F_{T(s, a)}^*$, and so $F_{T(s, a)}^*$ contains a copy of $\text{ND}(F_{T(s, a')}^*)$. Then apply lemma E.46 with $\Delta_1 := T(s, a')$ and $\Delta_2 := T(s, a)$ to conclude that $\forall \gamma \in [0, 1] : \mathbb{E}_{s_{a'} \sim T(s, a')} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_{a'}, \gamma)] \leq_{\text{most}} \mathbb{E}_{s_a \sim T(s, a)} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_a, \gamma)]$.

Suppose $\mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus \phi \cdot F_{a'})$ is non-empty. To apply the second condition of lemma E.46, we want to demonstrate that $\text{ND} \left(F_{T(s,a)}^* \right) \setminus \phi \cdot \text{ND} \left(F_{T(s,a')}^* \right)$ is also non-empty.

First consider $\mathbf{f} \in \mathcal{F}_{\text{nd}}(s) \cap F_a$. Because $F_a \subseteq \mathbf{e}_s + \gamma F_{T(s,a)}^*$, we have that $\gamma^{-1}(\mathbf{f} - \mathbf{e}_s) \in F_{T(s,a)}^*$. Because $\mathbf{f} \in \mathcal{F}_{\text{nd}}(s)$, by definition 3.6, $\exists \mathbf{r} \in \mathbb{R}^{|S|}, \gamma_x \in (0, 1)$ such that

$$\mathbf{f}(\gamma_x)^\top \mathbf{r} > \max_{\mathbf{f}' \in \mathcal{F}(s) \setminus \{\mathbf{f}\}} \mathbf{f}'(\gamma_x)^\top \mathbf{r}. \quad (183)$$

Then since $\gamma_x \in (0, 1)$,

$$\gamma_x^{-1}(\mathbf{f}(\gamma_x) - \mathbf{e}_s)^\top \mathbf{r} > \max_{\mathbf{f}' \in \mathcal{F}(s) \setminus \{\mathbf{f}\}} \gamma_x^{-1}(\mathbf{f}'(\gamma_x) - \mathbf{e}_s)^\top \mathbf{r} \quad (184)$$

$$= \max_{\mathbf{f}' \in \gamma_x^{-1}((\mathcal{F}(s) \setminus \{\mathbf{f}\}) - \mathbf{e}_s)} \mathbf{f}'(\gamma_x)^\top \mathbf{r} \quad (185)$$

$$\geq \max_{\mathbf{f}' \in \gamma_x^{-1}((F_a \setminus \{\mathbf{f}\}) - \mathbf{e}_s)} \mathbf{f}'(\gamma_x)^\top \mathbf{r} \quad (186)$$

$$= \max_{\mathbf{f}' \in F_{T(s,a)}^* \setminus \{\gamma_x^{-1}(\mathbf{f} - \mathbf{e}_s)\}} \mathbf{f}'(\gamma_x)^\top \mathbf{r}. \quad (187)$$

Equation (186) holds because $F_a \subseteq \mathcal{F}(s)$. By assumption, action a is optimal for \mathbf{r} at state s and at discount rate γ_x . Equation (181) shows that $F_{T(s,a)}^*$ potentially allows the agent a non-stationary policy choice at s , but non-stationary policies cannot increase optimal value [Puterman, 2014]. Therefore, eq. (187) holds.

We assumed that $\gamma^{-1}(\mathbf{f} - \mathbf{e}_s) \in \gamma^{-1}(\mathcal{F}_{\text{nd}}(s) - \mathbf{e}_s)$. Furthermore, since we just showed that $\gamma^{-1}(\mathbf{f} - \mathbf{e}_s) \in F_{T(s,a)}^*$ is strictly optimal over the other elements of $F_{T(s,a)}^*$ for reward function \mathbf{r} at discount rate $\gamma_x \in (0, 1)$, we conclude that it is an element of $\text{ND} \left(F_{T(s,a)}^* \right)$ by definition E.13.

Then we conclude that $\gamma^{-1}(\mathcal{F}_{\text{nd}}(s) - \mathbf{e}_s) \cap F_{T(s,a)}^* \subseteq \text{ND} \left(F_{T(s,a)}^* \right)$.

We now show that $\text{ND} \left(F_{T(s,a)}^* \right) \setminus \phi \cdot \text{ND} \left(F_{T(s,a')}^* \right)$ is non-empty.

$$0 < \left| \mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus \phi \cdot F_{a'}) \right| \quad (188)$$

$$= \left| \gamma^{-1} \left(\mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus \phi \cdot F_{a'}) - \mathbf{e}_s \right) \right| \quad (189)$$

$$\leq \left| \gamma^{-1} (\mathcal{F}_{\text{nd}}(s) - \mathbf{e}_s) \cap (F_{T(s,a)}^* \setminus \phi \cdot F_{T(s,a')}^*) \right| \quad (190)$$

$$= \left| \left(\gamma^{-1} (\mathcal{F}_{\text{nd}}(s) - \mathbf{e}_s) \cap F_{T(s,a)}^* \right) \setminus \phi \cdot F_{T(s,a')}^* \right| \quad (191)$$

$$\leq \left| \text{ND} \left(F_{T(s,a)}^* \right) \setminus \phi \cdot F_{T(s,a')}^* \right| \quad (192)$$

$$\leq \left| \text{ND} \left(F_{T(s,a)}^* \right) \setminus \phi \cdot \text{ND} \left(F_{T(s,a')}^* \right) \right|. \quad (193)$$

Equation (188) follows by the assumption that $\mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus \phi \cdot F_{a'})$ is non-empty. Let $\mathbf{f}, \mathbf{f}' \in \mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus \phi \cdot F_{a'})$ be distinct. Then we must have that for some $\gamma_x \in (0, 1)$, $\mathbf{f}(\gamma_x) \neq \mathbf{f}'(\gamma_x)$. This holds iff $\gamma_x^{-1}(\mathbf{f}(\gamma_x) - \mathbf{e}_s) \neq \gamma_x^{-1}(\mathbf{f}'(\gamma_x) - \mathbf{e}_s)$, and so eq. (189) holds.

Equation (190) holds because $F_a \subseteq \mathbf{e}_s + \gamma F_{T(s,a)}^*$ and $F_{a'} = \mathbf{e}_s + \gamma F_{T(s,a')}^*$ by eq. (182). Equation (192) holds because we showed above that $\gamma^{-1}(\mathcal{F}_{\text{nd}}(s) - \mathbf{e}_s) \cap F_{T(s,a)}^* \subseteq \text{ND} \left(F_{T(s,a)}^* \right)$.

Equation (193) holds because $\text{ND} \left(F_{T(s,a')}^* \right) \subseteq F_{T(s,a')}^*$ by definition E.13.

Therefore, $\text{ND} \left(F_{T(s,a)}^* \right) \setminus \phi \cdot \text{ND} \left(F_{T(s,a')}^* \right)$ is non-empty, and so apply the second condition of lemma E.46 to conclude that for all $\mathcal{D}_{X\text{-IID}} \in \mathfrak{D}_{C/B/\text{IID}}, \forall \gamma \in (0, 1) :$

$\mathbb{E}_{s_{a'} \sim T(s, a')} [\text{POWER}_{\mathcal{D}_{X\text{-IID}}}(s_{a'}, \gamma)] < \mathbb{E}_{s_a \sim T(s, a)} [\text{POWER}_{\mathcal{D}_{X\text{-IID}}}(s_a, \gamma)]$, and that $\forall \gamma \in (0, 1) : \mathbb{E}_{s_{a'} \sim T(s, a')} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_{a'}, \gamma)] \not\leq_{\text{most: } \mathcal{D}_{\text{bound}}} \mathbb{E}_{s_a \sim T(s, a)} [\text{POWER}_{\mathcal{D}_{\text{bound}}}(s_a, \gamma)]$.

Item 2. Let $\phi'(s_x) := \phi(s_x)$ when $s_x \in \text{REACH}(s, a') \cup \text{REACH}(s, a)$, and equal s_x otherwise. Since ϕ is an involution, so is ϕ' .

$$\phi' \cdot F_{a'} := \left\{ \mathbf{P}_{\phi'} \left(\mathbf{e}_s + \gamma \mathbb{E}_{s_{a'} \sim T(s, a')} [\mathbf{f}^{\pi, s_{a'}}] \right) \mid \pi \in \Pi, \pi(s) = a' \right\} \quad (194)$$

$$= \left\{ \mathbf{e}_s + \gamma \mathbb{E}_{s_{a'} \sim T(s, a')} [\mathbf{P}_{\phi'} \mathbf{f}^{\pi, s_{a'}}] \mid \pi \in \Pi, \pi(s) = a' \right\} \quad (195)$$

$$= \left\{ \mathbf{P}_{\phi} \mathbf{e}_s + \gamma \mathbb{E}_{s_{a'} \sim T(s, a')} [\mathbf{P}_{\phi} \mathbf{f}^{\pi, s_{a'}}] \mid \pi \in \Pi, \pi(s) = a' \right\} \quad (196)$$

$$=: \phi \cdot F_{a'} \quad (197)$$

$$\subseteq F_a. \quad (198)$$

Equation (195) follows because if $s \in \text{REACH}(s, a') \cup \text{REACH}(s, a)$, then we already showed that ϕ fixes s . Otherwise, $\phi'(s) = s$ by definition. Equation (196) follows by the definition of ϕ' on $\text{REACH}(s, a') \cup \text{REACH}(s, a)$ and because $\mathbf{e}_s = \mathbf{P}_{\phi} \mathbf{e}_s$. Next, we assumed that $\phi \cdot F_{a'} \subseteq F_a$, and so eq. (198) holds.

Therefore, F_a contains a copy of $F_{a'}$ via ϕ' fixing all $s_x \notin \text{REACH}(s, a') \cup \text{REACH}(s, a)$. Therefore, F_a contains a copy of $F_{\text{nd}, a'} := \mathcal{F}_{\text{nd}}(s) \cap F_{a'}$ via the same ϕ' . Then apply lemma E.49 with $s' := s$ to conclude that $\forall \gamma \in [0, 1] : \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_{a'}, \gamma) \leq_{\text{most: } \mathcal{D}_{\text{any}}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_a, \gamma)$. By lemma E.50, $\mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a', \gamma) = \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_{a'}, \gamma)$ and $\mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a, \gamma) = \mathbb{P}_{\mathcal{D}_{\text{any}}}(F_a, \gamma)$. Therefore, $\forall \gamma \in [0, 1] : \mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a', \gamma) \leq_{\text{most: } \mathcal{D}_{\text{any}}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a, \gamma)$.

If $\mathcal{F}_{\text{nd}}(s) \cap (F_a \setminus \phi \cdot F_{a'})$ is non-empty, then apply the second condition of lemma E.49 to conclude that for all $\gamma \in (0, 1)$, the inequality is strict for all $\mathcal{D}_{X\text{-IID}} \in \mathcal{D}_{C/B/IID}$, and $\mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a', \gamma) \not\leq_{\text{most: } \mathcal{D}_{\text{any}}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(s, a, \gamma)$. \square

E.4.2 When $\gamma = 1$, optimal policies tend to navigate towards “larger” sets of cycles

Lemma E.51 (POWER identity when $\gamma = 1$).

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, 1) = \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{d} \in \text{RSD}(s)} \mathbf{d}^\top \mathbf{r} \right] = \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{d} \in \text{RSD}_{\text{nd}}(s)} \mathbf{d}^\top \mathbf{r} \right]. \quad (199)$$

Proof.

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, 1) = \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{f}^{\pi, s} \in \mathcal{F}(s)} \lim_{\gamma \rightarrow 1} \frac{1 - \gamma}{\gamma} (\mathbf{f}^{\pi, s}(\gamma) - \mathbf{e}_s)^\top \mathbf{r} \right] \quad (200)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{d} \in \text{RSD}(s)} \mathbf{d}^\top \mathbf{r} \right] \quad (201)$$

$$= \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{d} \in \text{RSD}_{\text{nd}}(s)} \mathbf{d}^\top \mathbf{r} \right]. \quad (202)$$

Equation (200) follows by lemma E.45. Equation (201) follows by the definition of $\text{RSD}(s)$ (definition 6.10). Equation (202) follows because for all $\mathbf{r} \in \mathbb{R}^{|S|}$, corollary E.11 shows that $\max_{\mathbf{d} \in \text{RSD}(s)} \mathbf{d}^\top \mathbf{r} = \max_{\mathbf{d} \in \text{ND}(\text{RSD}(s))} \mathbf{d}^\top \mathbf{r} =: \max_{\mathbf{d} \in \text{RSD}_{\text{nd}}(s)} \mathbf{d}^\top \mathbf{r}$. \square

Proposition 6.12 (When $\gamma = 1$, RSDs control POWER). *If $\text{RSD}(s)$ contains a copy of $\text{RSD}_{\text{nd}}(s')$ via ϕ , then $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s, 1) \geq_{\text{most}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s', 1)$. If $\text{RSD}_{\text{nd}}(s) \setminus \phi \cdot \text{RSD}_{\text{nd}}(s')$ is non-empty, then the converse \leq_{most} statement does not hold.*

Proof. Suppose $\text{RSD}_{\text{nd}}(s')$ is similar to $D \subseteq \text{RSD}(s)$ via involution ϕ .

$$\text{POWER}_{\mathcal{D}_{\text{bound}}}(s', 1) = \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{d} \in \text{RSD}_{\text{nd}}(s')} \mathbf{d}^\top \mathbf{r} \right] \quad (203)$$

$$\leq_{\text{most: } \mathcal{D}_{\text{bound}}} \mathbb{E}_{\mathbf{r} \sim \mathcal{D}_{\text{bound}}} \left[\max_{\mathbf{d} \in \text{RSD}_{\text{nd}}(s)} \mathbf{d}^\top \mathbf{r} \right] \quad (204)$$

$$= \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, 1) \quad (205)$$

Equation (203) and eq. (205) follow from lemma E.51. By applying lemma E.24 with $A := \text{RSD}(s')$, $B' := D$, $B := \text{RSD}(s)$ and g the identity function, eq. (204) follows.

Suppose $\text{RSD}_{\text{nd}}(s) \setminus D$ is non-empty. By the same result, eq. (204) is a strict inequality for all $\mathcal{D}_{X\text{-IID}} \in \mathcal{D}_{C/B/\text{IID}}$, and we conclude that $\text{POWER}_{\mathcal{D}_{\text{bound}}}(s', 1) \not\leq_{\text{most: } \mathcal{D}_{\text{bound}}} \text{POWER}_{\mathcal{D}_{\text{bound}}}(s, 1)$. \square

Theorem 6.13 (Average-optimal policies tend to end up in “larger” sets of RSDs). *Let $D, D' \subseteq \text{RSD}(s)$. Suppose that D contains a copy of D' via ϕ , and that the sets $D \cup D'$ and $\text{RSD}_{\text{nd}}(s) \setminus (D' \cup D)$ have pairwise orthogonal vector elements (i.e. pairwise disjoint vector support). Then $\mathbb{P}_{\mathcal{D}_{\text{any}}}(D, \text{average}) \geq_{\text{most}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(D', \text{average})$. If $\text{RSD}_{\text{nd}}(s) \cap (D \setminus \phi \cdot D')$ is non-empty, the converse \leq_{most} statement does not hold.*

Proof. Let $D_{\text{sub}} := \phi \cdot D'$, where $D_{\text{sub}} \subseteq D$ by assumption. Let $X := \{s_i \in \mathcal{S} \mid \max_{\mathbf{d} \in D' \cup D} \mathbf{d}^\top \mathbf{e}_{s_i} > 0\}$. Define

$$\phi'(s_i) := \begin{cases} \phi(s_i) & \text{if } s_i \in X \\ s_i & \text{else.} \end{cases} \quad (206)$$

Since ϕ is an involution, ϕ' is also an involution. Furthermore, by the definition of X , $\phi' \cdot D' = D_{\text{sub}}$ and $\phi' \cdot D_{\text{sub}} = D'$ (because we assumed that both equalities hold for ϕ).

Let $D^* := D' \cup D_{\text{sub}} \cup (\text{RSD}_{\text{nd}}(s) \setminus (D' \cup D))$.

$$\phi' \cdot D^* := \phi' \cdot (D' \cup D_{\text{sub}} \cup (\text{RSD}_{\text{nd}}(s) \setminus (D' \cup D))) \quad (207)$$

$$= (\phi' \cdot D') \cup (\phi' \cdot D_{\text{sub}}) \cup \phi' \cdot (\text{RSD}_{\text{nd}}(s) \setminus (D' \cup D)) \quad (208)$$

$$= D_{\text{sub}} \cup D' \cup (\text{RSD}_{\text{nd}}(s) \setminus (D' \cup D)) \quad (209)$$

$$=: D^*. \quad (210)$$

In eq. (209), we know that $\phi' \cdot D' = D_{\text{sub}}$ and $\phi' \cdot D_{\text{sub}} = D'$. We just need to show that $\phi' \cdot (\text{RSD}_{\text{nd}}(s) \setminus (D' \cup D)) = \text{RSD}_{\text{nd}}(s) \setminus (D' \cup D)$.

Suppose $\exists s_i \in X, \mathbf{d}' \in \text{RSD}_{\text{nd}}(s) \setminus (D' \cup D) : \mathbf{d}'^\top \mathbf{e}_{s_i} > 0$. By the definition of X , $\exists \mathbf{d} \in D' \cup D : \mathbf{d}^\top \mathbf{e}_{s_i} > 0$. Then

$$\mathbf{d}^\top \mathbf{d}' = \sum_{j=1}^{|\mathcal{S}|} \mathbf{d}^\top (\mathbf{d}' \odot \mathbf{e}_{s_j}) \quad (211)$$

$$\geq \mathbf{d}^\top (\mathbf{d}' \odot \mathbf{e}_{s_i}) \quad (212)$$

$$= \mathbf{d}^\top ((\mathbf{d}'^\top \mathbf{e}_{s_i}) \mathbf{e}_{s_i}) \quad (213)$$

$$= (\mathbf{d}'^\top \mathbf{e}_{s_i}) \cdot (\mathbf{d}^\top \mathbf{e}_{s_i}) \quad (214)$$

$$> 0. \quad (215)$$

Equation (211) follows from the definitions of the dot and Hadamard products. Equation (212) follows because \mathbf{d} and \mathbf{d}' have non-negative entries. Equation (215) follows because $\mathbf{d}^\top \mathbf{e}_{s_i}$ and $\mathbf{d}'^\top \mathbf{e}_{s_i}$ are both positive. But eq. (215) shows that $\mathbf{d}^\top \mathbf{d}' > 0$, contradicting our assumption that \mathbf{d} and \mathbf{d}' are orthogonal.

Therefore, such an s_i cannot exist, and $X' := \left\{ s'_i \in \mathcal{S} \mid \max_{\mathbf{d}' \in \text{RSD}_{\text{nd}}(s) \setminus (D' \cup D)} \mathbf{d}'^\top \mathbf{e}_{s_i} > 0 \right\} \subseteq (\mathcal{S} \setminus X)$. By eq. (206), $\forall s'_i \in X' : \phi'(s'_i) = s'_i$. Thus, $\phi' \cdot (\text{RSD}_{\text{nd}}(s) \setminus (D' \cup D)) = \text{RSD}_{\text{nd}}(s) \setminus (D' \cup D)$, and eq. (209) follows. We conclude that $\phi' \cdot D^* = D^*$.

Consider $Z := (\text{RSD}_{\text{nd}}(s) \setminus (D' \cup D)) \cup D \cup D'$. First, $Z \subseteq \text{RSD}(s)$ by definition. Second, $\text{RSD}_{\text{nd}}(s) = \text{RSD}_{\text{nd}}(s) \setminus (D' \cup D) \cup (\text{RSD}_{\text{nd}}(s) \cap D') \cup (\text{RSD}_{\text{nd}}(s) \cap D) \subseteq Z$. Note that $D^* = Z \setminus (D \setminus D_{\text{sub}})$.

$$\mathbb{P}_{\mathcal{D}_{\text{any}}}(D', \text{average}) = p_{\mathcal{D}_{\text{any}}}(D' \geq \text{RSD}(s)) \quad (216)$$

$$\leq_{\text{most: } \mathfrak{D}_{\text{any}}} p_{\mathcal{D}_{\text{any}}}(D \geq \text{RSD}(s)) \quad (217)$$

$$= \mathbb{P}_{\mathcal{D}_{\text{any}}}(D, \text{average}). \quad (218)$$

Since $\phi \cdot D' \subseteq D$ and $\text{ND}(D') \subseteq D'$, $\phi \cdot \text{ND}(D') \subseteq D$. Then eq. (217) holds by applying lemma E.28 with $A := D'$, $B' := D_{\text{sub}}$, $B := D$, $C := \text{RSD}(s)$, and the previously defined Z which we showed satisfies $\text{ND}(C) \subseteq Z \subseteq C$. Furthermore, involution ϕ' satisfies $\phi' \cdot B^* = \phi' \cdot (Z \setminus (B \setminus B')) = Z \setminus (B \setminus B') = B^*$ by eq. (210).

When $\text{RSD}_{\text{nd}}(s) \cap (D \setminus D_{\text{sub}})$ is non-empty, since $B' \subseteq C$ by assumption, lemma E.28 also shows that eq. (217) is strict for all $\mathcal{D}_{X\text{-IID}} \in \mathfrak{D}_{C/B/\text{IID}}$, and that $\mathbb{P}_{\mathcal{D}_{\text{any}}}(D', \text{average}) \not\leq_{\text{most: } \mathfrak{D}_{\text{any}}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(D, \text{average})$. \square

Proposition E.52 (RSD properties). *Let $\mathbf{d} \in \text{RSD}(s)$. \mathbf{d} is element-wise non-negative and $\|\mathbf{d}\|_1 = 1$.*

Proof. \mathbf{d} has non-negative elements because it equals the limit of $\lim_{\gamma \rightarrow 1} (1 - \gamma) \mathbf{f}(\gamma)$, whose elements are non-negative by proposition E.3 item 1.

$$\|\mathbf{d}\|_1 = \left\| \lim_{\gamma \rightarrow 1} (1 - \gamma) \mathbf{f}(\gamma) \right\|_1 \quad (219)$$

$$= \lim_{\gamma \rightarrow 1} (1 - \gamma) \|\mathbf{f}(\gamma)\|_1 \quad (220)$$

$$= 1. \quad (221)$$

Equation (219) follows because the definition of RSDs (definition 6.10) ensures that $\exists \mathbf{f} \in \mathcal{F}(s) : \lim_{\gamma \rightarrow 1} (1 - \gamma) \mathbf{f}(\gamma) = \mathbf{d}$. Equation (220) follows because $\|\cdot\|_1$ is a continuous function. Equation (221) follows because $\|\mathbf{f}(\gamma)\|_1 = \frac{1}{1 - \gamma}$ by proposition E.3 item 2. \square

Lemma E.53 (When reachable with probability 1, 1-cycles induce non-dominated RSDs). *If $\mathbf{e}_{s'} \in \text{RSD}(s)$, then $\mathbf{e}_{s'} \in \text{RSD}_{\text{nd}}(s)$.*

Proof. If $\mathbf{d} \in \text{RSD}(s)$ is distinct from $\mathbf{e}_{s'}$, then $\|\mathbf{d}\|_1 = 1$ and \mathbf{d} has non-negative entries by proposition E.52. Since \mathbf{d} is distinct from $\mathbf{e}_{s'}$, then its entry for index s' must be strictly less than 1: $\mathbf{d}^\top \mathbf{e}_{s'} < 1 = \mathbf{e}_{s'}^\top \mathbf{e}_{s'}$. Therefore, $\mathbf{e}_{s'} \in \text{RSD}(s)$ is strictly optimal for the reward function $\mathbf{r} := \mathbf{e}_{s'}$, and so $\mathbf{e}_{s'} \in \text{RSD}_{\text{nd}}(s)$. \square

Corollary 6.14 (Average-optimal policies tend not to end up in any given 1-cycle). *Suppose $\mathbf{e}_{s_x}, \mathbf{e}_{s'} \in \text{RSD}(s)$ are distinct. Then $\mathbb{P}_{\mathcal{D}_{\text{any}}}(\text{RSD}(s) \setminus \{\mathbf{e}_{s_x}\}, \text{average}) \geq_{\text{most}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(\{\mathbf{e}_{s_x}\}, \text{average})$. If there is a third $\mathbf{e}_{s''} \in \text{RSD}(s)$, the converse \leq_{most} statement does not hold.*

Proof. Suppose $\mathbf{e}_{s_x}, \mathbf{e}_{s'} \in \text{RSD}(s)$ are distinct. Let $\phi := (s_x \ s')$, $D' := \{\mathbf{e}_{s_x}\}$, $D := \text{RSD}(s) \setminus \{\mathbf{e}_{s_x}\}$. $\phi \cdot D' = \{\mathbf{e}_{s'}\} \subseteq \text{RSD}(s) \setminus \{\mathbf{e}_{s_x}\} =: D$ since $s_x \neq s'$. $D' \cup D = \text{RSD}(s)$ and $\text{RSD}_{\text{nd}}(s) \setminus (D' \cup D) = \text{RSD}_{\text{nd}}(s) \setminus \text{RSD}(s) = \emptyset$ trivially have pairwise orthogonal vector elements. Then apply theorem 6.13 to conclude that $\mathbb{P}_{\mathcal{D}_{\text{any}}}(\{\mathbf{e}_{s_x}\}, \text{average}) \leq_{\text{most: } \mathfrak{D}_{\text{any}}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(\text{RSD}(s) \setminus \{\mathbf{e}_{s_x}\}, \text{average})$.

Suppose there exists another $\mathbf{e}_{s''} \in \text{RSD}(s)$. By lemma E.53, $\mathbf{e}_{s''} \in \text{RSD}_{\text{nd}}(s)$. Furthermore, since $s'' \notin \{s', s_x\}$, $\mathbf{e}_{s''} \in (\text{RSD}(s) \setminus \{\mathbf{e}_{s_x}\}) \setminus \{\mathbf{e}_{s'}\} = D \setminus \phi \cdot D'$. Therefore,

$\mathbf{e}_{s''} \in \text{RSD}_{\text{nd}}(s) \cap (D \setminus \phi \cdot D')$. Then apply the second condition of theorem 6.13 to conclude that $\mathbb{P}_{\mathcal{D}_{\text{any}}}(\{\mathbf{e}_{s_x}\}, \text{average}) \not\preceq_{\text{most: } \mathfrak{D}_{\text{bound}}} \mathbb{P}_{\mathcal{D}_{\text{any}}}(\text{RSD}(s) \setminus \{\mathbf{e}_{s_x}\}, \text{average})$. \square