

Statistical Inference for Data-adaptive Doubly Robust Estimators with Survival Outcomes

Iván Díaz ^{*1}

¹Division of Biostatistics, Weill Cornell Medicine.

January 30, 2019

Abstract

The consistency of doubly robust estimators relies on consistent estimation of at least one of two nuisance regression parameters. In moderate to large dimensions, the use of flexible data-adaptive regression estimators may aid in achieving this consistency. However, $n^{1/2}$ -consistency of doubly robust estimators is not guaranteed if one of the nuisance estimators is inconsistent. In this paper we present a doubly robust estimator for survival analysis with the novel property that it converges to a Gaussian variable at $n^{1/2}$ -rate for a large class of data-adaptive estimators of the nuisance parameters, under the only assumption that at least one of them is consistently estimated at a $n^{1/4}$ -rate. This result is achieved through adaptation of recent ideas in semiparametric inference, which amount to: (i) Gaussianizing (i.e., making asymptotically linear) a drift term that arises in the asymptotic analysis of the doubly robust estimator, and (ii) using cross-fitting to avoid entropy conditions on the nuisance estimators. We present the formula of the asymptotic variance of the estimator, which allows computation of doubly robust confidence intervals and p-values. We illustrate the finite-sample properties of the estimator in simulation studies, and demonstrate its use in a phase III clinical trial for estimating the effect of a novel therapy for the treatment of HER2 positive breast cancer.

1 Introduction

Doubly robust estimation is a widely used method for the estimation of causal effects and the analysis of missing outcome data. In survival analysis, doubly robust estimation [Díaz et al. \(2018\)](#); [Moore and van der Laan \(2011\)](#); [Parast et al. \(2014\)](#); [Zhang \(2014\)](#); [Cole and Hernán \(2004\)](#); [Xie and Liu \(2005\)](#); [Rotnitzky and Robins \(2005\)](#) proceeds by estimating two nuisance parameters: (i) the probability of treatment as a function of covariates and the probability of censoring conditional on covariates (henceforth referred to as treatment-censoring mechanism), and (ii) the probability of an outcome conditional on covariates (henceforth

^{*}corresponding author: ild2005@med.cornell.edu

referred to as the outcome mechanism). The asymptotic properties of the doubly robust estimator such as consistency and asymptotic distribution thus depends on the large sample behavior of functionals of these nuisance estimators. In low dimensional problems with categorical covariates, nuisance parameter estimation may be carried out using the nonparametric maximum likelihood estimator (NPMLE), and the Delta method yields asymptotic normality of the effect estimates. In moderate to high dimensions or with continuous covariates, the curse of dimensionality precludes the use of the NPMLE, making it necessary to use smoothing techniques. When (semi)-parametric smoothing methods are used (e.g., the Cox proportional hazards model), a simple application of the Delta method yields $n^{1/2}$ -consistency and asymptotic normality of doubly robust estimators, provided that at least one nuisance model is correctly specified. An influence function based approach or the bootstrap may be used to obtain asymptotically valid estimates of the variance, confidence intervals, and p-values. Under a moderate- to high-dimensional regime, the functional forms posed by many (semi)-parametric models are hardly supported by a-priori scientific knowledge, thus yielding inconsistent nuisance and doubly robust estimators. Data-adaptive regression methods have recently been adopted in the missing data and causal inference literature to tackle the problem of model misspecification [van der Laan et al. \(2005\)](#); [van der Laan \(2006\)](#); [Ridgeway and McCaffrey \(2007\)](#); [Bembom et al. \(2008\)](#); [Lee et al. \(2010\)](#); [van der Laan and Starmans \(2014\)](#); [Neugebauer et al. \(2016\)](#); [Belloni et al. \(2014, 2017\)](#); [Farrell \(2015\)](#). Techniques such as classification and regression trees, adaptive splines, neural networks, ℓ_1 regularization, support vector machines, boosting and ensembles, etc. offer a flexibility in functional form specification that is not available for traditional approaches such as the Cox proportional hazards model. However, under inconsistency of one nuisance estimator, the large sample analysis of the resulting data-adaptive doubly robust estimators requires empirical process conditions which are often not verifiable. This means that the finite sample and asymptotic distribution of cannot be established, and standard methods for computing confidence intervals and perform hypothesis testing (such as the bootstrap and influence function based approaches) cannot be guaranteed to be yield correct results.

We develop a doubly robust estimator of the exposure-specific survival curve under informative missingness and a non-randomly assigned exposure. Our estimator is asymptotically normal under the only assumption that the at least one of the nuisance parameters is estimated consistently at $n^{1/4}$ -rate. Our asymptotic analysis of the estimator avoids two empirical process conditions often made in the analysis data-adaptive doubly robust estimators: the Donsker condition (a condition on the entropy of the model) and the condition that a drift term defined as a functional of the nuisance estimators is asymptotically linear and therefore asymptotically Gaussian. To our knowledge, this is the first paper concerned with Gaussianization of this drift term, and with providing doubly-robust asymptotic distributions in the context of survival analysis.

Our work builds on the general framework of targeted learning [van der Laan and Rose \(2011\)](#), and is closely related to the methods in [van der Laan \(2014\)](#); [Benkeser et al. \(2017\)](#); [Díaz and van der Laan \(2017\)](#). These papers present a series of doubly robust estimators of the mean of an outcome from incomplete data in the setting of a cross-sectional study. We show that these ideas are generalizable to estimation with survival outcomes subject to informative right-censoring censoring. Generalizing previous results to more complex data structures has proven non-trivial, because the techniques used involve alternative represen-

tations of the doubly robust estimating equation, and are thus specific to the estimator and data structure considered. Our work provides insights and building blocks that are necessary to generalize the methods to even more complex data structures and doubly robust estimators. To remove the Donsker condition, which may limit the class of data-adaptive estimators allowed, we leverage previous work on cross-fitting, originally proposed in the context of cross-validated targeted minimum loss-based estimation (TMLE) by [Zheng and van der Laan \(2011\)](#), and later applied to the estimating equation approach by [Chernozhukov et al. \(2016\)](#). Our main contribution is to present a method to Gaussianize the drift term, a problem which cross-fitting does not solve.

Related to our methods, [Belloni et al. \(2014, 2017\)](#); [Farrell \(2015\)](#) study doubly robust estimators for cross-sectional studies in high-dimensional settings ($p \gg n$) under the assumption that the functional form of the nuisance parameters (outcome regression and treatment mechanism) can be approximated by a generalized linear model on a known transformation of the covariates. They show that lasso-type methods can be used to obtain estimators that are uniformly asymptotically normal under consistent estimation of both the outcome regression and the treatment mechanism. The work of [Avagyan and Vansteelandt \(2017\)](#); [Dukes et al. \(2018\)](#) extends these methods to obtain inference and tests that remain valid under misspecification of at most one of the nuisance models. The methods of [van der Laan \(2014\)](#); [Benkeser et al. \(2017\)](#); [Díaz and van der Laan \(2017\)](#) and this paper also achieve asymptotic normality under consistent estimation of only one of the working nuisance parameters. Unlike [Belloni et al. \(2014, 2017\)](#); [Farrell \(2015\)](#); [Avagyan and Vansteelandt \(2017\)](#); [Dukes et al. \(2018\)](#), we do not restrict our models to the class of generalized linear functions of the covariates, allowing for general data-adaptive methods such as those based on regression trees, adaptive splines, neural networks, etc.

The article is organized as follows. In Section 2 we introduce the notation and inference problem. In Section 3 we present existing doubly-robust estimators such as the TMLE and augmented inverse probability weighted (AIPW), and discuss their asymptotic properties, focusing on the definition of the “drift term” generated by inconsistent estimation of the nuisance parameters. In Section 4 we present an alternative representation of the drift term, a result that is fundamental to the construction of the repaired TMLE in Section 5. We conclude with a numerical study and an illustrative application in sections 6 and 7, as well as a brief discussion in Section 8.

2 Notation

Let T denote a discrete time-to-event outcome taking values on $\{1, \dots, K\}$. Let $C \in \{0, \dots, K\}$ denote the censoring time defined as the time at which the participant is last observed in the study. Let $A \in \{0, 1\}$ denote study arm assignment, and let W denote a vector of baseline variables which will be used to adjust for the confounding in treatment assignment and for informative censoring. The observed data vector for each participant is $O = (W, A, \Delta, \tilde{T})$, where $\tilde{T} = \min(C, T)$, and $\Delta = \mathbf{1}\{T \leq C\}$ is the indicator that the participant’s event time is observed (uncensored). Here $\mathbf{1}(X)$ is the indicator variable taking value 1 if X is true and 0 otherwise.

Equivalently, we encode a participant’s data vector O using the following longitudinal

data structure:

$$O = (W, A, R_0, L_1, R_1, L_2, \dots, R_{K-1}, L_K), \quad (1)$$

where $R_t = \mathbb{1}\{\tilde{T} = t, \Delta = 0\}$ and $L_t = \mathbb{1}\{\tilde{T} = t, \Delta = 1\}$, for $t \in \{0, \dots, K\}$. The sequence $R_0, L_1, R_1, L_2, \dots, R_{K-1}, L_K$ in the above display consists of all 0's until the first time that either the event is observed or censoring occurs. In the former case $L_t = 1$; otherwise $R_t = 1$. For a random variable X , we denote its history through time t as $\bar{X}_t = (X_0, \dots, X_t)$. For a given scalar x , the expression $\bar{X}_t = x$ denotes element-wise equality. Define the following indicator variables for each t :

$$I_t = \mathbb{1}\{\bar{R}_{t-1} = 0, \bar{L}_{t-1} = 0\}, \quad J_t = \mathbb{1}\{\bar{R}_{t-1} = 0, \bar{L}_t = 0\}.$$

The variable I_t is the indicator based on the data through time $t - 1$ that a participant is at risk of the event being observed at time t . Analogously, J_t is the indicator based on the outcome data through time t and censoring data before time t that a participant is at risk of censoring at time t . By convention we let $J_0 = 1$. We assume that $O \sim P_0$, where P_0 is a distribution in the non-parametric model defined as all distributions dominated by some measure ν . We assume we observe an i.i.d. sample O_1, \dots, O_n from P_0 , and denote \mathbb{P}_n its distribution function. For a function $f : O \mapsto \mathbb{R}$, we use the notation $Pf = \int f dP$.

Define the potential outcome T_1 as the event time that would have been observed had study arm assignment $A = 1$ and censoring time $C = K$ been externally set with probability one. For a given time point τ , we define the counterfactual survival curve under treatment arm $A = 1$ as

$$\theta_0 = P_0(T_1 > \tau).$$

We focus on estimating the survival probability for treatment arm $A = 1$, estimation of $P_0(T_0 > \tau)$, where T_0 as the event time that would have been observed had study arm assignment $A = 0$, may be obtained by symmetric arguments.

Define the conditional hazard function for survival at time t :

$$h(t, w) = P_0(L_t = 1 | I_t = 1, A = 1, W = w),$$

among the population at risk at time t within strata of study arm and baseline variables. Similarly, for the censoring variable C , define the censoring hazard at time $t \in \{0, \dots, K\}$:

$$g_{R,0}(t, w) = P_0(R_t = 1 | J_t = 1, A = 1, W = w).$$

We use the notation $g_{A,0}(w) = P_0(A = 1 | W = w)$ and $g_0(t, w) = (g_{A,0}(w), g_{R,0}(t, w))$. Let p_W denote the marginal distribution of the baseline variables W . We have added the subscript 0 to p_W, g, h to denote the corresponding quantities under P_0 , and will use p_W, g, h without a subscript to refer to generic quantities associated to any P in the non-parametric model. Likewise, we use E_0 to denote expectation under P_0 . Denote the survival function for T at time $t \in \{1, \dots, \tau - 1\}$ conditioned on study arm $A = 1$ and baseline variables w by

$$S_0(t, w) = P_0(T > t | A = 1, W = w). \quad (2)$$

Similarly, define the following function of the censoring distribution:

$$G_0(t, w) = P_0(C \geq t | A = 1, W = w). \quad (3)$$

Define the following assumptions, which are standard in the analysis of survival data under right censoring:

A1 (Consistency). $T = T_1$ in the event $A = 1$;

A2 (Treatment assignment randomization). A is independent of T_1 conditional on W ;

A3 (Random censoring). C is independent of T_1 conditional on $(A = 1, W)$;

A4 (Positivity). $P_0\{g_{A,0}(W) > \epsilon\} = P_0\{g_{R,0}(t, W) < 1 - \epsilon\} = 1$ for some $\epsilon > 0$, for each $t \in \{0, \dots, \tau - 1\}$

We make assumptions **A1-A4** throughout the manuscript. Assumption **A1** connects the potential outcomes to the observed outcome. Assumption **A2** holds by design in a randomized trial. Assumption **A3**, which is similar to that in Rubin (1987), holds if censoring is random within strata of treatment and baseline variables (which we abbreviate as “random censoring”). Assumption **A4** states that each treatment arm has a positive probability, and that every time point has a hazard of censoring smaller than one, within each baseline variable stratum $W = w$ with positive density under P_0 .

Under assumptions **A1-A4**, we have $T \perp\!\!\!\perp C | A, W$ and therefore $S_0(t, w)$ and $G_0(t, w)$ have the following product formula representations:

$$S_0(t, w) = \prod_{m=1}^t \{1 - h_0(m, w)\}; \quad G_0(t, w) = \prod_{m=0}^{t-1} \{1 - g_{R,0}(m, w)\}, \quad (4)$$

which leads to the following identification result:

$$\theta_0 = E_0[S_0(\tau, W)] = E_0 \left[\prod_{m=1}^{\tau} \{1 - h_0(m, W)\} \right].$$

We sometimes use the notation $\theta(P)$ to refer to the above parameter evaluated an any arbitrary distribution P of O in the non-parametric model.

For an estimator $\hat{\theta}$ of θ_0 , we refer to $n^{1/2}$ -consistency as the property that $n^{1/2}(\hat{\theta} - \theta_0)$ is bounded in probability. We describe the estimator as *asymptotic linear* if it admits the representation $n^{1/2}(\hat{\theta} - \theta_0) = \frac{1}{\sqrt{n}} \sum_i D(O_i) + o_P(1)$ for some function D . Asymptotically linear estimators are also referred to as *asymptotically normal* since the central limit theorem yields $n^{1/2}(\hat{\theta} - \theta_0) \rightsquigarrow N[0, \text{Var}\{D(O)\}]$. For an estimator $\hat{f}(t, o)$ of a parameter $f_0(t, o)$, and the $L_2(P_0)$ norm $\|f\|^2 = \sum_t \int f(t, o)^2 dP_0(o)$, we refer to $n^{1/4}$ -consistency as the property that $n^{1/4}\|\hat{f} - f_0\| = o_P(1)$. For a collection of functions (f_1, \dots, f_k) , the notation $\|(f_1, \dots, f_k)\|$ is used to denote the vector of element-wise norms.

3 Doubly robust consistency vs doubly robust inference

We start by presenting the efficient influence function for estimation of θ_0 in model the non-parametric model:

$$D_{\eta, \theta}(O) = - \sum_{t=1}^{\tau} \frac{\mathbf{1}(A=1)I_t}{g_A(W)G(t, W)} \frac{S(\tau, W)}{S(t, W)} \{L_t - h(t, W)\} + S(\tau, W) - \theta, \quad (5)$$

where we denote $\eta = (g, h)$ and $g = (g_A, g_R)$. This efficient influence function is a fundamental object for optimal estimation of θ_0 in the non-parametric model. First, for given estimators \hat{h} and \hat{g} , an estimator that solves for θ in the estimating equation $\mathbb{P}_n D_{\hat{\eta}, \theta} = 0$ is consistent if at least one of h_0 or g_0 is estimated consistently. Second, $\text{Var}\{D_{\eta_0, \theta_0}(O)\}$ is the efficiency bound for estimation of θ_0 in the model \mathcal{M} . Specifically, under consistent estimation of m_0 and g_0 at a fast enough rate (which we define below), an estimator that solves $\mathbb{P}_n D_{\hat{\eta}, \theta} = 0$ has variance smaller or equal to that of any regular, asymptotically linear estimator of θ_0 in the non-parametric model.

The estimator constructed by directly solving for θ in (the linear equation) $\mathbb{P}_n D_{\hat{\eta}, \theta} = 0$ is often referred to as the augmented IPW estimator, and we denote it by $\hat{\theta}_{\text{aipw}}$. The augmented IPW is sometimes problematic because directly solving the estimating equation can yield an estimate out of bounds of the parameter space [Gruber and van der Laan \(2010\)](#). Alternatives to repair the AIPW in cross-sectional analyses have been discussed by [Kang and Schafer \(2007\)](#); [Robins et al. \(2007\)](#); [Tan \(2010\)](#). In this paper we work under the targeted minimum loss based estimation (TMLE) framework of [van der Laan and Rubin \(2006\)](#); [van der Laan and Rose \(2011\)](#), which provides a general method to construct estimators that stay within the parameter space. In general, the TMLE of θ_0 is defined as a substitution estimator $\hat{\theta}_{\text{tmle}} = \theta(\tilde{P})$, where \tilde{P} is an estimate of P_0 constructed such that the corresponding $\tilde{\eta}$ and $\theta(\tilde{P})$ solve the estimating equation $\sum_{i=1}^n D_{\tilde{\eta}, \theta(\tilde{P})}(O_i) = o_P(n^{1/2})$. The estimator \tilde{P} is constructed by tilting an initial estimate \hat{P} towards a solution of the relevant estimating equation, by means of an empirical risk minimizer in a parametric submodel. The interested reader is referred to [Díaz et al. \(2018\)](#); [Moore and van der Laan \(2011\)](#) for more details on the construction of a TMLE for survival analysis. The preliminary estimator \hat{P} , or the component $\hat{\eta}$ necessary to evaluate $\theta(\hat{P})$, may be obtained based on data-adaptive regression methods. In this article we do not pursue the development of estimators of η_0 , but rather rely on estimators available in the literature. In particular, we advocate the use of stacked regression or learning ensembles, which poses desirable oracle guarantees [van der Laan et al. \(2007\)](#).

The analysis of the properties of the $\hat{\theta}_{\text{tmle}}$ estimator relies on (i) the fact that it solves the efficient influence function estimating equation, and (ii) the consistency and smoothness of the initial estimator $\hat{\eta}$. In particular, define the following conditions:

C1 (Doubly robust consistency of $\hat{\eta}$). Let $\|\cdot\|$ denote the $L_2(P_0)$ norm defined in the notation section. Assume $\|\hat{g}_A - g_{A,1}\| = o_P(1)$, $\|\hat{g}_R - g_{R,1}\| = o_P(1)$, and $\|\hat{h} - h_1\| = o_P(1)$, where either $(g_{A,1}, g_{R,1}) = (g_{A,0}, g_{R,0})$, or $h_1 = h_0$.

C2 (Donsker). Assume the class of functions $\{(g_A, g_R, h) : \|g_A - g_{A,1}\| < \delta, \|g_R - g_{R,1}\| < \delta, \|h - h_1\| < \delta\}$ is Donsker for some $\delta > 0$.

Under conditions **C1** and **C2**, an application of Theorems 5.9 and 5.31 of [van der Vaart \(1998\)](#) (see also example 2.10.10 in [van der Vaart and Wellner \(1996\)](#)) yields

$$\hat{\theta}_{\text{tmle}} - \theta_0 = \beta(\hat{\eta}) + (\mathbb{P}_n - P_0)D_{\eta_1, \theta_0} + o_P(n^{-1/2} + |\beta(\hat{\eta})|), \quad (6)$$

where $\beta(\hat{\eta}) = P_0 D_{\hat{\eta}, \theta_0}$. The term $(\mathbb{P}_n - P_0)D_{\eta_1, \theta_0}$ is an empirical average of mean zero i.i.d random variables, and thus converges to a normal random variable at $n^{1/2}$ -rate. Under **C1**,

$\beta(\hat{\eta})$ converges to zero in probability so that $\hat{\theta}_{\text{tmle}}$ is consistent. However, $n^{1/2}$ -consistency of $\hat{\theta}_{\text{tmle}}$ requires the stronger condition that $\beta(\hat{\eta}) = O_P(n^{-1/2})$. This can only be proved in general if *both* $(g_{A,1}, g_{R,1}) = (g_{A,0}, g_{R,0})$ and $h_1 = h_0$, in which case the stronger condition $\beta(\hat{\eta}) = o_P(n^{-1/2})$ holds. If $\hat{\eta}$ is estimated within a parametric model, the delta method yields asymptotic linearity of $\beta(\hat{\eta})$, which in turn yields asymptotic linearity of $\hat{\theta}_{\text{tmle}}$. However, in the doubly robust case of [C1](#) and under data-adaptive estimation of η_0 , an asymptotic analysis of this drift term is difficult, and the large sample distribution of the TMLE and AIPW is generally unknown. This means that typical doubly robust estimators are “doubly consistent”, but they cannot be used to obtain “doubly robust inference” such as confidence intervals that remain valid under inconsistent estimation of at most one nuisance parameter.

Our main achievement is to propose an estimation technique that Gaussianizes $\beta(\hat{\eta})$, i.e., it makes this term asymptotically linear. Gaussianizing drift terms such as $\beta(\hat{\eta})$ has been the subject of recent literature in targeted learning [van der Laan \(2014\)](#); [Benkeser et al. \(2017\)](#); [Díaz and van der Laan \(2017\)](#). These works develop estimation techniques for various problems in cross-sectional studies. In the next two sections we focus on the construction of drift-corrected estimators that endow the TMLE with a doubly robust asymptotic distribution through Gaussianization of the drift term $\beta(\hat{\eta})$. Extensions of cross-sectional techniques to the longitudinal setting are non-trivial, as they involve constructing asymptotic representations of $\beta(\hat{\eta})$ which are estimable. These representations depend on sequential conditional expectations of the efficient influence function, which for the longitudinal case involve carefully handling the at-risk sets for each time point. The alternative representation of $\beta(\hat{\eta})$ is achieved through representations in terms of score equations in the non-parametric model. Doubly robust inference is thus achieved through the construction estimators $\hat{\theta}$ that solve such score equations, thereby guaranteeing that $\beta(\hat{\eta})$ behaves as Gaussian variable asymptotically. The following remark provides an argument that Gaussianizing the drift term can also aid in reducing the bias of TMLE estimators when both nuisance estimators are inconsistent.

Remark 1 (Asymptotic bias of the TMLE under inconsistency of $\hat{\eta}$). *Assume $\hat{\eta}$ converges to some $\eta_1 \neq \eta_0$. Let θ_1 denote the solution to $P_0 D_{\eta_1, \theta} = 0$, and note that $D_{\eta_1, \theta_1} = D_{\eta_1, \theta_0} - \theta_1 + \theta_0$. Under [C2](#), an application of Theorem 5.31 of [van der Vaart \(1998\)](#) yields*

$$\hat{\theta}_{\text{tmle}} - \theta_1 = \beta(\hat{\eta}) + (\mathbb{P}_n - P_0)D_{\eta_1, \theta_1} + o_P(n^{-1/2} + |\beta(\hat{\eta})|).$$

Substituting $D_{\eta_1, \theta_1} = D_{\eta_1, \theta_0} - \theta_1 + \theta_0$ yields

$$\hat{\theta}_{\text{tmle}} - \theta_0 = \beta(\hat{\eta}) + (\mathbb{P}_n - P_0)D_{\eta_1, \theta_0} + o_P(n^{-1/2} + |\beta(\hat{\eta})|).$$

The empirical process term $(\mathbb{P}_n - P_0)D_{\eta_1, \theta_0}$ has mean zero. Thus, Gaussianizing $\beta(\hat{\eta})$ is expected to reduce the bias of $\hat{\theta}_{\text{tmle}}$ when $\hat{\eta}$ is doubly inconsistent.

4 Asymptotic representation of the drift term

Our proposed method to endow the TMLE with a doubly robust asymptotic distribution relies on an asymptotic representation of the drift term $\beta(\hat{\eta})$. This parameter is then estimated using targeted minimum loss based estimation. In [Theorem 1](#) below, we show that

this drift term may be written as a sum of score functions that depends on the true value of an additional nuisance parameter (defined below) and the estimator $\hat{\eta}$, plus a term that approaches zero at $n^{1/2}$ -rate. The insight that allows us to construct a TMLE with doubly robust asymptotic distribution is that Gaussianization of the drift term amounts to tilting the estimator $\hat{\eta}$ in a way such that it also targets a solution of these score equations, thereby estimating $\beta(\hat{\eta})$ at $n^{1/2}$ -rate. We introduce the following assumption regarding the convergence rate of $\hat{\eta}$ to η_1 :

C3 (Consistency rate for $\hat{\eta}$). Assume **C1**. In addition, assume that $\|\hat{g}_A - g_{A,1}\| = o_P(n^{-1/4})$, $\|\hat{g}_R - g_{R,1}\| = o_P(n^{-1/4})$, and $\|\hat{h} - h_1\| = o_P(n^{-1/4})$.

As discussed in the introduction, the above rate is achievable by many data-adaptive regression algorithms such as ℓ_1 regularization, tree-based methods, and neural networks. In particular, [Benkeser and van der Laan \(2016\)](#) show that a rate of $n^{-1/4-1/[8(d+1)]}$, where d is the dimension of W , is achievable under the mild assumption that the true regression function is right-hand continuous with left-hand limits and has variation norm bounded by a constant. Because it is generally not possible to know a-priori which regression algorithm will be more appropriate for a given problem, we propose to use an ensemble learner known as the super learner [van der Laan et al. \(2007\)](#). Super learning builds a combination of predictors in a user-given library of candidate estimators, where the weights minimize the cross-validated risk of the resulting combination. Super learner has been shown to have important theoretical guarantees [van der Laan & S. Dudoit & A.W. van der Vaart \(2006\)](#); [van der Vaart et al. \(2006\)](#) such as asymptotic equivalence to the oracle selector.

The following lemma provides a representation for the drift term in terms of score function in the tangent space of each of the models for $g_{A,0}$, $g_{R,0}$, and h_0 . Such approximation is achieved through the definition of the following univariate regression functions. For each time point t and k , define the time-dependent covariates $G_g(t, w) = g_{A,1}(w)G_1(t, w)$, $C_h(k, w) = S_1(\tau, w)/S_1(k, w)$, and define $M(w) = \sum_{t=1}^{\tau} S_1(t, w)$, where G_1 and S_1 denote the censoring and survival probabilities under the limits $g_{R,1}$ and h_1 of the estimators (**C1**). Define the following weighted error functions

$$\begin{aligned} e_{R,0}(k, w) &= E_0 \left[\frac{R_k - g_{R,1}(k, W)}{G_g(k+1, W)} \middle| J_k = 1, A = 1, C_h(k, W) = C_h(k, w) \right], \\ e_{L,0}(t, w) &= E_0 \left[C_h(t, W) \{L_t - h_1(t)\} \middle| I_t = 1, A = 1, G_g(t, W) = G_g(t, w) \right], \\ e_{A,0}(w) &= E_0 \left[\frac{A - g_{A,1}(W)}{g_{A,1}(W)} \middle| M(W) = M(w) \right]. \end{aligned} \quad (7)$$

For each k , define the conditional probabilities

$$\begin{aligned} d_{k,0}(t, w) &= P_0 [R_t = 1 \mid J_t = 1, A = 1, C_h(k, W) = C_h(k, w)], \\ b_{k,0}(t, w) &= P_0 [L_t = 1 \mid I_t = 1, A = 1, C_h(k, W) = C_h(k, w)], \\ u_{k,0}(t, w) &= P_0 [R_t = 1 \mid J_t = 1, A = 1, C_g(k, W) = C_g(k, w)], \\ v_{k,0}(t, w) &= P_0 [L_t = 1 \mid I_t = 1, A = 1, C_g(k, W) = C_g(k, w)], \\ q_0(w) &= P_0 [A = 1 \mid S_1(\tau, W) = S_1(\tau, w)], \end{aligned} \quad (8)$$

and the corresponding time-to-event functions

$$\begin{aligned} D_{k,0}(t) &= \prod_{m=0}^{t-1} \{1 - d_{k,0}(m)\}, & B_{k,0}(t) &= \prod_{m=1}^t \{1 - b_{k,0}(m)\}, \\ U_{k,0}(t) &= \prod_{m=0}^{t-1} \{1 - u_{k,0}(m)\}, & V_{k,0}(t) &= \prod_{m=1}^t \{1 - v_{k,0}(m)\}. \end{aligned}$$

We occasionally use the notation $\lambda_0 = (e_{R,0}, e_{L,0}, e_{A,0}, d_{k,0}, b_{k,0}, u_{k,0}, v_{k,0}, q_0)$ to refer to the collection of auxiliary nuisance parameters. We now present the asymptotic representation of the drift term.

Theorem 1 (Asymptotic approximation of the drift term). *Define the covariates*

$$\begin{aligned} H_A(w) &= \sum_{t=1}^{\tau} \frac{U_{t,0}(t, w)}{G_0(t, w)} \frac{V_{t,0}(t-1, w)}{g_{A,0}(w)} e_{L,0}(t, w), \\ H_R(k, w) &= \frac{1}{g_{A,0}(w) G_0(k+1, w)} \left\{ \sum_{t=k+1}^{\tau} \frac{V_{t,0}(t-1, w)}{V_{t,0}(k, w)} \frac{U_{t,0}(t, w)}{U_{t,0}(k, w)} \frac{G_0(k, w)}{G_0(t, w)} e_{L,0}(t, w) \right\}, \quad (9) \\ H_L(t, w) &= \frac{S_0(\tau, w)}{S_0(t, w)} \left\{ \sum_{k=0}^{t-1} \frac{S_0(t-1, w)}{S_0(k, w)} \frac{B_{k,0}(k, w)}{B_{k,0}(t-1, w)} \frac{D_{k,0}(k, w)}{D_{k,0}(t, w)} e_{R,0}(k, w) \right. \\ &\quad \left. + \frac{e_{A,0}(w)}{q_0(w) D_{t,0}(t, w)} \frac{S_0(t-1, w)}{B_{t,0}(t-1, w)} \right\} \end{aligned}$$

and define the following score functions:

$$\begin{aligned} D_{A,\hat{g}}(o) &= -H_A(w) \{a - \hat{g}_A(w)\}, \\ D_{R,\hat{g}}(o) &= - \sum_{k=0}^{\tau-1} a j_k H_R(k, w) \{r_k - \hat{g}_R(k, w)\}, \\ D_{L,\hat{h}}(o) &= - \sum_{t=1}^{\tau} a i_t H_L(t, w) \{l_t - \hat{h}(t, w)\}. \end{aligned}$$

Under **C3** we have $\beta(\hat{\eta}) = P_0\{D_{A,\hat{g}} + D_{R,\hat{g}} + D_{L,\hat{h}}\} + o_P(n^{-1/2})$.

The above approximation of the drift term depends only on λ_0 . Note that λ_0 depends on w only through one-dimensional transformations which are consistently estimable at $n^{1/4}$ -rate under **C3**. Thus, under condition **C3**, the parameters λ_0 can be estimated element-wise through non-parametric smoothing techniques. The asymptotic normality result that we present in Section 5 requires a consistency rate assumption for estimation of λ_0 . We now discuss two possible estimation techniques and introduce a rate assumption that will allow us to prove asymptotic normality.

The general method for estimating (7) and (8) proceeds by obtaining estimates of the covariates and outcomes, and then applying any non-parametric regression technique. For

instance, for a second-order kernel function K_h with bandwidth l the estimator of $b_{k,0}$ is given by

$$\hat{b}_k(t, w) = \frac{\sum_{i=1}^n I_{t,i} A_i K_l \{ \hat{C}_h(k, W_i) - \hat{C}_h(k, w) \} L_{t,i}}{\sum_{i=1}^n I_{t,i} A_i K_l \{ \hat{C}_h(k, W_i) - \hat{C}_h(k, w) \}}, \quad (10)$$

where $\hat{C}_h(k, w) = \hat{S}(\tau, w) / \hat{S}(k, w)$, and \hat{S} is constructed using the preliminary estimator \hat{h} and formula (4). The optimal bandwidth \hat{l}_{opt} may be chosen using K-fold cross-validation van der Vaart et al. (2006). The error functions may be estimated analogously by plugging in estimates $\hat{\eta}$ in all the quantities involved, and performing kernel smoothing. Alternatively, other non-parametric smoothing methods may be used for this purpose. For example, the highly adaptive lasso (HAL) Benkeser and van der Laan (2016) proceeds by constructing an alternative representation of the true function as a sum of basis functions that grows with the sample size, and then performing ℓ_1 regularization to select the appropriate basis functions.

The analysis of the drift-corrected estimators may be complicated due to the two-stage estimation process whereby the covariates C are estimated and then used in a univariate smoothing technique. We introduce the following assumption about the estimators of λ_0 , which helps us isolate this univariate non-parametric smoothing in (7) and (8) from the methods used to estimate the auxiliary covariates and outcomes in the same expressions.

C4 (Convergence rate for auxiliary nuisance parameter estimators). Let $\hat{\lambda}_0$ denote (7) and (8) with the auxiliary covariates G_g , C_h , and M replaced by estimates \hat{C}_g , \hat{C}_h , and \hat{M} . For example,

$$\hat{b}_{k,0} = P_0 \left[L_t = 1 \mid I_t = 1, A = 1, \hat{C}_h(k, W) = \hat{C}_h(k, w) \right].$$

Assume that the smoothing method used to obtain $\hat{\lambda}$ satisfies $\|\hat{\lambda} - \lambda_0\| = o_P(n^{-1/4})$.

Note that the above assumption is purely about the consistency of the smoothing method used to obtain $\hat{\lambda}$, because the covariates \hat{C}_h , \hat{C}_h , and \hat{M} are the same in $\hat{\lambda}$ and λ_0 . Non-parametric smoothing methods can be expected to satisfy this assumption in certain situations. For example, under the assumption that the map $c \mapsto P_0 \left[L_t = 1 \mid I_t = 1, A = 1, \hat{C}_h(k, W) = c \right]$ is twice differentiable, a kernel regression estimator with optimal bandwidth guarantees the desired convergence rate $\|\hat{b} - b_0\| = o_P(n^{-1/4})$. The HAL also achieves the desired rate under the weaker assumption that $c \mapsto P_0 \left[L_t = 1 \mid I_t = 1, A = 1, \hat{C}_h(k, W) = c \right]$ is càdlàg with bounded sectional variation norm Benkeser and van der Laan (2016). We make assumption C4 through the remainder of the manuscript.

Remark 2. A substitution estimator of the drift term may be constructed by plugging in all the nuisance estimates in the alternative representation given in Theorem 1. An intuitive solution to the doubly robust inference problem would then be to subtract this term from the $\hat{\theta}_{\text{tmle}}$. While this makes intuitive sense, it does not guarantee that the resulting estimator will have the desired properties. The reason is that this strategy fails to control the term $|\beta(\hat{\eta})|$ that shows up in the $o_P(\cdot)$ expression in (6). The authors of van der Laan (2014); Benkeser et al. (2017); Díaz and van der Laan (2017) also noticed this problem in the cross-sectional setting, a more in-depth explanation of the issue may be found in these references.

5 TMLE with doubly robust inference

We now proceed to present an estimation method to Gaussianize the drift term $\beta(\hat{\eta})$. As discussed in the previous section, it is necessary to construct estimators $\hat{\eta}$ such that $\beta(\hat{\eta})$ is asymptotically Gaussian. Lemma 1 in the Supplementary Materials shows that, for any $\hat{\eta}$,

$$\beta(\hat{\eta}) - \hat{\beta}(\hat{\eta}) = -(\mathbb{P}_n - P_0)\{D_{R,\hat{g}} + D_{A,\hat{g}} + D_{L,\hat{h}}\} + o_P(n^{-1/2}), \quad (11)$$

where $\hat{\beta}(\hat{\eta})$ is constructed by plugging in estimates of $D_{R,\hat{g}}$, $D_{A,\hat{g}}$, and $D_{L,\hat{h}}$ in the result of Theorem 1. In light of expression (6), an asymptotically linear estimator $\hat{\eta}$ can be achieved through the construction of an estimator $\tilde{\eta}$ that satisfies $\hat{\beta}(\tilde{\eta}) = 0$. In the following, this construction is based on the fact that $D_{R,\hat{g}}$, $D_{A,\hat{g}}$, and $D_{L,\hat{h}}$ are score equations in the model for $g_{R,0}$, $g_{A,0}$, and h_0 , respectively. As a result, adding the corresponding H covariates to a logistic tilting model will tilt an initial estimator $\hat{\eta} = (\hat{g}_A, \hat{g}_R, \hat{h})$ towards a solution $\tilde{\eta}$ of the Gaussianizing equations $\hat{\beta}(\tilde{\eta}) = 0$. Our method for solving this estimating equations is rooted in the ideas of targeted learning van der Laan and Rose (2011). Readers familiar with targeted learning will see the similarities between the iterative procedure below and the estimators presented, e.g., in Moore and van der Laan (2011). As in Moore and van der Laan (2011), use the framework of targeted learning to solve the relevant estimating equations. Unlike Moore and van der Laan (2011), here we are not only interested in solving the efficient influence function estimating equation, but also in simultaneously solving the Gaussianizing equation $\hat{\beta}(\tilde{\eta}) = 0$. In what follows we will use the following modified representation of the data set:

$$\{(t, W_i, A_i, J_{t,i}, R_{t,i}, I_{t+1,i}, L_{t+1,i}) : t = 0, \dots, K-1; i = 1, \dots, n\}. \quad (12)$$

This data set is referred to as the long form, and the original data set

$$\{(W_i, A_i, \Delta_i, \tilde{T}_i) : i = 1, \dots, n\} \quad (13)$$

is referred to as the short form. The proposed targeted TMLE is defined by the following algorithm:

- Step 1. *Initial estimators.* Obtain initial estimators \hat{g}_A , \hat{g}_R , and \hat{h} of $g_{A,0}$, $g_{R,0}$, and h_0 . These estimators may be based on data-adaptive predictive methods that allow flexibility in the specification of the corresponding functional forms. Construct estimators \hat{e}_A , \hat{e}_R , \hat{e}_L by fitting a univariate regression method regression as described in the previous subsection. Similarly, for each k , compute estimators of $d_{k,0}$, $d_{k,0}$, $d_{k,0}$, $d_{k,0}$, and $d_{k,0}$ by also running univariate regressions.
- Step 2. *Compute auxiliary covariates.* For each subject i , compute the auxiliary covariates $\hat{H}_A(W_i)$, $\hat{H}_R(t, W_i)$, and $\hat{H}_L(t, W_i)$ by plugging in the estimators of the previous step in the definitions given in (9). In addition, compute the covariate

$$\hat{Z}(t, W_i) = \frac{\hat{S}(\tau, W_i)}{\hat{g}_A(W_i)\hat{S}(t, W_i)\hat{G}(t, W_i)}.$$

The covariate $\hat{Z}(t, W_i)$ is fundamental to obtain an estimator that solves the efficient influence function estimating equation (see Moore and van der Laan (2011)).

Step 3. *Solve estimating equations.* Estimate the parameter $\epsilon = (\epsilon_A, \epsilon_R, \epsilon_L)$ in the following logistic tilting models $g_{A,\epsilon}$, $g_{R,\epsilon}$, and h_ϵ for $g_{A,0}$, $g_{R,0}$, and h_0 :

$$\text{logit } g_{A,\epsilon}(W_i) = \text{logit } \hat{g}_A(W_i) + \epsilon_A \hat{H}_A(W_i) \quad (14)$$

$$\text{logit } g_{R,\epsilon}(t, W_i) = \text{logit } \hat{g}_R(t, W_i) + \epsilon_R \hat{H}_R(t, W_i). \quad (15)$$

$$\text{logit } h_\epsilon(t, W_i) = \text{logit } \hat{h}(t, W_i) + \epsilon_{L,1} \hat{H}_L(t, W_i) + \epsilon_{L,2} \hat{Z}(t, W_i). \quad (16)$$

where $\text{logit}(p) = \log\{p(1-p)^{-1}\}$. Here, $\text{logit } \hat{g}_R(t, w)$, $\text{logit } \hat{g}_A(w)$, and $\text{logit } \hat{h}(w)$ are offset variables (i.e., variables with known parameter equal to one). The above parameters may be estimated by fitting standard logistic regression models. For example, ϵ_R is estimated through a logistic regression model of R_t on $\hat{H}_R(t, W_i)$ with no intercept and an offset term equal to $\text{logit } \hat{g}_R(t, W)$ among observations with $(J_t, A) = (1, 1)$ in the long form dataset. Analogously, ϵ_A may be estimated by fitting a logistic regression model of A on \hat{H}_A with no intercept and an offset term equal to $\text{logit } \hat{g}_A(W)$ using all observations in the short form dataset, and ϵ_L is estimated through a logistic regression model of L_t on $(\hat{H}_L(t, W_i), \hat{Z}(t, W_i))$ with no intercept and an offset term equal to $\text{logit } \hat{h}(t, W)$ among observations with $(I_t, A) = (1, 1)$. Let $\hat{\epsilon}$ denote these estimates.

Step 4. *Update estimators and iterate.* Define the updated estimators as $\hat{g}_R = g_{R,\hat{\epsilon}}$, $\hat{g}_A = g_{A,\hat{\epsilon}}$, and $\hat{h} = h_{\hat{\epsilon}}$. Repeat steps 2-4 until convergence. In practice, we stop the iteration once $\max\{|\hat{\epsilon}_R|, |\hat{\epsilon}_A|, |\hat{\epsilon}_L|\} < 10^{-4}n^{-3/5}$.

Step 5. *Compute IPW.* Denote the estimators in the last step of the iteration with \tilde{g}_R , \tilde{g}_A , and \tilde{h} . The drift-corrected TMLE of θ_0 is defined as

$$\hat{\theta}_{\text{dr}} = \frac{1}{n} \sum_{i=1}^n \prod_{m=1}^{\tau} \{1 - \tilde{h}(m, W_i)\}.$$

The large sample distribution of the above TMLE is given in the following theorem:

Theorem 2 (Asymptotic Distribution of $\hat{\theta}_{\text{dr}}$). *Assume **C2** and **C3** hold for $\tilde{\eta}$, and **C4** holds for $\hat{\lambda}$. Then*

$$n^{1/2}(\hat{\theta}_{\text{dr}} - \theta_0) \rightarrow N(0, \sigma^2),$$

where $\sigma^2 = \text{Var}\{\text{IF}(O)\}$ and $\text{IF}(O) = D_{\eta_1, \theta_0}(O) - D_{L, h_1}(O) - D_{R, g_1}(O) - D_{A, g_1}(O)$. Furthermore:

(i) if $(g_{A,1}, g_{R,1}) = (g_{A,0}, g_{R,0})$ then $D_{R, g_1}(O) = D_{A, g_1}(O) = 0$, and

(ii) if $h_1 = h_0$, then $D_{L, h_1}(O) = 0$.

Thus, if $\eta_1 = \eta_0$ then $\text{IF} = D_{\eta_0, \theta_0}$ and $\hat{\theta}_{\text{dr}}$ is efficient.

The proof of this theorem is presented in the Supplementary Materials. Broadly, the proof proceeds as follows. First, inclusion of the covariate \hat{Z} guarantees that the submodel $\{h_\epsilon : \epsilon\}$ generates a score which is equal to the first term in the right hand side of (5). This is

used in the proof to show that the estimator solves the efficient influence function estimating equation and therefore satisfies (6). Then, we show that $\hat{\beta}(\tilde{g})$ is an asymptotically linear estimator of $\beta(\tilde{g})$ with influence function $D_{L,h_1}(O) + D_{R,g_1}(O) + D_{A,g_1}(O)$. Since $\hat{\beta}(\tilde{g}) = 0$, this asymptotic linearity result also implies $|\beta(\tilde{g})| = O_P(n^{-1/2})$, which according to the discussion in the previous section is a requisite for asymptotic linearity of $\hat{\theta}_{\text{dr}}$. An important part of this theorem is that, in the double consistency case in which $\eta_1 = \eta_0$, we have $D_{L,h_1}(O) = D_{R,g_1}(O) = D_{A,g_1}(O) = 0$, and the estimator $\hat{\theta}_{\text{dr}}$ is asymptotically equivalent to the $\hat{\theta}_{\text{tmle}}$, both being efficient. Unlike $\hat{\theta}_{\text{tmle}}$, the distribution of the estimator $\hat{\theta}_{\text{dr}}$ under condition C3 is known, and the variance given in the theorem can be used to compute doubly robust standard errors and to perform hypothesis tests. That is, the Wald-type confidence interval $\hat{\theta}_{\text{dr}} \pm z_\alpha \hat{\sigma}_{\text{dr}} / \sqrt{n}$, where $\hat{\sigma}_{\text{dr}}^2$ is the empirical variance of $\hat{\text{IF}}(O)$ has correct asymptotic coverage $(1 - \alpha)100\%$, whenever at least one of \tilde{g} or \tilde{h} converges to their true value at the stated rate.

5.1 Removing the Donsker Condition

Asymptotic linearity of $\hat{\theta}_{\text{dr}}$ requires Donsker condition C2. This is a powerful empirical processes condition that allows the analysis of many estimators in semi-parametric models van der Vaart (1998). However, this condition may be restrictive in high-dimensional settings, or when the estimator of the censoring mechanism is in a large class of function. For example functions classes with unbounded variation are generally not Donsker, and highly adaptive estimators such as random forests may have unbounded variation. Fortunately, C2 may be avoided by introducing cross-fitting into our estimation procedure. Cross-fitting was first proposed in the context of targeted minimum loss-based estimation in Zheng and van der Laan (2011), and was subsequently applied to estimating equations in Chernozhukov et al. (2016).

Our cross-fitting procedure proceeds as follows. Let $\mathcal{V}_1, \dots, \mathcal{V}_J$ denote a random partition of the index set $\{1, \dots, n\}$ into J validation sets of approximately the same size. That is, $\mathcal{V}_j \subset \{1, \dots, n\}$; $\bigcup_{j=1}^J \mathcal{V}_j = \{1, \dots, n\}$; and $\mathcal{V}_j \cap \mathcal{V}_{j'} = \emptyset$. In addition, for each j , the associated training sample is given by $\mathcal{T}_j = \{1, \dots, n\} \setminus \mathcal{V}_j$. Denote by $\hat{\eta}_{\mathcal{T}_j}$ the estimator of η_0 obtained by training the corresponding prediction algorithms using only data in the sample \mathcal{T}_j . Let also $j(i)$ denote the index of the validation set which contains observation i . The cross-fitted TMLE estimator is constructed replacing $\hat{\eta}$ by its cross-fitted in steps 2 and 3 of the first iteration of the TMLE algorithm described in Section 5. Let $\hat{\theta}_{\text{cfdR}}$ denote the resulting estimator. We have the following theorem.

Theorem 3 (Asymptotic Distribution of $\hat{\theta}_{\text{cfdR}}$). *Assume C3 holds for $\tilde{\eta}$ and C4 holds for $\hat{\lambda}$. Then $n^{1/2}(\hat{\theta}_{\text{cfdR}} - \theta_0) \rightarrow N(0, \sigma^2)$, where σ^2 is defined as in Theorem 2.*

The proof of this theorem is a straightforward adaptation of the proofs in Zheng and van der Laan (2011) to our Theorem 2. The proof rests on the key observation that for each validation set \mathcal{V}_j , the estimators $\hat{g}_{A,\mathcal{T}_j}$ and $\hat{g}_{R,\mathcal{T}_j}$ are fixed functions, and thus no entropy conditions are required in the application of empirical process results. The interested reader is encouraged to consult the original articles Zheng and van der Laan (2011); Chernozhukov et al. (2016) for more details and general proofs on cross-fitting.

6 Numerical study

In this section we present the results of a simulation experiment to illustrate the finite sample performance of statistical inference based on the asymptotic distribution given in Theorem 2.

We evaluate our method using a covariate vector of dimension $d = 10$, where the data generating mechanism for $g_{R,0}$, $g_{A,0}$, and h_0 is sparse; and ℓ_1 regularized logistic regression is used to estimate these nuisance parameters. This estimator satisfies condition C3 (see e.g., Theorem 4.1 of Rigollet et al. (2011)).

For each sample size $n \in \{400, 900, 1600, 2500, 3600, 4900\}$, we generate 1000 datasets from a conditional distribution defined as follows. First, a covariate vector W was generated from $TN(0, \Sigma)$ where TN is a multivariate normal distribution with each margin truncated at $(-1.5, 1.5)$, and Σ is a 10×10 symmetric Toeplitz matrix with first row equal to $(10, \dots, 1)/10$. We then define the unobserved variables

$$\begin{aligned} U_1 &= |W_1 W_2|^{1/2} - |W_{10}|^{1/2} + \cos(W_5) - \cos(W_6) \cos(W_5) \\ U_2 &= |W_1 W_{10}|^{1/2} - |W_9|^{1/2} + \cos(W_5) - \cos(W_7) \cos(W_6). \end{aligned}$$

The data are generated as

$$\begin{aligned} A \mid W = w &\sim \text{Ber}\{g_{A,0}(u_1)\} \\ R_t \mid J_t = 1, A = a, W = w &\sim \text{Ber}\{g_{R,0}(t, a, u_2)\} \\ L_t \mid I_t = 1, A = a, W = w &\sim \text{Ber}\{h_0(t, a, u_1)\}, \end{aligned}$$

where $\text{Ber}(p)$ denotes the Bernoulli distribution with parameter p and

$$\begin{aligned} g_{A,0}(u) &= \text{expit}(-2u) \\ g_{R,0}(t, a, u) &= \text{expit}\{-4 + a + a \cos(t) - aut^{1/2}\} \\ h_0(t, a, u) &= \text{expit}\{-3 + a - 2u \log(t) + 0.5au - 0.6(a+1)u \sin(t)\}, \end{aligned}$$

As previously discussed, $g_{R,0}$, $g_{A,0}$, and h_0 are estimated through ℓ_1 regularized logistic regression. For consistent estimation of $g_{A,0}$ the design matrix contains all W covariates in addition to their absolute squared root and cosine transformations as well as all two-way interactions between all these terms. For consistent estimation of $g_{R,0}$ and h_0 , the design matrix is constructed by considering all main effects and interactions of: (i) time as a categorical variable, (ii) the treatment indicator A , and (iii) all the terms considered for $g_{A,0}$. Inconsistent estimators were obtained through standard logistic regression with main terms only. We considered three scenarios for estimation of the nuisance parameters: (a) all $g_{A,0}$, $g_{R,0}$, and h_0 consistently estimated, (b) only h_0 consistently estimated, and (c) only $g_{A,0}$ and $g_{R,0}$ consistently estimated. We also performed a simulation where all nuisance parameters are inconsistently estimated, but the results are uninformative and are not presented.

We compute two estimators: a doubly robust $\hat{\theta}_{\text{tmle}}$ Moore and van der Laan (2011) and our proposed $\hat{\theta}_{\text{dr}}$. The $\hat{\theta}_{\text{tmle}}$ estimator has been shown to outperform the $\hat{\theta}_{\text{aipw}}$ estimator at finite samples in simulation studies Porter et al. (2011), and both are expected to have similar asymptotic behavior. We evaluate the performance of the estimators in terms of bias, variance, mean squared error, and coverage of the 90%, 95%, and 99% confidence intervals.

Some of these quantities are multiplied by $n^{1/2}$ to evaluate $n^{1/2}$ -consistency. We also evaluate $\hat{\sigma}_{\text{dr}}$ as an estimator of the standard error of the estimators.

The results are presented in Figure 1. Some expected properties of the estimators, which we corroborate in the simulation study are:

- The best behavior in terms of all metrics is obtained in scenario (a) for both estimators. In this case, both estimators have very similar asymptotic performance, with $\hat{\theta}_{\text{dr}}$ having slightly better bias in the smaller sample sizes.
- $\hat{\theta}_{\text{dr}}$ has significantly smaller bias than $\hat{\theta}_{\text{tmle}}$ for scenarios (b) and (c).
- The proposed estimator of the standard error $\hat{\sigma}_{\text{dr}}$ seems to consistently estimate the standard error of $\hat{\theta}_{\text{dr}}$ in all three scenarios, whereas the naïve estimator for $\hat{\theta}_{\text{tmle}}$ seems to be inconsistent in scenarios (b) and (c).
- The coverage probabilities for $\hat{\theta}_{\text{dr}}$ are closer to the nominal level for all sample sizes and all three scenarios. Of particular relevance, $\hat{\theta}_{\text{dr}}$ seems to provide very important small-sample gains in scenarios (b) and (c).

According to Remark 1, solving the debiasing equation $\beta(\hat{\eta}) = 0$ could reduce this bias of $\hat{\theta}_{\text{dr}}$, in comparison to the bias of $\hat{\theta}_{\text{tmle}}$ in the case of double inconsistency (results not shown). However, the MSE of both estimators was identical, and increased linearly in $n^{1/2}$ -scale. Identifying scenarios under which this bias reduction can be expected is an open problem.




Figure 1: Results of the numerical study. The scenarios under study are: (a) all $g_{A,0}$, $g_{R,0}$, and h_0 consistently estimated, (b) only h_0 consistently estimated, and (c) only $g_{A,0}$ and $g_{R,0}$ consistently estimated. $\text{Cov}(p)$ stands for coverage of a Wald-type 100p% confidence interval, and $\text{sd}(\hat{\theta})$ stands for the standard deviation of the estimator $\hat{\theta}$.

7 Motivating Application

Different types of human breast cancer tumors have been shown to have heterogeneous response to treatments [Perou et al. \(2000\)](#); [Sotiriou and Pusztai \(2009\)](#). Amplification of ERBB2 gene and associated overexpression of human epidermal growth factor receptor (HER2) encoded by this gene occur in 25-30% of breast cancers [Slamon et al. \(2001\)](#). HER2-positive breast cancer is an aggressive form of the disease and the prognosis for such patients is generally poor [Slamon \(1987\)](#); [Seshadri et al. \(1993\)](#). The clinical efficacy of adjuvant trastuzumab, a recombinant monoclonal antibody, in early stage HER2-positive patients was demonstrated by several large clinical trials [Perez et al. \(2011\)](#); [Romond et al. \(2005\)](#). We illustrate our methods using data for 1390 patients from the North Central Cancer Treatment Group N9831 study, a phase III randomized clinical trial testing the addition of trastuzumab to chemotherapy in stage I-III HER2-positive breast cancer. Recruitment started in 2000, and the maximum follow-up time was 16 years. The treatment group comprised 907 patients. The trial was subject to right censoring because patients drop out of the study and because enrollment spanned several years. We adjusted for 12 baseline variables which included demographic variables such as age, ethnicity, and race; as well as clinical variables such as tumor grade, nodal status, tumor size, and histology.

We estimated the treatment probabilities as well as the hazard of the event and censoring using an ensemble predictor known as the super learner [van der Laan et al. \(2007\)](#), implemented in the R package SuperLearner. Super learning builds a convex combination of candidate predictors in a user-given library, where the weights are chosen to minimize the cross-validated log-likelihood of the resulting ensemble. We present the results of the ensemble in Table 1, which includes some of the most popular statistical learning algorithms. The tuning parameters of each algorithm are chosen using internal cross-validation. In order to fully account for treatment-covariate interactions, we fitted separate models for the censoring probabilities in the treated and control arms.

Table 1: Super learning ensemble coefficients. RF: random forests, XGB: extreme gradient boosting, MLP: multi-layer perceptron, GLM: logistic regression, MARS: multivariate adaptive splines, Lasso: L_1 regularized logistic regression.

	RF	XGB	MLP	GLM	MARS	Lasso
g_A	0.00	0.47	0.15	0.00	0.09	0.28
$g_R, A = 1$	0.00	0.29	0.00	0.00	0.49	0.21
$g_R, A = 0$	0.00	0.41	0.00	0.00	0.09	0.50
$h, A = 1$	0.17	0.00	0.00	0.27	0.36	0.20
$h, A = 0$	0.27	0.13	0.00	0.00	0.42	0.18

We computed the $\hat{\theta}_{\text{dr}}$ and $\hat{\theta}_{\text{tmle}}$ estimators separately for the treated and untreated groups at time $\tau = 12$ years. We obtained an estimated difference ($A = 1$ vs $A = 0$) in survival probability of $\hat{\theta}_{\text{dr}} = 0.107$ (s.e. 0.036) and 0.098 (s.e. 0.032) years in the treatment arm for each estimator, respectively. The Kaplan-Meier estimator is equal to 0.044 (s.e. 0.032), highlighting the possible bias due to informative censoring.

8 Discussion

Our method assumes that censoring is confounded with the time to event only by baseline variables. In the presence of time dependent confounding between censoring and the event time, our proposal may be adapted by augmenting the censoring and outcome models to include time-varying confounders. In these settings, it may be possible to retain the asymptotic linearity to proper control of the drift term by generalizing the techniques used to prove our theorems. Such generalizations are unfortunately not trivial because the representations of the drift term and therefore its targeting algorithm varies with the estimating equation considered.

Our methods for doubly robust estimators trivially translate into $n^{1/2}$ -consistency rates for inverse probability weighted estimators under $n^{1/4}$ -consistent estimation of the g components of the nuisance parameter η . Specifically, such IPW estimator may be obtained through our proposal by setting $\hat{h}(t, a, w) = 1$, and omitting the tilting model (16) in the iterative procedure that defines the proposed estimator.

Most clinical research studies use discrete time scales to measure the time to event. This is the case of our application and simulation studies. If time is measured on a continuous scale, implementation of our methods requires discretization. The specific choice of the discretization intervals may be guided by what is clinically relevant. For example, in clinical applications with time to death outcomes, the clinically relevant scale would typically be a day. In the absence of clinical criteria to guide the choice of discretization level, a concern is that too coarse of a discretization may lead to potentially meaningful information losses. A question for future research is how to optimally set the level of discretization in order to trade off information loss versus estimator precision. Another area for future research is to consider discretization levels that get finer with sample size.

Existing doubly robust estimators cannot be proved regular or $n^{1/2}$ -consistent in general under inconsistent estimation of one of the nuisance parameters. While we do not tackle the regularity problem, we do solve the $n^{1/2}$ -consistency problem. This is done by proving a doubly robust asymptotic linearity result for our estimator, under the only assumption that at least one of the nuisance estimators is consistent at $n^{1/4}$ -rate. The regularity of our estimator remains an open problem along with that of all doubly robust estimators based on data-adaptive estimation of nuisance parameters under inconsistency of at least one nuisance estimator.

The $n^{1/4}$ -rate required by our estimators may still be considered a restrictive assumption. However, this rate is achievable by several data-adaptive regression algorithms under certain assumptions on the true regression functions. See for example [Bickel et al. \(2009\)](#) for results on ℓ_1 regularization, [Wager and Walther \(2015\)](#) for results on regression trees, and [Chen and White \(1999\)](#) for neural networks. This convergence rate is also achievable by the highly adaptive lasso [Benkeser and van der Laan \(2016\)](#) under the mild assumption that the true regression function is right-hand continuous with left-hand limits and has variation norm bounded by a constant.

9 Software

Software in the form of R code, together with a sample input data set and complete documentation is available at <https://github.com/idiazst/survdr>.

Supplementary Material

Theorem 1

Proof. For notational simplicity, in this proof we omit the dependence of all functions on w . E.g., $h_0(t, w)$ is denoted with $h_0(t)$. Lemma 1 in the Supplementary materials of [Díaz et al. \(2018\)](#) shows

$$P_0 D_{\hat{\eta}, \theta_0} = \sum_{t=1}^{\tau} E_0 \left[-\frac{\hat{S}(\tau)}{\hat{g}_A \hat{S}(t) \hat{G}(t)} S_0(t-1) \{h_0(t) - \hat{h}(t)\} \left\{ g_{A,0} G_0(t) - \hat{g}_A \hat{G}(t) \right\} \right].$$

First, note that

$$\begin{aligned} \{h_0(t) - \hat{h}(t)\} \{g_{A,0} G_0(t) - \hat{g}_A \hat{G}(t)\} &= \{h_0(t) - h_1(t)\} \{g_{A,0} G_0(t) - \hat{g}_A \hat{G}(t)\} \\ &\quad + \{h_0(t) - \hat{h}(t)\} \{g_{A,0} G_0(t) - g_{A,1} G_1(t)\} \\ &\quad + \{h_0(t) - h_1(t)\} \{g_{A,0} G_0(t) - g_{A,1} G_1(t)\} \end{aligned} \quad (17)$$

$$+ \{h_1(t) - \hat{h}(t)\} \{g_{A,1} G_1(t) - \hat{g}_A \hat{G}(t)\}. \quad (18)$$

By assumption, the expectation of (17) with respect to P_0 is zero, and the expectation of (18) is $o_P(n^{-1/2})$. Define

$$\begin{aligned} \beta_g(\hat{g}) &= \sum_{t=1}^{\tau} E_0 \left[-\frac{\hat{S}(\tau)}{\hat{g}_A \hat{S}(t) \hat{G}(t)} S_0(t-1) \{h_0(t) - h_1(t)\} \left\{ g_{A,0} G_0(t) - \hat{g}_A \hat{G}(t) \right\} \right], \\ \beta_h(\hat{h}) &= \sum_{t=1}^{\tau} E_0 \left[-\frac{\hat{S}(\tau)}{\hat{g}_A \hat{S}(t) \hat{G}(t)} S_0(t-1) \{h_0(t) - \hat{h}(t)\} \left\{ g_{A,0} G_0(t) - g_{A,1} G_1(t) \right\} \right] \end{aligned}$$

Assume first that $g_1 = g_0$. Denote

$$\begin{aligned} \hat{e}_{R,0}(k, w) &= E_0 \left[\frac{R_k - g_{R,1}(k, W)}{g_{A,1}(W) G_1(k+1, W)} \middle| J_k = 1, A = 1, \hat{C}_h(k, W) = \hat{C}_h(k, w) \right] \\ \hat{e}_{L,0}(t, w) &= E_0 \left[\frac{S_1(\tau)}{S_1(t)} \{L_t - h_1(t)\} \middle| I_t = 1, A = 1, \hat{C}_g(t, W) = \hat{C}_g(t, w) \right], \end{aligned}$$

where the expectation is taken with respect to P_0 taking \hat{C}_h and \hat{C}_g as fixed functions. We have

$$\begin{aligned}
\beta_g(\hat{g}) &= \sum_{t=1}^{\tau} E_0 \left[-\frac{S_1(\tau)}{S_1(t)} S_0(t-1) \{h_0(t) - h_1(t)\} \left\{ \frac{g_{A,0}}{\hat{g}_A} \frac{G_0(t)}{\hat{G}(t)} - 1 \right\} \right] + o_P(n^{-1/2}) \\
&= \sum_{t=1}^{\tau} E_0 \left[-\frac{S_1(\tau)}{S_1(t)} A I_t \{L_t - h_1(t)\} \left\{ \frac{1}{\hat{g}_A \hat{G}(t)} - \frac{1}{g_{A,0} G_0(t)} \right\} \right] + o_P(n^{-1/2}) \\
&= \sum_{t=1}^{\tau} E_0 \left[-A I_t e_{L,0}(t) \left\{ \frac{1}{\hat{g}_A \hat{G}(t)} - \frac{1}{g_{A,0} G_0(t)} \right\} \right] - \sum_{t=1}^{\tau} E_0 \left[\frac{A I_t}{\hat{g}_A \hat{G}(t)} \{ \hat{e}_{L,0}(t) - e_{L,0}(t) \} \right] + o_P(n^{-1/2}) \\
&= \sum_{t=1}^{\tau} E_0 \left[-A I_t e_{L,0}(t) \left\{ \frac{1}{\hat{g}_A \hat{G}(t)} - \frac{1}{g_{A,0} G_0(t)} \right\} \right] + o_P(n^{-1/2}) \\
&= \sum_{t=1}^{\tau} E_0 \left[-A V_{t,0}(t-1) U_{t,0}(t) e_{L,0}(t) \left\{ \frac{1}{\hat{g}_A \hat{G}(t)} - \frac{1}{g_{A,0} G_0(t)} \right\} \right] + o_P(n^{-1/2}) \\
&= \sum_{t=1}^{\tau} E_0 \left[-A V_{t,0}(t-1) U_{t,0}(t) \frac{e_{L,0}(t)}{g_{A,0} G_0(t)} \left\{ \frac{g_{A,0}}{\hat{g}_A \hat{G}(t)} \{G_0(t) - \hat{G}(t)\} + \frac{1}{\hat{g}_A} (g_{A,0} - \hat{g}_A) \right\} \right] + o_P(n^{-1/2}) \\
&= \sum_{t=1}^{\tau} E_0 \left[-A V_{t,0}(t-1) U_{t,0}(t) \frac{e_{L,0}(t)}{g_{A,0} G_0(t)} \left\{ \frac{g_{A,0}}{\hat{g}_A \hat{G}(t)} \{G_0(t) - \hat{G}(t)\} \right\} \right] \tag{19}
\end{aligned}$$

$$- \sum_{t=1}^{\tau} E_0 \left[V_{t,0}(t-1) U_{t,0}(t) \frac{e_{L,0}(t)}{G_0(t)} \frac{1}{g_{A,0}} (A - \hat{g}_A) \right] + o_P(n^{-1/2}), \tag{20}$$

where we get

$$\sum_{t=1}^{\tau} E_0 \left[\frac{A I_t}{\hat{g}_A \hat{G}(t)} \{ \hat{e}_{L,0}(t) - e_{L,0}(t) \} \right] = 0$$

using the law of iterated expectation. The term (20) is in the desired form. It remains to prove the result for (19). Define

$$\begin{aligned}
M_0(k, w) &= \{g_{R,0}(k, w) - \hat{g}_R(k, w)\} \frac{G_0(k, w)}{G_0(k+1, w)}, \\
\tilde{u}_{k,0}(t, w) &= P_0 [R_t = 1 \mid J_t = 1, A = 1, C_g(k, W) = C_g(k, w), M_0(k, W) = M_0(k, w)], \\
\tilde{v}_{k,0}(t, w) &= P_0 [L_t = 1 \mid I_t = 1, A = 1, C_g(k, W) = C_g(k, w), M_0(k, W) = M_0(k, w)],
\end{aligned}$$

and notice that $P_0\{\tilde{v}_{k,0}(t) - v_{k,0}(t)\} = O_P(\|\hat{g}_R - g_{R,0}\|)$, $P_0\{\tilde{u}_{k,0}(t) - u_{k,0}(t)\} = O_P(\|\hat{g}_R - g_{R,0}\|)$. Then (19) is equal to

$$\begin{aligned}
& \sum_{t=1}^{\tau} E_0 \left[-AV_{t,0}(t-1)U_{t,0}(t) \frac{e_{L,0}(t)}{g_{A,0}G_0(t)} \left\{ \frac{g_{A,0}}{\hat{g}_A \hat{G}(t)} \{G_0(t) - \hat{G}(t)\} \right\} \right] \\
&= \sum_{t=1}^{\tau} E_0 \left[-AV_{t,0}(t-1)U_{t,0}(t) \frac{e_{L,0}(t)}{g_{A,0}G_0(t)} \left\{ \frac{g_{A,0}}{\hat{g}_A \hat{G}(t)} \sum_{k=0}^{t-1} G_0(k) \{g_{R,0}(k) - \hat{g}_R(k)\} \frac{\hat{G}(t)}{\hat{G}(k+1)} \right\} \right] \\
&= \sum_{t=1}^{\tau} \sum_{k=0}^{t-1} E_0 \left[-\frac{A}{g_{A,0}} V_{t,0}(t-1)U_{t,0}(t) \frac{e_{L,0}(t)}{G_0(t)} \left\{ \{g_{R,0}(k) - \hat{g}_R(k)\} \frac{G_0(k)}{G_0(k+1)} \right\} \right] \\
&= \sum_{t=1}^{\tau} \sum_{k=0}^{t-1} E_0 \left[-\frac{A}{g_{A,0}} V_{t,0}(t-1)U_{t,0}(t) \frac{e_{L,0}(t)}{G_0(t)} \frac{1-R_0}{1-\tilde{u}_{t,0}(0)} \left\{ \{g_{R,0}(k) - \hat{g}_R(k)\} \frac{G_0(k)}{G_0(k+1)} \right\} \right] \\
&= \sum_{t=1}^{\tau} \sum_{k=0}^{t-1} E_0 \left[-\frac{A}{g_{A,0}} V_{t,0}(t-1)U_{t,0}(t) \frac{e_{L,0}(t)}{G_0(t)} \frac{1-R_0}{1-u_{t,0}(0)} \left\{ \{g_{R,0}(k) - \hat{g}_R(k)\} \frac{G_0(k)}{G_0(k+1)} \right\} \right] + O_P(\|\hat{g}_R - g_{R,0}\|^2) \\
&= \sum_{t=1}^{\tau} \sum_{k=0}^{t-1} E_0 \left[-\frac{A}{g_{A,0}} V_{t,0}(t-1)U_{t,0}(t) \frac{e_{L,0}(t)}{G_0(t)} \frac{1-R_0}{1-u_{t,0}(0)} \left\{ \{g_{R,0}(k) - \hat{g}_R(k)\} \frac{G_0(k)}{G_0(k+1)} \right\} \right] + o_P(n^{-1/2}) \\
&= \sum_{t=1}^{\tau} \sum_{k=0}^{t-1} E_0 \left[-\frac{A}{g_{A,0}} V_{t,0}(t-1)U_{t,0}(t) \frac{e_{L,0}(t)}{G_0(t)} \frac{1-R_0}{1-u_{t,0}(0)} \frac{1-L_1}{1-v_{t,0}(1)} \left\{ \{g_{R,0}(k) - \hat{g}_R(k)\} \frac{G_0(k)}{G_0(k+1)} \right\} \right] + o_P(n^{-1/2}) \\
&= \sum_{t=1}^{\tau} \sum_{k=0}^{t-1} E_0 \left[-\frac{A}{g_{A,0}} \frac{V_{t,0}(t-1)}{V_{t,0}(k)} \frac{U_{t,0}(t)}{U_{t,0}(k)} \frac{e_{L,0}(t)}{G_0(t)} J_k \left\{ \{g_{R,0}(k) - \hat{g}_R(k)\} \frac{G_0(k)}{G_0(k+1)} \right\} \right] + o_P(n^{-1/2}) \\
&= \sum_{t=1}^{\tau} \sum_{k=0}^{t-1} E_0 \left[-\frac{A}{g_{A,0}} \frac{V_{t,0}(t-1)}{V_{t,0}(k)} \frac{U_{t,0}(t)}{U_{t,0}(k)} \frac{e_{L,0}(t)}{G_0(t)} J_k \left\{ \{R_k - \hat{g}_R(k)\} \frac{G_0(k)}{G_0(k+1)} \right\} \right] + o_P(n^{-1/2}) \\
&= \sum_{k=0}^{\tau-1} \sum_{t=k+1}^{\tau} E_0 \left[-\frac{A}{g_{A,0}} \frac{V_{t,0}(t-1)}{V_{t,0}(k)} \frac{U_{t,0}(t)}{U_{t,0}(k)} \frac{e_{L,0}(t)}{G_0(t)} J_k \left\{ \{R_k - \hat{g}_R(k)\} \frac{G_0(k)}{G_0(k+1)} \right\} \right] + o_P(n^{-1/2}),
\end{aligned}$$

where the first equality follows from Lemma 2.

Assume now $h_1 = h_0$. Then

$$\beta_h(\hat{h}) = \sum_{t=1}^{\tau} E_0 \left[-\frac{\hat{S}(\tau)}{\hat{S}(t)} S_0(t-1) \{h_0(t) - \hat{h}(t)\} \left\{ \frac{g_{A,0}}{g_{A,1}} \sum_{k=0}^{t-1} \{g_{R,0}(k) - g_{R,1}(k)\} \frac{G_0(k)}{G_1(k+1)} \right\} \right] \quad (21)$$

$$+ \sum_{t=1}^{\tau} E_0 \left[-\frac{\hat{S}(\tau)}{\hat{S}(t)} S_0(t-1) \{h_0(t) - \hat{h}(t)\} \frac{1}{g_{A,1}} (g_{A,0} - g_{A,1}) \right]. \quad (22)$$

We first tackle the term in (21). This term is equal to

$$E_0 \left[\frac{g_{A,0}}{g_{A,1}} \sum_{k=0}^{\tau-1} \{g_{R,0}(k) - g_{R,1}(k)\} \frac{G_0(k)}{G_1(k+1)} \sum_{t=k+1}^{\tau} \frac{\hat{S}(\tau)}{\hat{S}(t)} S_0(t-1) \{\hat{h}(t) - h_0(t)\} \right].$$

We have

$$\begin{aligned}
\sum_{t=k+1}^{\tau} \frac{\hat{S}(\tau)}{\hat{S}(t)} S_0(t-1) \{\hat{h}(t) - \hat{h}_0(t)\} &= \sum_{t=1}^{\tau} \frac{\hat{S}(\tau)}{\hat{S}(t)} S_0(t-1) \{\hat{h}(t) - h_0(t)\} - \sum_{t=1}^k \frac{\hat{S}(\tau)}{\hat{S}(t)} S_0(t-1) \{\hat{h}(t) - h_0(t)\} \\
&= S_0(\tau) - \hat{S}(\tau) - \frac{\hat{S}(\tau)}{\hat{S}(k)} \{S_0(k) - \hat{S}(k)\} \\
&= S_0(k) \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\},
\end{aligned}$$

where the second equality follows from Lemma 2. Thus, (21) equals

$$\begin{aligned}
& E_0 \left[\sum_{k=0}^{\tau-1} \left\{ \frac{g_{A,0}}{g_{A,1}} G_0(k) S_0(k) \{g_{R,0}(k) - g_{R,1}(k)\} \frac{1}{G_1(k+1)} \right\} \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] \\
&= E_0 \left[\sum_{k=0}^{\tau-1} \left\{ \frac{A J_k}{g_{A,1}} \{R_k - g_{R,1}(k)\} \frac{1}{G_1(k+1)} \right\} \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] \\
&= E_0 \left[\sum_{k=0}^{\tau-1} A J_k e_{R,0}(k) \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] + E_0 \left[\sum_{k=0}^{\tau-1} A J_k \frac{\hat{S}(\tau)}{\hat{S}(k)} \{\hat{e}_{R,0}(k) - e_{R,0}(k)\} \right] \\
&= E_0 \left[\sum_{k=0}^{\tau-1} A \mathbf{1}\{\bar{R}_{k-1} = 0, \bar{L}_k = 0\} e_{R,0}(k) \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] \\
&= E_0 \left[\sum_{k=0}^{\tau-1} A \mathbf{1}\{\bar{R}_{k-1} = 0, \bar{L}_{k-1} = 0\} (1 - L_k) e_{R,0}(k) \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] \\
&= E_0 \left[\sum_{k=0}^{\tau-1} A \mathbf{1}\{\bar{R}_{k-1} = 0, \bar{L}_{k-1} = 0\} \{1 - b_{k,0}(k)\} e_{R,0}(k) \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] \\
&= E_0 \left[\sum_{k=0}^{\tau-1} A \mathbf{1}\{\bar{R}_{k-2} = 0, \bar{L}_{k-1} = 0\} (1 - R_{k-1}) \{1 - b_{k,0}(k)\} e_{R,0}(k) \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] \\
&= E_0 \left[\sum_{k=0}^{\tau-1} A \mathbf{1}\{\bar{R}_{k-2} = 0, \bar{L}_{k-1} = 0\} \{1 - d_{k,0}(k-1)\} \{1 - b_{k,0}(k)\} e_{R,0}(k) \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] \\
&= E_0 \left[\sum_{k=0}^{\tau-1} A \mathbf{1}\{\bar{R}_{k-2} = 0, \bar{L}_{k-1} = 0\} \frac{D_{k,0}(k)}{D_{k,0}(k-1)} \frac{B_{k,0}(k)}{B_{k,0}(k-1)} e_{R,0}(k) \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] \\
&= E_0 \left[\sum_{k=0}^{\tau-1} A \mathbf{1}\{\bar{R}_{k-3} = 0, \bar{L}_{k-2} = 0\} \frac{D_{k,0}(k)}{D_{k,0}(k-2)} \frac{B_{k,0}(k)}{B_{k,0}(k-2)} e_{R,0}(k) \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] \\
&\vdots \\
&= E_0 \left[\sum_{k=0}^{\tau-1} A \frac{B_{k,0}(k)}{S_0(k)} D_{k,0}(k) e_{R,0}(k) S_0(k) \left\{ \frac{\hat{S}(\tau)}{\hat{S}(k)} - \frac{S_0(\tau)}{S_0(k)} \right\} \right] \\
&= E_0 \left[\sum_{t=1}^{\tau} -A \frac{\hat{S}(\tau)}{\hat{S}(t)} S_0(t-1) \{h_0(t) - \hat{h}(t)\} \left\{ \sum_{k=0}^{t-1} \frac{B_{k,0}(k)}{S_0(k)} D_{k,0}(k) e_{R,0}(k) \right\} \right] \\
&= E_0 \left[\sum_{t=1}^{\tau} \sum_{k=0}^{t-1} -A \frac{1 - R_0}{1 - d_{k,0}(0)} S_0(t-1) \{h_0(t) - \hat{h}(t)\} \frac{\hat{S}(\tau)}{\hat{S}(t)} \frac{B_{k,0}(k)}{S_0(k)} D_{k,0}(k) e_{R,0}(k) \right] \\
&= E_0 \left[\sum_{t=1}^{\tau} \sum_{k=0}^{t-1} -A \frac{1 - R_0}{1 - d_{k,0}(0)} \frac{1 - L_1}{1 - b_{k,0}(1)} S_0(t-1) \{h_0(t) - \hat{h}(t)\} \frac{\hat{S}(\tau)}{\hat{S}(t)} \frac{B_{k,0}(k)}{S_0(k)} D_{k,0}(k) e_{R,0}(k) \right] \\
&= E_0 \left[\sum_{t=1}^{\tau} \sum_{k=0}^{t-1} -A \frac{I_t}{D_{k,0}(t) B_{k,0}(t-1)} S_0(t-1) \{h_0(t) - \hat{h}(t)\} \frac{\hat{S}(\tau)}{\hat{S}(t)} \frac{B_{k,0}(k)}{S_0(k)} D_{k,0}(k) e_{R,0}(k) \right] \\
&= E_0 \left[\sum_{t=1}^{\tau} -A I_t \sum_{k=0}^{t-1} \left\{ \frac{S_0(t-1)}{B_{k,0}(t-1)} \frac{S_0(\tau)}{S_0(t)} \frac{B_{k,0}(k)}{S_0(k)} \frac{D_{k,0}(k)}{S_{k,0}(t)} e_{R,0}(k) \right\} \{L_t - \hat{h}(t)\} \right].
\end{aligned}$$

Similar arguments can be used to show that (22) equals

$$-E_0 \left[\frac{e_{A,0}}{q_0} A \sum_{t=1}^{\tau} \frac{S_0(\tau)}{S_0(t)} \frac{S_0(t-1)}{B_{t,0}(t-1)} \frac{I_t}{D_{t,0}(t)} \{L_t - \hat{h}(t)\} \right],$$

concluding the proof of the theorem. \square

Theorem 2

This result follows from (6) in the main document and Lemma 1 below.

Other results

Lemma 1. *Assume C2 and C3. Then we have*

$$\beta(\hat{\eta}) = -(\mathbb{P}_n - P_0)\{D_{A,g_1} + D_{R,g_1} + D_{L,h_1}\} + o_P(n^{-1/2}).$$

Proof. From Theorem 1 we have $\beta(\hat{\eta}) = P_0\{D_{A,\hat{g}} + D_{R,\hat{g}} + D_{L,\hat{h}}\} + o_P(n^{-1/2})$. We will show that

$$P_0 D_{R,\hat{g}} = -(\mathbb{P}_n - P_0) D_{R,g_1} + o_P(n^{-1/2}). \quad (23)$$

The proof for the other components of $\beta(\hat{\eta})$ follows analogous steps. Assume first that $g_1 = g_0$. Denote

$$\hat{D}_{R,\hat{g}}(o) = - \sum_{k=0}^{\tau-1} a j_k \hat{H}_R(k, w) \{r_k - \hat{g}_R(k, w)\}.$$

Note that, by construction, $\mathbb{P}_n \hat{D}_{R,\hat{g}} = 0$. Thus we have

$$P_0 D_{R,\hat{g}} = -(\mathbb{P}_n - P_0) \hat{D}_{R,\hat{g}} + P_0(D_{R,\hat{g}} - \hat{D}_{R,\hat{g}}),$$

where we added and subtracted $P_0 \hat{D}_{R,\hat{g}}$. We have

$$P_0(D_{R,\hat{g}} - \hat{D}_{R,\hat{g}}) = \int \sum_{k=0}^{\tau-1} a j_k \{\hat{H}_R(k, w) - H_R(k, w)\} \{g_{R,0}(k, w) - \hat{g}_R(k, w)\} dP_0(o).$$

The Cauchy-Schwartz inequality shows

$$P_0(D_{R,\hat{g}} - \hat{D}_{R,\hat{g}}) = O_P \left(\|\hat{g}_R - g_{R,0}\| \|\hat{H}_R - H_R\| \right).$$

We will now argue that $\|\hat{H}_R - H_R\|$ may be decomposed as a sum of two terms: one exclusively related to estimation of g_0 , and one exclusively related to the smoothing method used to obtain $\hat{\lambda}$. Lemma 2 along with the Cauchy-Schwartz inequality and the definition of H_R show that

$$\|\hat{H}_R - H_R\| = O_P(\|\hat{v}_k - v_{k,0}\| + \|\hat{u}_k - u_{k,0}\| + \|\hat{g}_R - g_{R,0}\| + \|\hat{e}_L - e_{L,0}\|)$$

Recall the definition of $\hat{v}_{k,0}$, $\hat{u}_{k,0}$, and $\hat{e}_{L,0}$ as the corresponding true expectations conditional on the estimated covariate \hat{C}_g . The triangle inequality shows

$$\|\hat{v}_k - v_{k,0}\| \leq \|\hat{v}_{k,0} - v_{k,0}\| + \|\hat{v}_k - \hat{v}_{k,0}\|,$$

where the first term in the right hand side converges as $\|\hat{C}_g - C_g\|$, and the second term is assumed $o_P(n^{-1/4})$ (C4). Analogous inequalities hold for $\hat{u}_{k,0}$ and \hat{e}_L . Since $\|\hat{C}_g - C_g\| = O_P(\|\hat{g}_R - g_{R,0}\| + \|\hat{g}_A - g_{A,0}\|)$, we get

$$P_0(D_{R,\hat{g}} - \hat{D}_{R,\hat{g}}) = O_P(\|\hat{g}_R - g_{R,0}\| \{ \|\hat{g}_R - g_{R,0}\| + \|\hat{g}_A - g_{A,0}\| + o_P(n^{-1/4}) \}).$$

Under condition C3 this term is $o_P(n^{-1/2})$. Under C2 and C3, example 2.10.10 of van der Vaart and Wellner (1996) yields that $\hat{D}_{R,\hat{g}}$ is in a Donsker class. Thus, according to theorem 19.24 of van der Vaart (1998):

$$P_0 D_{R,\hat{g}} = -(\mathbb{P}_n - P_0) D_{R,g_0} + o_P(n^{-1/2}).$$

If $h_1 = h_0$, then $e_L(t, w) = 0$, which implies $H_R(t, w) = 0$. Thus, $D_{R,\hat{g}}(o) = D_{R,g_1}(o) = 0$, and (23) follows trivially, concluding the proof of the lemma. \square

Lemma 2. For two sequences a_1, \dots, a_m and b_1, \dots, b_m such that $a_t \neq 1$ and $b_t \neq 1$, we have

$$\prod_{t=1}^m (1 - a_t) - \prod_{t=1}^m (1 - b_t) = \sum_{t=1}^m \left\{ \prod_{k=1}^{t-1} (1 - a_k) (b_t - a_t) \prod_{k=t+1}^m (1 - b_k) \right\}.$$

Proof. Replace $(b_t - a_t)$ by $(1 - a_t) - (1 - b_t)$ in the right hand side and expand the sum to notice it is a telescoping sum. \square

References

- Vahe Avagyan and Stijn Vansteelandt. Honest data-adaptive inference for the average treatment effect under model misspecification using penalised bias-reduced double-robust estimation. *arXiv preprint arXiv:1708.03787*, 2017.
- Alexandre Belloni, Victor Chernozhukov, and Christian Hansen. Inference on treatment effects after selection among high-dimensional controls. *The Review of Economic Studies*, 81(2):608–650, 2014. doi: 10.1093/restud/rdt044. URL <http://dx.doi.org/10.1093/restud/rdt044>.
- Alexandre Belloni, Victor Chernozhukov, Ivan Fernández-Val, and Christian Hansen. Program evaluation and causal inference with high-dimensional data. *Econometrica*, 85(1): 233–298, 2017.
- O. Bembom, J.W. Fessel, R.W. Shafer, and M.J. van der Laan. Data-adaptive selection of the adjustment set in variable importance estimation. 2008. URL <http://www.bepress.com/ucbbiostat/paper231>.

- David Benkeser and Mark van der Laan. The highly adaptive lasso estimator. In *2016 IEEE International Conference on Data Science and Advanced Analytics (DSAA)*, pages 689–696. IEEE, 2016.
- David Benkeser, Marco Carone, MJ Van Der Laan, and PB Gilbert. Doubly robust non-parametric inference on the average treatment effect. *Biometrika*, 104(4):863–880, 2017.
- Peter J Bickel, Yaacov Ritov, Alexandre B Tsybakov, et al. Simultaneous analysis of lasso and dantzig selector. *The Annals of Statistics*, 37(4):1705–1732, 2009.
- Xiaohong Chen and Halbert White. Improved rates and asymptotic normality for non-parametric neural network estimators. *IEEE Transactions on Information Theory*, 45(2):682–691, 1999.
- Victor Chernozhukov, Denis Chetverikov, Mert Demirer, Esther Duflo, Christian Hansen, et al. Double machine learning for treatment and causal parameters. *arXiv preprint arXiv:1608.00060*, 2016.
- Stephen R Cole and Miguel A Hernán. Adjusted survival curves with inverse probability weights. *Computer methods and programs in biomedicine*, 75(1):45–49, 2004.
- Iván Díaz and Mark J van der Laan. Doubly robust inference for targeted minimum loss-based estimation in randomized trials with missing outcome data. *Statistics in medicine*, 36(24):3807–3819, 2017.
- Iván Díaz, Elizabeth Colantuoni, Daniel F. Hanley, and Michael Rosenblum. Improved precision in the analysis of randomized trials with survival outcomes, without assuming proportional hazards. *Lifetime Data Analysis*, Feb 2018. ISSN 1572-9249. doi: 10.1007/s10985-018-9428-5. URL <https://doi.org/10.1007/s10985-018-9428-5>.
- Iván Díaz, Oleksandr Savenkov, and Karla Ballman. Targeted learning ensembles for optimal individualized treatment rules with time-to-event outcomes. *Biometrika*, 105(3):723–738, 2018.
- Oliver Dukes, Vahe Avagyan, and Stijn Vansteelandt. High-dimensional doubly robust tests for regression parameters. *arXiv preprint arXiv:1805.06714*, 2018.
- Max H Farrell. Robust inference on average treatment effects with possibly more covariates than observations. *Journal of Econometrics*, 189(1):1–23, 2015.
- Susan Gruber and Mark J van der Laan. A targeted maximum likelihood estimator of a causal effect on a bounded continuous outcome. *The International Journal of Biostatistics*, 6(1), 2010.
- J. Kang and J. Schafer. Demystifying double robustness: A comparison of alternative strategies for estimating a population mean from incomplete data (with discussion). *Statistical Science*, 22:523–39, 2007.

- Brian K Lee, Justin Lessler, and Elizabeth A Stuart. Improving propensity score weighting using machine learning. *Statistics in medicine*, 29(3):337–346, 2010.
- Kelly L. Moore and Mark J. van der Laan. RCTs with time-to-event outcomes. In *Targeted Learning*, Springer Series in Statistics, pages 259–269. Springer New York, 2011. ISBN 978-1-4419-9781-4.
- Romain Neugebauer, Julie A Schmittdiel, and Mark J van der Laan. A case study of the impact of data-adaptive versus model-based estimation of the propensity scores on causal inferences from three inverse probability weighting estimators. *The international journal of biostatistics*, 12(1):131–155, 2016.
- Layla Parast, Lu Tian, and Tianxi Cai. Landmark estimation of survival and treatment effect in a randomized clinical trial. *Journal of the American Statistical Association*, 109(505):384–394, 2014.
- Edith A Perez, Edward H Romond, Vera J Suman, Jong-Hyeon Jeong, Nancy E Davidson, Charles E Geyer Jr, Silvana Martino, Eleftherios P Mamounas, Peter A Kaufman, and Norman Wolmark. Four-year follow-up of trastuzumab plus adjuvant chemotherapy for operable human epidermal growth factor receptor 2–positive breast cancer: Joint analysis of data from ncctg n9831 and nsabp b-31. *Journal of Clinical Oncology*, 29(25):3366–3373, 2011.
- Charles M Perou, Therese Sørlie, Michael B Eisen, Matt van de Rijn, Stefanie S Jeffrey, Christian A Rees, Jonathan R Pollack, Douglas T Ross, Hilde Johnsen, Lars A Akslen, et al. Molecular portraits of human breast tumours. *Nature*, 406(6797):747–752, 2000.
- Kristin E. Porter, Susan Gruber, Mark J. van der Laan, and Jasjeet S. Sekhon. The relative performance of targeted maximum likelihood estimators. *The International Journal of Biostatistics*, 7(1):1–34, 2011.
- Greg Ridgeway and Daniel F. McCaffrey. Comment: Demystifying double robustness: A comparison of alternative strategies for estimating a population mean from incomplete data. *Statist. Sci.*, 22(4):540–543, 11 2007. doi: 10.1214/07-STS227C. URL <http://dx.doi.org/10.1214/07-STS227C>.
- Philippe Rigollet, Alexandre Tsybakov, et al. Exponential screening and optimal rates of sparse estimation. *The Annals of Statistics*, 39(2):731–771, 2011.
- James Robins, Mariela Sued, Quanhong Lei-Gomez, and Andrea Rotnitzky. Comment: Performance of double-robust estimators when” inverse probability” weights are highly variable. *Statistical Science*, 22(4):544–559, 2007.
- Edward H Romond, Edith A Perez, John Bryant, Vera J Suman, Charles E Geyer Jr, Nancy E Davidson, Elizabeth Tan-Chiu, Silvana Martino, Soonmyung Paik, Peter A Kaufman, et al. Trastuzumab plus adjuvant chemotherapy for operable her2-positive breast cancer. *New England Journal of Medicine*, 353(16):1673–1684, 2005.

- Andrea Rotnitzky and James M Robins. Inverse probability weighting in survival analysis. *Encyclopedia of Biostatistics*, 2005.
- Donald B Rubin. *Multiple Imputation for Nonresponse in Surveys*. John Wiley & Sons, 1987.
- Ram Seshadri, FA Firgaira, DJ Horsfall, K McCaul, V Setlur, and P Kitchen. Clinical significance of her-2/neu oncogene amplification in primary breast cancer. the south australian breast cancer study group. *Journal of Clinical Oncology*, 11(10):1936–1942, 1993.
- Dennis J Slamon, Brian Leyland-Jones, Steven Shak, Hank Fuchs, Virginia Paton, Alex Bajamonde, Thomas Fleming, Wolfgang Eiermann, Janet Wolter, Mark Pegram, et al. Use of chemotherapy plus a monoclonal antibody against her2 for metastatic breast cancer that overexpresses her2. *New England Journal of Medicine*, 344(11):783–792, 2001.
- DJ Slamon. Human breast cancer: correlation of relapse and. *Science*, 3798106(177):235, 1987.
- Christos Sotiriou and Lajos Pusztai. Gene-expression signatures in breast cancer. *New England Journal of Medicine*, 360(8):790–800, 2009.
- Zhiqiang Tan. Bounded, efficient and doubly robust estimation with inverse weighting. *Biometrika*, 97(3):661–682, 2010.
- Mark J van der Laan. Targeted estimation of nuisance parameters to obtain valid statistical inference. *The international journal of biostatistics*, 10(1):29–57, 2014.
- Mark J van der Laan and Richard JCM Starmans. Entering the era of data science: Targeted learning and the integration of statistics and computational data analysis. *Advances in Statistics*, 2014, 2014.
- M.J. van der Laan and S. Rose. *Targeted Learning: Causal Inference for Observational and Experimental Data*. Springer, New York, 2011.
- M.J. van der Laan and D. Rubin. Targeted maximum likelihood learning. *The International Journal of Biostatistics*, 2(1):Article 11, 2006.
- M.J. van der Laan, M.L. Petersen, and M.M. Joffe. History-adjusted marginal structural models & statically-optimal dynamic treatment regimens. *The International Journal of Biostatistics*, 1(1):10–20, 2005.
- M.J. van der Laan, E. Polley, and A. Hubbard. Super learner. *Statistical Applications in Genetics & Molecular Biology*, 6(25):Article 25, 2007.
- M.J. van der Laan & S. Dudoit & A.W. van der Vaart. The cross-validated adaptive epsilon-net estimator. *Statistics & Decisions*, 24(3):373–395, 2006.
- Y. Wang & O. Bembom & M.J. van der Laan. Data adaptive estimation of the treatment specific mean. *Journal of Statistical Planning & Inference*, 2006.

- A. W. van der Vaart. *Asymptotic Statistics*. Cambridge University Press, 1998.
- A. W. van der Vaart and J. A. Wellner. *Weak Convergence and Empirical Processes*. Springer-Verlag New York, 1996.
- A.W. van der Vaart, S. Dudoit, and M.J. van der Laan. Oracle inequalities for multi-fold cross-validation. *Statistics & Decisions*, 24(3):351–371, 2006.
- Stefan Wager and Guenther Walther. Adaptive concentration of regression trees, with application to random forests. *arXiv preprint arXiv:1503.06388*, 2015.
- Jun Xie and Chaofeng Liu. Adjusted kaplan–meier estimator and log-rank test with inverse probability of treatment weighting for survival data. *Statistics in medicine*, 24(20):3089–3110, 2005.
- Min Zhang. Robust methods to improve efficiency and reduce bias in estimating survival curves in randomized clinical trials. *Lifetime data analysis*, pages 1–19, 2014.
- Wenjing Zheng and Mark J van der Laan. Cross-validated targeted minimum-loss-based estimation. In *Targeted Learning*, pages 459–474. Springer, 2011.